

Part II — Algebraic Topology

Based on lectures by H. Wilton

Notes taken by Dexter Chua

Michaelmas 2015

These notes are not endorsed by the lecturers, and I have modified them (often significantly) after lectures. They are nowhere near accurate representations of what was actually lectured, and in particular, all errors are almost surely mine.

Part IB Analysis II is essential, and Metric and Topological Spaces is highly desirable

The fundamental group

Homotopy of continuous functions and homotopy equivalence between topological spaces. The fundamental group of a space, homomorphisms induced by maps of spaces, change of base point, invariance under homotopy equivalence. [3]

Covering spaces

Covering spaces and covering maps. Path-lifting and homotopy-lifting properties, and their application to the calculation of fundamental groups. The fundamental group of the circle; topological proof of the fundamental theorem of algebra. *Construction of the universal covering of a path-connected, locally simply connected space*. The correspondence between connected coverings of X and conjugacy classes of subgroups of the fundamental group of X . [5]

The Seifert-Van Kampen theorem

Free groups, generators and relations for groups, free products with amalgamation. Statement *and proof* of the Seifert-Van Kampen theorem. Applications to the calculation of fundamental groups. [4]

Simplicial complexes

Finite simplicial complexes and subdivisions; the simplicial approximation theorem. [3]

Homology

Simplicial homology, the homology groups of a simplex and its boundary. Functorial properties for simplicial maps. *Proof of functoriality for continuous maps, and of homotopy invariance*. [4]

Homology calculations

The homology groups of S^n , applications including Brouwer's fixed-point theorem. The Mayer-Vietoris theorem. *Sketch of the classification of closed combinatorial surfaces*; determination of their homology groups. Rational homology groups; the Euler-Poincaré characteristic and the Lefschetz fixed-point theorem. [5]

Contents

0	Introduction	3
1	Definitions	4
1.1	Some recollections and conventions	4
1.2	Cell complexes	4
2	Homotopy and the fundamental group	7
2.0	Motivation	7
2.1	Homotopy	7
2.2	Paths	11
2.3	The fundamental group	15
3	Covering spaces	22
3.1	Covering space	22
3.2	The fundamental group of the circle and its applications	30
3.3	Universal covers	32
3.4	The Galois correspondence	35
4	Some group theory	39
4.1	Free groups and presentations	39
4.2	Another view of free groups	41
4.3	Free products with amalgamation	44
5	Seifert-van Kampen theorem	46
5.1	Seifert-van Kampen theorem	46
5.2	The effect on π_1 of attaching cells	50
5.3	A refinement of the Seifert-van Kampen theorem	52
5.4	The fundamental group of all surfaces	54
6	Simplicial complexes	56
6.1	Simplicial complexes	56
6.2	Simplicial approximation	62
7	Simplicial homology	66
7.1	Simplicial homology	66
7.2	Some homological algebra	71
7.3	Homology calculations	74
7.4	Mayer-Vietoris sequence	76
7.5	Continuous maps and homotopy invariance	82
7.6	Homology of spheres and applications	87
7.7	Homology of surfaces	89
7.8	Rational homology, Euler and Lefschetz numbers	92

0 Introduction

In topology, a typical problem is that we have two spaces X and Y , and we want to know if $X \cong Y$, i.e. if X and Y are homeomorphic. If they are homeomorphic, we can easily show this by writing down a homeomorphism. But what if they are not? How can we prove that two spaces are not homeomorphic?

For example, are \mathbb{R}^m and \mathbb{R}^n homeomorphic (for $m \neq n$)? Intuitively, they should not be, since they have different dimensions, and in fact they are not. But how can we actually prove this?

The idea of algebraic topology is to translate these non-existence problems in topology to non-existence problems in algebra. It turns out we are much better at algebra than topology. It is *much* easier to show that two groups are not isomorphic. For example, we will be able to reduce the problem of whether \mathbb{R}^m and \mathbb{R}^n are homeomorphic (for $m \neq n$) to the question of whether \mathbb{Z} and $\{e\}$ are isomorphic, which is very easy.

While the statement that $\mathbb{R}^m \not\cong \mathbb{R}^n$ for $n \neq m$ is intuitively obvious, algebraic topology can be used to prove some less obvious results.

Let D^n be the n dimensional unit disk, and S^{n-1} be the $n-1$ dimensional unit sphere. We will be able to show that there is no continuous map $F : D^n \rightarrow S^{n-1}$ such that the composition

$$S^{n-1} \hookrightarrow D^n \xrightarrow{F} S^{n-1}$$

is the identity, where the first arrow is the inclusion map. Alternatively, this says that we cannot continuously map the disk onto the boundary sphere such that the boundary sphere is fixed by the map.

Using algebraic topology, we can translate this statement into an algebraic statement: there is no homomorphism $F : \{0\} \rightarrow \mathbb{Z}$ such that

$$\mathbb{Z} \hookrightarrow \{0\} \xrightarrow{F} \mathbb{Z}$$

is the identity. This is something we can prove in 5 seconds.

By translating a non-existence problem of a continuous map to a non-existence problem of a homomorphism, we have made our life much easier.

In algebraic topology, we will be developing a lot of machinery to do this sort of translation. However, this machinery is not easy. It will take some hard work, and will be rather tedious and boring at the beginning. So keep in mind that the point of all that hard work is to prove all these interesting theorems.

In case you are completely uninterested in topology, and don't care if \mathbb{R}^m and \mathbb{R}^n are homeomorphic, further applications of algebraic topology include *solving equations*. For example, we will be able to prove the fundamental theorem of algebra (but we won't), as well as Brouwer's fixed point theorem (which says that every continuous function from $D^2 \rightarrow D^2$ has a fixed point). If you are not interested in these as well, you may as well drop this course.

1 Definitions

1.1 Some recollections and conventions

We will start with some preliminary definitions and conventions.

Definition (Map). In this course, the word *map* will always refer to continuous maps. We are doing topology, and never care about non-continuous functions.

We are going to build a lot of continuous maps in algebraic topology. To do so, we will often need to glue maps together. The *gluing lemma* tells us that this works.

Lemma (Gluing lemma). If $f : X \rightarrow Y$ is a function of topological spaces, $X = C \cup K$, C and K are both closed, then f is continuous if and only if the restrictions $f|_C$ and $f|_K$ are continuous.

Proof. Suppose f is continuous. Then for any closed $A \subseteq Y$, we have

$$f|_C^{-1}(A) = f^{-1}(A) \cap C,$$

which is closed. So $f|_C$ is continuous. Similarly, $f|_K$ is continuous.

If $f|_C$ and $f|_K$ are continuous, then for any closed $A \subseteq Y$, we have

$$f^{-1}(A) = f|_C^{-1}(A) \cup f|_K^{-1}(A),$$

which is closed. So f is continuous. □

This lemma is also true with “closed” replaced with “open”. The proof is the same proof with “closed” replaced with “open”.

We will also need the following technical lemma about metric spaces.

Lemma. Let (X, d) be a compact metric space. Let $\mathcal{U} = \{U_\alpha\}_{\alpha \in A}$ be an open cover of X . Then there is some δ such that for each $x \in X$, there is some $\alpha \in I$ such that $B_\delta(x) \subseteq U_\alpha$. We call δ a *Lebesgue number* of this cover.

Proof. Suppose not. Then for each $n \in \mathbb{N}$, there is some $x_n \in X$ such that $B_{1/n}(x_n)$ is not contained in any U_α . Since X is compact, the sequence (x_n) has a convergent subsequence. Suppose this subsequence converges to y .

Since \mathcal{U} is an open cover, there is some $\alpha \in A$ such that $y \in U_\alpha$. Since U_α is open, there is some $r > 0$ such that $B_r(y) \subseteq U_\alpha$. But then we can find a sufficiently large n such that $\frac{1}{n} < \frac{r}{2}$ and $d(x_n, y) < \frac{r}{2}$. But then

$$B_{1/n}(x_n) \subseteq B_r(y) \subseteq U_\alpha.$$

Contradiction. □

1.2 Cell complexes

In topology, we can construct some really horrible spaces. Even if we require them to be compact, Hausdorff etc, we can often still produce really ugly topological spaces with weird, unexpected behaviour. In algebraic topology, we will often restrict our attention to some nice topological spaces, known as *cell complexes*.


To build cell complexes, we are not just gluing maps, but spaces.

Definition (Cell attachment). For a space X , and a map $f : S^{n-1} \rightarrow X$, the space obtained by *attaching an n -cell* to X along f is

$$X \cup_f D^n = (X \amalg D^n) / \sim,$$


where the equivalence relation \sim is the equivalence relation generated by $x \sim f(x)$ for all $x \in S^{n-1} \subseteq D^n$ (and \amalg is the disjoint union).

Intuitively, a map $f : S^{n-1} \rightarrow X$ just picks out a subset X that looks like the sphere. So we are just sticking a disk onto X by attaching the boundary of the disk onto a sphere within X .



Definition (Cell complex). A (finite) *cell complex* is a space X obtained by


- (i) Start with a discrete finite set $X^{(0)}$.




- (ii) Given $X^{(n-1)}$, for $X^{(n)}$ by taking a finite set of maps $\{f_\alpha : S^{n-1} \rightarrow X^{(n-1)}\}$ by attaching n -cells along the f_α :

$$X^{(n)} = \left(X^{(n-1)} \amalg \coprod_{\alpha} D_{\alpha}^n \right) / \{x \sim f_{\alpha}(x)\}.$$

For example, given the $X^{(0)}$ above, we can attach some loops and lines to obtain the following $X^{(1)}$



We can add surfaces to obtain the following $X^{(2)}$




- (iii) Stop at some $X = X^{(k)}$. The minimum such k is the *dimension* of X .

To define non-finite cell complexes, we just have to remove the words “finite” in the definition and remove the final stopping condition.

We have just had an example of a cell complex above, and it is easy to produce many more examples, such as the n -sphere. So instead let’s look at a non-cell complex.

Example. The following is *not* a cell complex: we take \mathbb{R}^2 , and add a circle with radius $\frac{1}{2}$ and center $(0, \frac{1}{2})$. Then we add another circle with radius $\frac{1}{4}$ and center $(0, \frac{1}{4})$, then a circle with radius $\frac{1}{8}$ and center $(0, \frac{1}{8})$ etc. We obtain something like



This is known as the *Hawaiian Earring*.

Why is this not an (infinite) cell complex? We did obtain it by attaching lots of 1-cells to the single point $(0, 0)$. However, in the definition of a cell complex, the cells are supposed to be completely unrelated and disjoint, apart from intersecting at the origin. However, here the circles clump together at the origin.

In particular, if we take the following sequence $(0, 1), (0, \frac{1}{2}), (0, \frac{1}{4}), \dots$, it converges to $(0, 0)$. If this were a cell complex, then this shouldn’t happen because the cells are unrelated, and picking a point from each cell should not produce a convergent sequence (if you are not convinced, if we actually did produce by attaching cells, then note that during the attaching process, we needn’t have attached them this way. We could have made it such that the n th cell has radius n . Then clearly picking the topmost point of each cell will not produce a convergent sequence).

We will see that the Hawaiian Earring will be a counterexample to a lot of our theorems here.

2 Homotopy and the fundamental group

This will be our first trick to translate topological spaces into groups.

2.0 Motivation


Recall that we wanted to prove that $\mathbb{R}^n \not\cong \mathbb{R}^m$ for $n \neq m$. Let's first do the simple case, where $m = 1, n = 2$. We want to show that $\mathbb{R} \not\cong \mathbb{R}^2$.

This is not hard. We know that \mathbb{R} is a line, while \mathbb{R}^2 is a plane. Let's try to remove a point from each of them. If we remove a point from \mathbb{R} , the space stops being path connected. However, removing a point does not have this effect on \mathbb{R}^2 . Since being path connected is a topological property, we have now showed that \mathbb{R} and \mathbb{R}^2 are not homeomorphic.

Unfortunately, this does not extend very far. We cannot use this to show that \mathbb{R}^2 and \mathbb{R}^3 are not homeomorphic. What else can we do?

Notice that when we remove a point from \mathbb{R}^2 , sure it is still connected, but something has changed.

Consider a circle containing the origin in $\mathbb{R}^2 \setminus \{0\}$. If the origin were there, we can keep shrinking the circle down until it becomes a point. However, we cannot do this if the origin is missing.



The strategy now is to exploit the fact that $\mathbb{R}^2 \setminus \{0\}$ has circles which cannot be deformed to points.

2.1 Homotopy

We have just talked about the notion of “deforming” circles to a point. We can think of a circle in X as a map $S^1 \rightarrow X$, and we want to “deform” this map to a point. This process of deformation is known as *homotopy*. Here we are going to use the interval $[0, 1] \subseteq \mathbb{R}$ a lot, and we will just call it I .

Notation.

$$I = [0, 1] \subseteq \mathbb{R}.$$

Definition (Homotopy). Let $f, g : X \rightarrow Y$ be maps. A *homotopy* from f to g is a map

$$H : X \times I \rightarrow Y$$

such that

$$H(x, 0) = f(x), \quad H(x, 1) = g(x).$$

We think of the interval I as time. For each time t , $H(\cdot, t)$ defines a map $X \rightarrow Y$. So we want to start from f , move with time, and eventually reach g .

If such an H exists, we say f is *homotopic* to g , and write $f \simeq g$. If we want to make it explicit that the homotopy is H , we write $f \simeq_H g$.

As mentioned at the beginning, by calling H a map, we are requiring it to be continuous.

Sometimes, we don't want a general homotopy. We might want to make sure that when we are deforming from a path f to g , the end points of the path don't move. In general, we have

Definition (Homotopy rel A). We say f is *homotopic to g rel A* , written $f \simeq g \text{ rel } A$, if for all $a \in A \subseteq X$, we have

$$H(a, t) = f(a) = g(a).$$

This notion will find itself useful later, but we don't have to pay too much attention to this yet.

Our notation suggests that homotopy is an equivalence relation. Indeed, we have

Proposition. For spaces X, Y , and $A \subseteq X$, the “homotopic rel A ” relation is an equivalence relation. In particular, when $A = \emptyset$, homotopy is an equivalence relation.

Proof.

- (i) Reflexivity: $f \simeq f$ since $H(x, t) = f(x)$ is a homotopy.
- (ii) Symmetry: if $H(x, t)$ is a homotopy from f to g , then $H(x, 1 - t)$ is a homotopy from g to f .
- (iii) Transitivity: Suppose $f, g, h : X \rightarrow Y$ and $f \simeq_H g \text{ rel } A$, $g \simeq_{H'} h \text{ rel } A$. We want to show that $f \simeq h \text{ rel } A$. The idea is to “glue” the two maps together.

We know how to continuously deform f to g , and from g to h . So we just do these one after another. We define $H'' : X \times I \rightarrow Y$ by

$$H(x, t) = \begin{cases} H(x, 2t) & 0 \leq t \leq \frac{1}{2} \\ H'(x, 2t - 1) & \frac{1}{2} \leq t \leq 1 \end{cases}$$

This is well-defined since $H(x, 1) = g(x) = H'(x, 0)$. This is also continuous by the gluing lemma. It is easy to check that H'' is a homotopy rel A .

□

We now have a notion of equivalence of maps — two maps are equivalent if they are homotopic. We can extend the notion of homotopy to spaces as well.


Recall that when we defined homeomorphism, we required that there is some f, g such that $f \circ g = \text{id}$, $g \circ f = \text{id}$. Here, we replace equality by homotopy.

Definition (Homotopy equivalence). A map $f : X \rightarrow Y$ is a *homotopy equivalence* if there exists a $g : Y \rightarrow X$ such that $f \circ g \simeq \text{id}_Y$ and $g \circ f \simeq \text{id}_X$. We call g a *homotopy inverse* for f .

If a homotopy equivalence $f : X \rightarrow Y$ exists, we say that X and Y are homotopy equivalent and write $X \simeq Y$.

We are soon going to prove that this is indeed an equivalence relation on spaces, but we first look at some examples of homotopy equivalent spaces. Clearly, homeomorphic spaces are homotopy equivalent. However, we will see that we can do much more “violent” things to a space and still be homotopy equivalent.

Example. Let $X = S^1$, $Y = \mathbb{R}^2 \setminus \{0\}$. We have a natural inclusion map $i : X \hookrightarrow Y$. To obtain a map $Y \rightarrow X$, we can project each point onto the circle.



In particular, we define $r : Y \rightarrow X$ by

$$r(y) = \frac{y}{\|y\|}.$$

We immediately have $r \circ i = \text{id}_X$. We will now see that $i \circ r \simeq \text{id}_Y$. The composition $i \circ r$ first projects each object into S^1 , and then includes it back into $\mathbb{R}^2 \setminus \{0\}$. So this is just the projection map. We can define a homotopy $H : Y \times I \rightarrow Y$ by

$$H(y, t) = \frac{y}{t + (1 - t)\|y\|}.$$

This is continuous, and $H(\cdot, 0) = i \circ r$, $H(\cdot, 1) = \text{id}_Y$.

As we have said, homotopy equivalence can do really “violent” things to a space. We started with a 2-dimensional $\mathbb{R}^2 \setminus \{0\}$ space, and squashed it into a one-dimensional sphere.

Hence we see that homotopy equivalence doesn’t *care* about dimensions. Dimensions seems to be a rather fundamental thing in geometry, and we are discarding it here. So what is left? What does homotopy equivalence preserve?

While S^1 and $\mathbb{R}^2 \setminus \{0\}$ seem rather different, they have something in common — they both have a “hole”. We will later see that this is what homotopy equivalence preserves.

Example. Let $Y = \mathbb{R}^n$, $X = \{0\} = *$. Let $X \hookrightarrow Y$ be the inclusion map, and $r : Y \rightarrow X$ be the unique map that sends everything to $\{0\}$. Again, we have $r \circ i = \text{id}_X$. We can also obtain a homotopy from $i \circ r$ to id_Y by

$$H(y, t) = ty.$$

Again, from the point of view of homotopy theory, \mathbb{R}^n is just the same as a point! You might think that this is something crazy to do — we have just given up a lot of structure of topological spaces. However, by giving up these structures, it is easier to focus on what we really want to care about — holes. For example,

it is often much easier to argue about the one-point space $*$ than the whole of \mathbb{R}^2 ! By studying properties that are preserved by homotopy equivalence, and not just homeomorphism, we can simplify our problems by reducing complicated spaces to simpler ones via homotopy equivalence.

In general things homotopy equivalent to a single point are known as *contractible spaces*.

Notation. $*$ denotes the one-point space $\{0\}$.

Definition (Contractible space). If $X \simeq *$, then X is *contractible*.

We now show that homotopy equivalence of spaces is an equivalence relation. To do this, we first need a lemma.

Lemma. Consider the spaces and arrows

$$\begin{array}{ccccc} X & \xrightarrow{f_0} & Y & \xrightarrow{g_0} & Z \\ & \searrow f_1 & & \searrow g_1 & \\ & & Y & \xrightarrow{g_1} & Z \end{array}$$

If $f_0 \simeq_H f_1$ and $g_0 \simeq_{H'} g_1$, then $g_0 \circ f_0 \simeq g_1 \circ f_1$.

Proof. We will show that $g_0 \circ f_0 \simeq g_0 \circ f_1 \simeq g_1 \circ f_1$. Then we are done since homotopy between maps is an equivalence relation. So we need to write down two homotopies.

(i) Consider the following composition:

$$X \times I \xrightarrow{H} Y \xrightarrow{g_0} Z$$

It is easy to check that this is the first homotopy we need to show $g_0 \circ f_0 \simeq g_0 \circ f_1$.

(ii) The following composition is a homotopy from $g_0 \circ f_1$ to $g_1 \circ f_1$:

$$X \times I \xrightarrow{f_1 \times \text{id}_I} Y \times I \xrightarrow{H'} Z$$

□

Proposition. Homotopy equivalence on spaces is an equivalence relation.

Proof. Symmetry and reflexivity are trivial. To show transitivity, let $f : X \rightarrow Y$ and $h : Y \rightarrow Z$ be homotopy equivalences, and $g : Y \rightarrow X$ and $k : Z \rightarrow Y$ be their homotopy inverses. We will show that $h \circ f : X \rightarrow Z$ is a homotopy equivalence with homotopy inverse $g \circ k$. We have

$$(h \circ f) \circ (g \circ k) = h \circ (f \circ g) \circ k \simeq h \circ \text{id}_Y \circ k = h \circ k \simeq \text{id}_Z.$$

Similarly,

$$(g \circ k) \circ (h \circ f) = g \circ (k \circ h) \circ f \simeq g \circ \text{id}_Y \circ f = g \circ f \simeq \text{id}_X.$$

So done. □

Definition (Retraction). Let $A \subseteq X$ be a subspace. A *retraction* $r : X \rightarrow A$ is a map such that $r \circ i = \text{id}_A$, where $i : A \hookrightarrow X$ is the inclusion.

This map sends everything in X to A without moving things in A . Roughly speaking, if such a retraction exists, then A is no more complicated than X .

Definition (Deformation retraction). The retraction r is a *deformation retraction* if $i \circ r \simeq \text{id}_X$. A deformation retract is *strong* if we require this homotopy to be a homotopy rel A .


Roughly, this says that A is as complicated as X .

Example. Take X any space, and $A = \{x\} \subseteq X$. Then the constant map $r : X \rightarrow A$ is a retraction. If X is contractible, then this is a deformation retract.

2.2 Paths

Definition (Path). A *path* in a space X is a map $\gamma : I \rightarrow X$. If $\gamma(0) = x_0$ and $\gamma(1) = x_1$, we say γ is a path from x_0 to x_1 , and write $\gamma : x_0 \rightsquigarrow x_1$.

If $\gamma(0) = \gamma(1)$, then γ is called a *loop* (based at x_0).



Note that this map does not have to be injective. It can be self intersecting or do all sorts of weird stuff.

Recall that the basic idea of algebraic topology is to assign to each space X an (algebraic) object, which is (hopefully) easier to deal with. In homotopy theory, we do so using the idea of paths.


To do so, we need to be able to perform operations on paths.

Definition (Concatenation of paths). If we have two paths γ_1 from x_0 to x_1 ; and γ_2 from x_1 to x_2 , we define the *concatenation* to be

$$(\gamma_1 \cdot \gamma_2)(t) = \begin{cases} \gamma_1(2t) & 0 \leq t \leq \frac{1}{2} \\ \gamma_2(2t - 1) & \frac{1}{2} \leq t \leq 1. \end{cases}$$

This is continuous by the gluing lemma.

Note that we concatenate left to right, but function composition goes from right to left.



Definition (Inverse of path). The *inverse* of path $\gamma : I \rightarrow X$ is defined by

$$\gamma^{-1}(t) = \gamma(1 - t).$$

This is exactly the same path but going in the opposite direction.


What else do we need? We need an identity.

Definition (Constant path). The *constant path* at a point $x \in X$ is given by $c_x(t) = x$.

We haven't actually got a good algebraic system. We have γ and γ^{-1} , but when we compose them, we get a path from x_1 to x_2 and back, and not the identity. Also, we are not able to combine arbitrary paths in a space.

Before we make these into proper algebraic operations, we will talk about something slightly different. We can view this as a first attempt at associating things to topological spaces.

Definition (Path components). We can define a relation on X : $x_1 \sim x_2$ if there exists a path from x_1 to x_2 . By the concatenation, inverse and constant paths, \sim is an equivalence relation. The equivalence classes $[x]$ are called *path components*. We denote the quotient X/\sim by $\pi_0(X)$.



In the above space, we have three path components.

This isn't really a very useful definition, since most spaces we care about are path-connected, i.e. only have one path component. However, this is a first step at associating *something* to spaces. We can view this as a "toy model" for the more useful definitions we will later have.

One important property of this π_0 is that not only does it associate a set to each topological space, but also associates a function between the corresponding sets to each continuous map.

Proposition. For any map $f : X \rightarrow Y$, there is a well-defined function

$$\pi_0(f) : \pi_0(X) \rightarrow \pi_0(Y),$$

defined by

$$\pi_0(f)([x]) = [f(x)].$$

Furthermore,

- (i) If $f \simeq g$, then $\pi_0(f) = \pi_0(g)$.
- (ii) For any maps $A \xrightarrow{h} B \xrightarrow{k} C$, we have $\pi_0(k \circ h) = \pi_0(k) \circ \pi_0(h)$.
- (iii) $\pi_0(\text{id}_X) = \text{id}_{\pi_0(X)}$

Proof. To show this is well-defined, suppose $[x] = [y]$. Then let $\gamma : I \rightarrow X$ be a path from x to y . Then $f \circ \gamma$ is a path from $f(x)$ to $f(y)$. So $[f(x)] = [f(y)]$.

- (i) If $f \simeq g$, let $H : X \times I \rightarrow Y$ be a homotopy from f to g . Let $x \in X$. Then $H(x, \cdot)$ is a path from $f(x)$ to $g(x)$. So $[f(x)] = [g(x)]$, i.e. $\pi_0(f)([x]) = \pi_0(g)([x])$. So $\pi_0(f) = \pi_0(g)$.
- (ii) $\pi_0(k \circ h)([x]) = \pi_0(k) \circ \pi_0(h)([x]) = [k(h(x))]$.
- (iii) $\pi_0(\text{id}_X)([x]) = [\text{id}_X(x)] = [x]$. So $\pi_0(\text{id}_X) = \text{id}_{\pi_0(X)}$.

□


Corollary. If $f : X \rightarrow Y$ is a homotopy equivalence, then $\pi_0(f)$ is a bijection.

Example. The two point space $X = \{-1, 1\}$ is not contractible, because $|\pi_0(X)| = 2$, but $|\pi_0(*)| = 1$.

This is a rather silly example, since we can easily prove it directly. However, this is an example to show how we can use these machinery to prove topological results.

Now let's return to our operations on paths, and try to make them algebraic.


Definition (Homotopy of paths). Paths $\gamma, \gamma' : I \rightarrow X$ are *homotopic as paths* if they are homotopic rel $\{0, 1\} \subseteq I$, i.e. the end points are fixed. We write $\gamma \simeq \gamma'$.




Note that we would necessarily want to fix the two end points. Otherwise, if we allow end points to move, we can shrink any path into a constant path, and our definition of homotopy would be rather silly.

This homotopy works well with our previous operations on paths.

Proposition. Let $\gamma_1, \gamma_2 : I \rightarrow X$ be paths, $\gamma_1(1) = \gamma_2(0)$. Then if $\gamma_1 \simeq \gamma'_1$ and $\gamma_2 \simeq \gamma'_2$, then $\gamma_1 \cdot \gamma_2 \simeq \gamma'_1 \cdot \gamma'_2$.



Proof. Suppose that $\gamma_1 \simeq_{H_1} \gamma'_1$ and $\gamma_2 \simeq_{H_2} \gamma'_2$. Then we have the diagram



We can thus construct a homotopy by

$$H(s, t) = \begin{cases} H_1(s, 2t) & 0 \leq t \leq \frac{1}{2} \\ H_2(s, 2t - 1) & \frac{1}{2} \leq t \leq 1 \end{cases}.$$

□


To solve all our previous problems about operations on paths not behaving well, we can look at paths *up to homotopy*.

Proposition. Let $\gamma_0 : x_0 \rightsquigarrow x_1, \gamma_1 : x_1 \rightsquigarrow x_2, \gamma_2 : x_2 \rightsquigarrow x_3$ be paths. Then


- (i) $(\gamma_0 \cdot \gamma_1) \cdot \gamma_2 \simeq \gamma_0 \cdot (\gamma_1 \cdot \gamma_2)$
- (ii) $\gamma_0 \cdot c_{x_1} \simeq \gamma_0 \simeq c_{x_0} \cdot \gamma_0$.
- (iii) $\gamma_0 \cdot \gamma_0^{-1} \simeq c_{x_0}$ and $\gamma_0^{-1} \cdot \gamma_0 \simeq c_{x_1}$.

Proof.


- (i) Consider the following diagram:



- (ii) Consider the following diagram:



(iii) Consider the following diagram:



□

Turning these into proper proofs is left as an exercise for the reader.

2.3 The fundamental group

The idea is to take spaces and turn them into groups. We want to try to do this using paths. We've seen that if we want to do this, we should not work directly with paths, but paths up to homotopy. According to our proposition, this operation satisfies associativity, inverses and identity. The last issue we have to resolve is that we can't actually put two paths together unless they have the same start and end points.

The idea is to fix one of the points x_0 in our space, and only think about loops that start and end at x_0 . So we can always join two paths together.

This tells us that we aren't going to just think about spaces, but spaces *with basepoints*. We've now got everything we need.

Definition (Fundamental group). Let X be a space and $x_0 \in X$. The *fundamental group* of X (based at x_0), denoted $\pi_1(X, x_0)$, is the set of homotopy classes of loops in X based at x_0 (i.e. $\gamma(0) = \gamma(1) = x_0$). The group operations are defined as follows:

We define an operation by $[\gamma_0][\gamma_1] = [\gamma_0 \cdot \gamma_1]$; inverses by $[\gamma]^{-1} = [\gamma^{-1}]$; and the identity as the constant path $e = [c_{x_0}]$.

Often, when we write the homotopy classes of paths $[\gamma]$, we just get lazy and write γ .

Theorem. The fundamental group is a group.

Proof. Immediate from our previous lemmas. □

Often, in mathematics, after defining a term, we give lots of examples of it. Unfortunately, it is rather difficult to prove that a space has a non-trivial fundamental group, until we have developed some relevant machinery. Hence we will have to wait for a while before we have some concrete examples. Instead, we will look at some properties of the fundamental group first.

Definition (Based space). A *based space* is a pair (X, x_0) of a space X and a point $x_0 \in X$, the *basepoint*. A *map of based spaces*

$$f : (X, x_0) \rightarrow (Y, y_0)$$

is a continuous map $f : X \rightarrow Y$ such that $f(x_0) = y_0$. A *based homotopy* is a homotopy $\text{rel } \{x_0\}$.

Recall that for π_0 , for every map $f : X \rightarrow Y$, we can associate a function $\pi_0(f) : \pi_0(X) \rightarrow \pi_0(Y)$. We can do the same for π_1 .

Proposition. To a based map

$$f : (X, x_0) \rightarrow (Y, y_0),$$

there is an associated function

$$f_* = \pi_1(f) : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0),$$

defined by $[\gamma] \mapsto [f \circ \gamma]$. Moreover, it satisfies

- (i) $\pi_1(f)$ is a homomorphism of groups.
- (ii) If $f \simeq f'$, then $\pi_1(f) = \pi_1(f')$.
- (iii) For any maps $(A, a) \xrightarrow{h} (B, b) \xrightarrow{k} (C, c)$, we have $\pi_1(k \circ h) = \pi_1(k) \circ \pi_1(h)$.
- (iv) $\pi_1(\text{id}_X) = \text{id}_{\pi_1(X, x_0)}$

Proof. Exercise. □


In category-theoretic language, we say that π_1 is a *functor*.

So far so good. However, to define the fundamental group, we had to make a compromise and pick a basepoint. But we just care about the space. We don't want a basepoint! Hence, we should look carefully what happens when we change the basepoint, and see if we can live without it.

The first observation is that $\pi_1(X, x_0)$ only “sees” the path component of x_0 , since all loops based at x_0 can only live inside the path component of x_0 . Hence, the first importance of picking a basepoint is picking a path component.

For all practical purposes, we just assume that X is path connected, since if we weren't, the fundamental group just describes a particular path component of the original space.

Now we want to compare fundamental groups with different basepoints. Suppose we have two basepoints x_0 and x_1 . Suppose we have a loop γ at x_0 . How can we turn this into a loop based at x_1 ? This is easy. We first pick a path $u : x_0 \rightsquigarrow x_1$. Then we can produce a new loop at x_1 by going along u^{-1} to x_0 , take the path γ , and then return to x_0 by u , i.e. consider $u^{-1} \cdot \gamma \cdot u$.



Proposition. A path $u : x_0 \rightsquigarrow x_1$ induces a group *isomorphism*

$$u_{\#} : \pi_1(X, x_0) \rightarrow \pi_1(X, x_1)$$

by

$$[\gamma_0] \mapsto [u^{-1} \cdot \gamma \cdot u].$$

This satisfies

- (i) If $u \simeq u'$, then $u_{\#} = u'_{\#}$.
- (ii) $(c_{x_0})_{\#} = \text{id}_{\pi_1(X, x_0)}$
- (iii) If $v : x_1 \rightsquigarrow x_2$. Then $(u \cdot v)_{\#} = v_{\#} \circ u_{\#}$.
- (iv) If $f : X \rightarrow Y$ with $f(x_0) = y_0$, $f(x_1) = y_1$, then

$$(f \circ u)_{\#} \circ f_{*} = f_{*} \circ u_{\#} : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_1).$$

A nicer way of writing this is

$$\begin{array}{ccc} \pi_1(X, x_0) & \xrightarrow{f_*} & \pi_1(Y, y_0) \\ \downarrow u_{\#} & & \downarrow (f \circ u)_{\#} \\ \pi_1(X, x_1) & \xrightarrow{f_*} & \pi_1(Y, y_1) \end{array}$$

The property says that the composition is the same no matter which way we go from $\pi_1(X, x_0)$ to $\pi_1(Y, y_1)$. We say that the square is a *commutative diagram*. These diagrams will appear all of the time in this course.

- (v) If $x_1 = x_0$, then $u_{\#}$ is an automorphism of $\pi_1(X, x_0)$ given by conjugation by u .

It is important (yet difficult) to get the order of concatenation and composition right. Path concatenation is from left to right, while function composition is from right to left.

Proof. Yet another exercise. Note that $(u^{-1})_{\#} = (u_{\#})^{-1}$, which is why we have an isomorphism. \square

The main takeaway is that if x_0 and x_1 are in the same path component, then

$$\pi_1(X, x_0) \cong \pi_1(X, x_1).$$

So the basepoint isn't really too important. However, we have to be careful. While the two groups are isomorphic, the actual isomorphism depends on *which* path $u : x_0 \rightsquigarrow x_1$ we pick. So there is no natural isomorphism between the two groups. In particular, we cannot say "let $\alpha \in \pi_1(X, x_0)$. Now let α' be the corresponding element in $\pi_1(X, x_1)$ ".

We can also see if (X, x_0) and (Y, y_0) are based homotopy equivalent, then $\pi_1(X, x_0) \cong \pi_1(Y, y_0)$. Indeed, if they are homotopy equivalent, then there is some $f : X \rightarrow Y$, $g : Y \rightarrow X$ such that

$$f \circ g \simeq \text{id}_Y, \quad g \circ f \simeq \text{id}_X.$$

So

$$f_* \circ g_* = \text{id}_{\pi_1(Y, y_0)}, \quad g_* \circ f_* = \text{id}_{\pi_1(X, x_0)},$$


and f_* and g_* are the isomorphisms we need.

However, can we do this with non-based homotopies? Suppose we have a space X , and a space Y , and functions $f, g : X \rightarrow Y$, with a homotopy $H : X \times I \rightarrow Y$ from f to g . If this were a based homotopy, we know

$$f_* = g_* : \pi_1(X, x_0) \rightarrow \pi_1(Y, f(x_0)).$$

Now we don't insist that this homotopy fixes x_0 , and we could have $f(x_0) \neq g(x_0)$. How can we relate $f_* : \pi_1(X, x_0) \rightarrow \pi_1(Y, f(x_0))$ and $g_* : \pi_1(X, x_0) \rightarrow \pi_1(Y, g(x_0))$?

First of all, we need to relate the groups $\pi_1(Y, f(x_0))$ and $\pi_1(Y, g(x_0))$. To do so, we need find a path from $f(x_0)$ to $g(x_0)$. To produce this, we can use the homotopy H . We let $u : f(x_0) \rightsquigarrow g(x_0)$ with $u = H(x_0, \cdot)$. Now we have three maps f_*, g_* and $u_\#$. Fortunately, these do fit together well.



Lemma. The following diagram commutes:

$$\begin{array}{ccc} & \pi_1(Y, f(x_0)) & \\ f_* \nearrow & \downarrow u_\# & \\ \pi_1(X, x_0) & & \pi_1(Y, g(x_0)) \\ g_* \searrow & & \end{array}$$

In algebra, we say

$$g_* = u_\# \circ f_*.$$

Proof. Suppose we have a loop $\gamma : I \rightarrow X$ based at x_0 .

We need to check that

$$g_*([\gamma]) = u_\# \circ f_*([\gamma]).$$

In other words, we want to show that

$$g \circ \gamma \simeq u^{-1} \cdot (f \circ \gamma) \cdot u.$$

To prove this result, we want to build a homotopy.

Consider the composition:


$$F : I \times I \xrightarrow{\gamma \times \text{id}_I} X \times I \xrightarrow{H} Y.$$

Our plan is to exhibit two homotopic paths ℓ^+ and ℓ^- in $I \times I$ such that

$$F \circ \ell^+ = g \circ \gamma, \quad F \circ \ell^- = u^{-1} \cdot (f \circ \gamma) \cdot u.$$

This is in general a good strategy — X is a complicated and horrible space we don't understand. So to construct a homotopy, we make ourselves work in a much nicer space $I \times I$.

Our ℓ^+ and ℓ^- are defined in a rather simple way.



More precisely, ℓ^+ is the path $s \mapsto (s, 1)$, and ℓ^- is the concatenation of the paths $s \mapsto (0, 1 - s)$, $s \mapsto (s, 0)$ and $s \mapsto (1, s)$.

Note that ℓ^+ and ℓ^- are homotopic as paths. If this is not obvious, we can manually check the homotopy

$$L(s, t) = t\ell^+(s) + (1 - t)\ell^-(s).$$

This works because $I \times I$ is convex. Hence $F \circ \ell^+ \simeq_{F \circ L} F \circ \ell^-$ as paths.

Now we check that the compositions $F \circ \ell^\pm$ are indeed what we want. We have

$$F \circ \ell^+(s) = H(\gamma(s), 1) = g \circ \gamma.$$

Similarly, we can show that

$$F \circ \ell^-(s) = u^{-1} \cdot (f \circ \gamma) \cdot u.$$

So done. □

It is worth looking at this proof hard and truly understand what is going on, since this is a really good example of how we can construct interesting homotopies.

With this lemma, we can show that fundamental groups really respect homotopies.


Theorem. If $f : X \rightarrow Y$ is a homotopy equivalence, and $x_0 \in X$, then the induced map

$$f_* : \pi_1(X, x_0) \rightarrow \pi_1(Y, f(x_0)).$$

is an isomorphism.

While this seems rather obvious, it is actually non-trivial if we want to do it from scratch. While we are given a homotopy equivalence, we are given no guarantee that the homotopy respects our basepoints. So this proof involves some real work.

Proof. Let $g : Y \rightarrow X$ be a homotopy inverse. So $f \circ g \simeq_H \text{id}_Y$ and $g \circ f \simeq'_H \text{id}_X$.



We have no guarantee that $g \circ f(x_0) = x_0$, but we know that our homotopy H' gives us $u' = H'(x_0, \cdot) : x_0 \rightsquigarrow g \circ f(x_0)$.

Applying our previous lemma with id_X for “ f ” and $g \circ f$ for “ g ”, we get

$$u'_{\#} \circ (\text{id}_X)_* = (g \circ f)_*$$

Using the properties of the $*$ operation, we get that

$$g_* \circ f_* = u'_{\#}.$$

However, we know that $u'_{\#}$ is an isomorphism. So f_* is injective and g_* is surjective.

Doing it the other way round with $f \circ g$ instead of $g \circ f$, we know that g_* is injective and f_* is surjective. So both of them are isomorphisms. \square

With this theorem, we can finally be sure that the fundamental group is a property of the space, without regards to the basepoint (assuming path connectedness), and is preserved by arbitrary homotopies.

We now use the fundamental group to define several simple notions.

Definition (Simply connected space). A space X is *simply connected* if it is path connected and $\pi_1(X, x_0) \cong 1$ for some (any) choice of $x_0 \in X$.

Example. Clearly, a point $*$ is simply connected since there is only one path on $*$ (the constant path). Hence, any contractible space is simply connected since it is homotopic to $*$. For example, \mathbb{R}^n is simply connected for all spaces.

There is a useful characterization of simply connected spaces:

Lemma. A path-connected space X is simply connected if and only if for any $x_0, x_1 \in X$, there exists a unique homotopy class of paths $x_0 \rightsquigarrow x_1$.

Proof. Suppose X is simply connected, and let $u, v : x_0 \rightsquigarrow x_1$ be paths. Now note that $u \cdot v^{-1}$ is a loop based at x_0 , it is homotopic to the constant path, and $v^{-1} \cdot v$ is trivially homotopic to the constant path. So we have

$$u \simeq u \cdot v^{-1} \cdot v \simeq v.$$

On the other hand, suppose there is a unique homotopy class of paths $x_0 \rightsquigarrow x_1$ for all $x_0, x_1 \in X$. Then in particular there is a unique homotopy class of loops based at x_0 . So $\pi_1(X, x_0)$ is trivial. \square

3 Covering spaces

We can ask ourselves a question — what are groups? We can write down a definition in terms of operations and axioms, but this is not what groups *are*. Groups were created to represent *symmetries* of objects. In particular, a group should be *acting* on something. If not, something wrong is probably going on.


We have created the fundamental group. So what do they act on? Can we find something on which these fundamental groups act on?

An answer to this would also be helpful in more practical terms. So far we have not exhibited a non-trivial fundamental group. This would be easy if we can make the group act on something — if the group acts non-trivially on our thing, then clearly the group cannot be trivial.

These things we act on are *covering spaces*.

3.1 Covering space

Intuitively, a *covering space* of X is a pair $(\tilde{X}, p : \tilde{X} \rightarrow X)$, such that if we take any $x_0 \in X$, there is some neighbourhood U of x_0 such that the pre-image of the neighbourhood is “many copies” of U .



Definition (Covering space). A *covering space* of X is a pair $(\tilde{X}, p : \tilde{X} \rightarrow X)$, such that each $x \in X$ has a neighbourhood U which is *evenly covered*.

Whether we require p to be surjective is a matter of taste. However, it will not matter if X is path-connected, which is the case we really care about.


Definition (Evenly covered). U is *evenly covered* by p if

$$p^{-1}(U) \cong \coprod_{\alpha \in \Lambda} V_{\alpha},$$

where $p|_{V_{\alpha}} : V_{\alpha} \rightarrow U$ is a homeomorphism, and each of the V_{α} is open.

Example. Homeomorphisms are covering maps. Duh.


Example. Consider $p : \mathbb{R} \rightarrow S^1 \subseteq \mathbb{C}$ defined by $t \mapsto e^{2\pi it}$. This is a covering space.



Here we have $p^{-1}(1) = \mathbb{Z}$.

As we said, we are going to use covering spaces to determine non-trivial fundamental groups. Before we do that, we can guess what the fundamental group of S^1 is. If we have a loop on S^1 , it could be nothing, it can loop around the circle once, or loop around many times. So we can characterize each loop by the number of times it loops around the circle, and it should not be difficult to convince ourselves that two loops that loop around the same number of times can be continuously deformed to one another. So it is not unreasonable to guess that $\pi_1(S, 1) \cong \mathbb{Z}$. However, we also know that $p^{-1}(1) = \mathbb{Z}$. In fact, for any point $z \in S^1$, $p^{-1}(z)$ is just “ \mathbb{Z} many copies” of z . We will show that this is not a coincidence.


Example. Consider $p_n : S^1 \rightarrow S^1$ (for any $n \in \mathbb{Z} \setminus \{0\}$) defined by $z \mapsto z^n$. We can consider this as “winding” the circle n times, or as the following covering map:



where we join the two red dots together.

This time the pre-image of 1 would be n copies of 1, instead of \mathbb{Z} copies of 1.

Example. Consider $X = \mathbb{RP}^2$, which is the real projective plane. This is defined by S^2/\sim , where we identify every $x \simeq -x$, i.e. every pair of antipodal point. We can also think of this as the space of lines in \mathbb{R}^3 . This is actually really difficult to draw (and in fact impossible to fully embed in \mathbb{R}^3), so we will not attempt. There is, however, a more convenient way of thinking about \mathbb{RP}^2 . We just use the definition — we imagine an S^2 , but this time, instead of a point being a “point” on the sphere, it is a *pair* of antipodal points.



We define $p : S^2 \rightarrow \mathbb{RP}^2$ to be the quotient map. Then this is a covering map, since the pre-image of a small neighbourhood of any x_0 is just two copies of the neighbourhood.


As we mentioned, a covering space of X is (locally) like many “copies” of X . Hence, given any function $f : Y \rightarrow X$, a natural question is whether we can “lift” it to a function $\tilde{f} : Y \rightarrow \tilde{X}$, by mapping to one of the copies of X in \tilde{X} .

This is not always possible, but when it is, we call this a lift.

Definition (Lifting). Let $f : Y \rightarrow X$ be a map, and $p : \tilde{X} \rightarrow X$ a covering space. A *lift* of f is a map $\tilde{f} : Y \rightarrow \tilde{X}$ such that $f = p \circ \tilde{f}$, i.e. the following diagram commutes:

$$\begin{array}{ccc} & & \tilde{X} \\ & \nearrow \tilde{f} & \downarrow p \\ Y & \xrightarrow{f} & X \end{array}$$

We can visualize this in the case where Y is the unit interval I and the map is just a path in X .



It feels that if we know which “copy” of X we lifted our map to, then we already know everything about \tilde{f} , since we are just moving our f from X to that particular copy of X . This is made precise by the following lemma:

Lemma. Let $p : \tilde{X} \rightarrow X$ be a covering map, $f : Y \rightarrow X$ be a map, and \tilde{f}_1, \tilde{f}_2 are both lifts of f . Then

$$S = \{y \in Y : \tilde{f}_1(y) = \tilde{f}_2(y)\}$$

is both open and closed. In particular, if Y is connected, \tilde{f}_1 and \tilde{f}_2 agree either everywhere or nowhere.

This is sort of a “uniqueness statement” for a lift. If we know a point in the lift, then we know the whole path. This is since once we’ve decided our starting point, ie, which “copy” of X we work in, the rest of \tilde{f} has follow what f does.

Proof. First we show it is open. Let y be such that $\tilde{f}_1(y) = \tilde{f}_2(y)$. Then there is an evenly covered open neighbourhood $U \subseteq X$ of $f(y)$. Let \tilde{U} be such that $\tilde{f}_1(y) \in \tilde{U}$, $p(\tilde{U}) = U$ and $p|_{\tilde{U}} : \tilde{U} \rightarrow U$ is a homeomorphism. Let $V = \tilde{f}_1^{-1}(\tilde{U}) \cap \tilde{f}_2^{-1}(\tilde{U})$. We will show that $\tilde{f}_1 = \tilde{f}_2$ on V .

Indeed, by construction

$$p|_{\tilde{U}} \circ \tilde{f}_1|_V = p|_{\tilde{U}} \circ \tilde{f}_2|_V.$$

Since $p|_{\tilde{U}}$ is a homeomorphism, it follows that

$$\tilde{f}_1|_V = \tilde{f}_2|_V.$$

Now we show S is closed. Suppose not. Then there is some $y \in \bar{S} \setminus S$. So $\tilde{f}_1(y) \neq \tilde{f}_2(y)$. Let U be an evenly covered neighbourhood of $f(y)$. Let $p^{-1}(U) = \coprod U_\alpha$. Let $\tilde{f}_1(y) \in U_\beta$ and $\tilde{f}_2(y) \in U_\gamma$, where $\beta \neq \gamma$. Then $V = \tilde{f}_1^{-1}(U_\beta) \cap \tilde{f}_2^{-1}(U_\gamma)$ is an open neighbourhood of y , and hence intersects S by definition of closure. So there is some $x \in V$ such that $\tilde{f}_1(x) = \tilde{f}_2(x)$. But $\tilde{f}_1(x) \in U_\beta$ and $\tilde{f}_2(x) \in U_\gamma$, and hence U_β and U_γ have a non-trivial intersection. This is a contradiction. So S is closed. \square

We just had a uniqueness statement. How about existence? Given a map, is there guarantee that we can lift it to something? Moreover, if I have fixed a “copy” of X I like, can I also lift my map to that copy? We will later come up with a general criterion for when lifts exist. However, it turns out homotopies can always be lifted.

Lemma (Homotopy lifting lemma). Let $p : \tilde{X} \rightarrow X$ be a covering space, $H : Y \times I \rightarrow X$ be a homotopy from f_0 to f_1 . Let \tilde{f}_0 be a lift of f_0 . Then there exists a *unique* homotopy $\tilde{H} : Y \times I \rightarrow \tilde{X}$ such that

- (i) $\tilde{H}(\cdot, 0) = \tilde{f}_0$; and
- (ii) \tilde{H} is a lift of H , i.e. $p \circ \tilde{H} = H$.

This lemma might be difficult to comprehend at first. We can look at the special case where $Y = *$. Then a homotopy is just a path. So the lemma specializes to

Lemma (Path lifting lemma). Let $p : \tilde{X} \rightarrow X$ be a covering space, $\gamma : I \rightarrow X$ a path, and $\tilde{x}_0 \in \tilde{X}$ such that $p(\tilde{x}_0) = x_0 = \gamma(0)$. Then there exists a *unique* path $\tilde{\gamma} : I \rightarrow \tilde{X}$ such that

- (i) $\tilde{\gamma}(0) = \tilde{x}_0$; and
- (ii) $\tilde{\gamma}$ is a lift of γ , i.e. $p \circ \tilde{\gamma} = \gamma$.

This is exactly the picture we were drawing before. We just have to start at a point \tilde{x}_0 , and then everything is determined because locally, everything upstairs in \tilde{X} is just like X . Note that we have already proved uniqueness. So we just need to prove existence.

In theory, it makes sense to prove homotopy lifting, and path lifting comes immediately as a corollary. However, the proof of homotopy lifting is big and scary. So instead, we will prove path lifting, which is something we can more easily visualize and understand, and then use that to prove homotopy lifting.

Proof. Let

$$S = \{s \in I : \tilde{\gamma} \text{ exists on } [0, s] \subseteq I\}.$$

Observe that

- (i) $0 \in S$.
- (ii) S is open. If $s \in S$ and $\tilde{\gamma}(s) \in V_\beta \subseteq p^{-1}(U)$, we can define $\tilde{\gamma}$ on some small neighbourhood of s by

$$\tilde{\gamma}(t) = (p|_{V_\beta})^{-1} \circ \gamma$$

- (iii) S is closed. If $s \notin S$, then pick an evenly covered neighbourhood U of $\gamma(s)$. Suppose $\gamma((s - \varepsilon, s)) \subseteq U$. So $s - \frac{\varepsilon}{2} \notin S$. So $(s - \frac{\varepsilon}{2}, 1] \cap S = \emptyset$. So S is closed.

Since S is both open and closed, and is non-empty, we have $S = I$. So $\tilde{\gamma}$ exists. \square

How can we promote this to a proof of the homotopy lifting lemma? At every point $y \in Y$, we know what to do, since we have path lifting. So $\tilde{H}(y, \cdot)$ is defined. So the thing we have to do is to show that this is continuous. Steps of the proof are

- (i) Use compactness of I to argue that the proof of path lifting works on small neighbourhoods in Y .
- (ii) For each y , we pick an open neighbourhood U of y , and define a good path lifting on $U \times I$.
- (iii) By uniqueness of lifts, these path liftings agree when they overlap. So we have one big continuous lifting.

With the homotopy lifting lemma in our toolkit, we can start to use it to do stuff. So far, we have covering spaces and fundamental groups. We are now going to build a bridge between these two, and show how covering spaces can be used to reflect some structures of the fundamental group.

At least one payoff of this work is that we are going to exhibit some non-trivial fundamental groups.


We have just showed that we are allowed to lift homotopies. However, what we are really interested in is homotopy *as paths*. The homotopy lifting lemma does not tell us that the lifted homotopy preserves basepoints. This is what we are going to show.

Corollary. Suppose $\gamma, \gamma' : I \rightarrow X$ are paths $x_0 \rightsquigarrow x_1$ and $\tilde{\gamma}, \tilde{\gamma}' : I \rightarrow \tilde{X}$ are lifts of γ and γ' respectively, both starting at $\tilde{x}_0 \in p^{-1}(x_0)$.

If $\gamma \simeq \gamma'$ as *paths*, then $\tilde{\gamma}$ and $\tilde{\gamma}'$ are homotopic as paths. In particular, $\tilde{\gamma}(1) = \tilde{\gamma}'(1)$.

Note that if we cover the words “as paths” and just talk about homotopies, then this is just the homotopy lifting lemma. So we can view this as a stronger form of the homotopy lifting lemma.

Proof. The homotopy lifting lemma gives us a \tilde{H} , a lift of H with $\tilde{H}(\cdot, 0) = \tilde{\gamma}$.




In this diagram, we by assumption know the bottom of the \tilde{H} square is $\tilde{\gamma}$. To show that this is a path homotopy from $\tilde{\gamma}$ to $\tilde{\gamma}'$, we need to show that the other edges are $c_{\tilde{x}_0}$, $c_{\tilde{x}_1}$ and $\tilde{\gamma}'$ respectively.

Now $\tilde{H}(\cdot, 1)$ is a lift of $H(\cdot, 1) = \gamma'$, starting at \tilde{x}_0 . Since lifts are unique, we must have $\tilde{H}(\cdot, 1) = \tilde{\gamma}'$. So this is indeed a homotopy between $\tilde{\gamma}$ and $\tilde{\gamma}'$. Now we need to check that this is a homotopy of paths.

We know that $\tilde{H}(0, \cdot)$ is a lift of $H(0, \cdot) = c_{x_0}$. We are aware of one lift of c_{x_0} , namely $c_{\tilde{x}_0}$. By uniqueness of lift, we must have $\tilde{H}(0, \cdot) = c_{\tilde{x}_0}$. Similarly, $\tilde{H}(1, \cdot) = c_{\tilde{x}_1}$. So this is a homotopy of paths. \square


So far, our picture of covering spaces is like this:



Except... is it? Is it possible that we have four copies of x_0 but just three copies of x_1 ? This is obviously possible if X is not path connected — the component containing x_0 and the one containing x_1 are completely unrelated. But what if X is path connected?

Corollary. If X is a path connected space, $x_0, x_1 \in X$, then there is a bijection $p^{-1}(x_0) \rightarrow p^{-1}(x_1)$.

Proof. Let $\gamma : x_0 \rightsquigarrow x_1$ be a path. We want to use this to construct a bijection between each preimage of x_0 and each preimage of x_1 . The obvious thing to use is to use lifts of the path γ .



Define a map $f_\gamma : p^{-1}(x_0) \rightarrow p^{-1}(x_1)$ that sends \tilde{x}_0 to the end point of the unique lift of γ at \tilde{x}_0 .

The inverse map is obtained by replacing γ with γ^{-1} , i.e. $f_{\gamma^{-1}}$. To show this is an inverse, suppose we have some lift $\tilde{\gamma} : \tilde{x}_0 \rightsquigarrow \tilde{x}_1$, so that $f_\gamma(\tilde{x}_0) = \tilde{x}_1$. Now notice that $\tilde{\gamma}^{-1}$ is a lift of γ^{-1} starting at \tilde{x}_1 and ending at \tilde{x}_0 . So $f_{\gamma^{-1}}(\tilde{x}_1) = \tilde{x}_0$. So $f_{\gamma^{-1}}$ is an inverse to f_γ , and hence f_γ is bijective. \square

Definition (n -sheeted). A covering space $p : \tilde{X} \rightarrow X$ of a path-connected space X is n -sheeted if $|p^{-1}(x)| = n$ for all $x \in X$.

Each covering space has a number associated to it, namely the number of sheets. Is there any number we can assign to fundamental groups? Well, the index of a subgroup might be a good candidate. We'll later see if this is the case.

One important property of covering spaces is the following:

Lemma. If $p : \tilde{X} \rightarrow X$ is a covering map and $\tilde{x}_0 \in \tilde{X}$, then

$$p_* : \pi_1(\tilde{X}, \tilde{x}_0) \rightarrow \pi_1(X, x_0)$$

is injective.


Proof. To show that a group homomorphism p_* is injective, we have to show that if $p_*(x)$ is trivial, then x must be trivial.

Consider a based loop $\tilde{\gamma}$ in \tilde{X} . We let $\gamma = p \circ \tilde{\gamma}$. If γ is trivial, i.e. $\gamma \simeq c_{x_0}$ as paths, The homotopy lifting lemma then gives us a homotopy upstairs between $\tilde{\gamma}$ and $c_{\tilde{x}_0}$. So $\tilde{\gamma}$ is trivial. \square

As we have originally said, our objective is to make our fundamental group act on something. We are almost there already.

Let's look again at the proof that there is a bijection between $p^{-1}(x_0)$ and $p^{-1}(x_1)$. What happens if γ is a loop? For any $\tilde{x}_0 \in p^{-1}(x_0)$, we can look at the end point of the lift. This end point may or may not be our original \tilde{x}_0 . So each loop γ "moves" our \tilde{x}_0 to another \tilde{x}'_0 .

However, we are not really interested in paths themselves. We are interested in equivalence classes of paths under homotopy of paths. However, this is fine. If γ is homotopic to γ' , then this homotopy can be lifted to get a homotopy between $\tilde{\gamma}$ and $\tilde{\gamma}'$. In particular, these have the same end points. So each (based) homotopy class gives a well-defined endpoint.



Now this gives an action of $\pi_1(X, x_0)$ on $p^{-1}(x_0)$! Note, however, that this will not be the sort of actions we are familiar with. We usually work with *left*-actions, where the group acts on the left, but now we will have *right*-actions, which may confuse you a lot. To see this, we have to consider what happens when we perform two operations one after another, which you shall check yourself. We write this action as $\tilde{x}_0 \cdot [\gamma]$.

When we have an action, we are interested in two things — the orbits, and the stabilizers. This is what the next lemma tells us about.

Lemma. Suppose X is path connected and $x_0 \in X$.

- (i) The action of $\pi_1(X, x_0)$ on $p^{-1}(x_0)$ is transitive if and only if \tilde{X} is path connected. Alternatively, we can say that the orbits of the action correspond to the path components.
- (ii) The stabilizer of $\tilde{x}_0 \in p^{-1}(x_0)$ is $p_*(\pi(\tilde{X}, \tilde{x}_0)) \subseteq \pi(X, x_0)$.
- (iii) If \tilde{X} is path connected, then there is a bijection

$$p_*\pi_1(\tilde{X}, \tilde{x}_0) \backslash \pi_1(X, x_0) \rightarrow p^{-1}(x_0).$$

Note that $p_*\pi_1(\tilde{X}, \tilde{x}_0) \backslash \pi_1(X, x_0)$ is not a quotient, but simply the set of cosets. We write it the “wrong way round” because we have right cosets instead of left cosets.

Note that this is great! If we can find a covering space p and a point x_0 such that $p^{-1}(x_0)$ is non-trivial, then we immediately know that $\pi_1(X, x_0)$ is non-trivial!

Proof.

- (i) If $\tilde{x}_0, \tilde{x}'_0 \in p^{-1}(x_0)$, then since \tilde{X} is path connected, we know that there is some $\tilde{\gamma} : \tilde{x}_0 \rightsquigarrow \tilde{x}'_0$. Then we can project this to $\gamma = p \circ \tilde{\gamma}$. Then γ is a path from $x_0 \rightsquigarrow x_0$, i.e. a loop. Then by the definition on the action, $\tilde{x}_0 \cdot [\gamma] = \tilde{\gamma}(1) = \tilde{x}'_0$.
- (ii) Suppose $[\gamma] \in \text{stab}(\tilde{x}_0)$. Then $\tilde{\gamma}$ is a loop based at \tilde{x}_0 . So $\tilde{\gamma}$ defines $[\tilde{\gamma}] \in \pi_1(\tilde{X}, \tilde{x}_0)$ and $\gamma = p \circ \tilde{\gamma}$.
- (iii) This follows directly from the orbit-stabilizer theorem.

□

We now want to use this to determine that the fundamental group of a space is non-trivial. We can be more ambitious, and try to actually *find* $\pi_1(X, x_0)$. In the best possible scenario, we would have $\pi_1(\tilde{X}, \tilde{x}_0)$ trivial. Then we have a bijection between $\pi_1(X, x_0)$ and $p^{-1}(x_0)$. In other words, we want our covering space \tilde{X} to be simply connected.

Definition (Universal cover). A covering map $p : \tilde{X} \rightarrow X$ is a *universal cover* if \tilde{X} is simply connected.


We will look into universal covers in depth later and see what they really are.

Corollary. If $p : \tilde{X} \rightarrow X$ is a universal cover, then there is a bijection $\ell : \pi_1(X, x_0) \rightarrow p^{-1}(x_0)$.

Note that the orbit stabilizer theorem does not provide a canonical bijection between $p^{-1}(x_0)$ and $p_*\pi_1(\tilde{X}, \tilde{x}_0) \cong p_*\pi_1(\tilde{X}/\sim, \tilde{x}_0/\sim)$. To obtain a bijection, we need to pick a starting point $\tilde{x}_0 \in p^{-1}(x_0)$. So the above bijection ℓ depends on a choice of \tilde{x}_0 .

3.2 The fundamental group of the circle and its applications

Finally, we can exhibit a non-trivial fundamental group. We are going to consider the space S^1 and a universal covering \mathbb{R} .



Then our previous corollary gives


Corollary. There is a bijection $\pi_1(S^1, 1) \rightarrow p^{-1}(1) = \mathbb{Z}$.

What's next? We just know that $\pi_1(S^1, 1)$ is countably infinite, but can we work out the group structure?

We can, in fact, prove a stronger statement:

Theorem. The map $\ell : \pi_1(S^1, 1) \rightarrow p^{-1}(1) = \mathbb{Z}$ is a group isomorphism.

Proof. We know it is a bijection. So we need to check it is a group homomorphism. The idea is to write down representatives for what we think the elements should be.



Let $\tilde{u}_n : I \rightarrow \mathbb{R}$ be defined by $t \mapsto nt$, and let $u_n = p \circ \tilde{u}_n$. Since \mathbb{R} is simply connected, so there is a unique homotopy class between any two points. So for any $[\gamma] \in \pi_1(S^1, 1)$, if $\tilde{\gamma}$ is the lift to \mathbb{R} at 0 and $\tilde{\gamma}(1) = n$, then $\tilde{\gamma} \simeq \tilde{u}_n$ as paths. So $[\gamma] = [u_n]$.

To show that this has the right group operation, we can easily see that $\widehat{u_m \cdot u_n} = \tilde{u}_{m+n}$, since we are just moving by $n + m$ in both cases. Therefore

$$\ell([u_m][u_n]) = \ell([u_m \cdot u_n]) = m + n = \ell([u_{m+n}]).$$

So ℓ is a group isomorphism. \square

What have we done? In general, we might be given a horrible, crazy loop in S^1 . It would be rather difficult to work with it directly in S^1 . So we pull it up to the universal covering \mathbb{R} . Since \mathbb{R} is nice and simply connected, we can easily produce a homotopy that “straightens out” the path. We then project this homotopy down to S^1 , to get a homotopy from γ to u_n .


It is indeed possible to produce a homotopy directly inside S^1 from each loop to some u_n , but that would be tedious work that involves messing with a lot of algebra and weird, convoluted formulas.

With the fundamental group of the circle, we do many things. An immediate application is that we can properly define the “winding number” of a closed curve. Since $\mathbb{C} \setminus \{0\}$ is homotopy equivalent to S^1 , its fundamental group is \mathbb{Z} as well. Any closed curve $S^1 \rightarrow \mathbb{C} \setminus \{0\}$ thus induces a group homomorphism $\mathbb{Z} \rightarrow \mathbb{Z}$. Any such group homomorphism must be of the form $t \mapsto nt$, and the winding number is given by n . If we stare at it long enough, it is clear that this is exactly the number of times the curve winds around the origin.

Also, we have the following classic application:

Theorem (Brouwer’s fixed point theorem). Let $D^2 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$ be the unit disk. If $f : D^2 \rightarrow D^2$ is continuous, then there is some $x \in D^2$ such that $f(x) = x$.

Proof. Suppose not. So $x \neq f(x)$ for all $x \in D^2$.



We define $g : D^2 \rightarrow S^1$ as in the picture above. Then we know that g is continuous and g is a retraction from D^2 onto S^1 . In other words, the following composition is the identity:

$$\begin{array}{ccc} S^1 & \xleftarrow{\iota} & D^2 \xrightarrow{g} S^1 \\ & \searrow \text{id}_{S^1} & \nearrow \end{array}$$

Then this induces a homomorphism of groups whose composition is the identity:

$$\begin{array}{ccc} \mathbb{Z} & \xleftarrow{\iota_*} & \{0\} \xrightarrow{g_*} \mathbb{Z} \\ & \searrow \text{id}_{\mathbb{Z}} & \nearrow \end{array}$$

But this is clearly nonsense! So we must have had a fixed point. \square


But we have a problem. What about D^3 ? Can we prove a similar theorem? Here the fundamental group is of little use, since we can show that the fundamental group of S^n for $n \geq 2$ is trivial. Later in the course, we will be able to prove this theorem for higher dimensions, when we have developed more tools to deal with stuff.

3.3 Universal covers


We have defined universal covers mysteriously as covers that are simply connected. We have just shown that $p : \mathbb{R} \rightarrow S^1$ is a universal cover. In general, what do universal covers look like?

Let's consider a slightly more complicated example. What would be a universal cover of the torus $S^1 \times S^1$? An obvious guess would be $p \times p : \mathbb{R} \times \mathbb{R} \rightarrow S^1 \times S^1$. How can we visualize this?

First of all, how can we visualize a torus? Often, we just picture it as the surface of a doughnut. Alternatively, we can see it as a quotient of the square, where we identify the following edges:



Then what does it *feel* like to live in the torus? If you live in a torus and look around, you wouldn't see a boundary. The space just extends indefinitely for ever, somewhat like \mathbb{R}^2 . The difference is that in the torus, you aren't actually seeing free space out there, but just seeing copies of the same space over and over again. If you live inside the square, the universe actually looks like this:



As we said, this looks somewhat like \mathbb{R}^2 , but we *know* that this is not \mathbb{R}^2 , since we can see some symmetry in this space. Whenever we move one unit horizontally or vertically, we get back to “the same place”. In fact, we can move horizontally by n units and vertically by m units, for any $n, m \in \mathbb{Z}$, and still get back to the same place. This space has a huge translation symmetry. What is this symmetry? It is exactly $\mathbb{Z} \times \mathbb{Z}$.

We see that if we live inside the torus $S^1 \times S^1$, it feels like we are actually living in the universal covering space $\mathbb{R} \times \mathbb{R}$, except that we have an additional symmetry given by the fundamental group $\mathbb{Z} \times \mathbb{Z}$.

Hopefully, you are convinced that universal covers are nice. We would like to say that universal covers always exist. However, this is not always true.

Firstly, we should think — what would having a universal cover imply? Suppose X has a universal cover \tilde{X} . Pick any point $x_0 \in X$, and pick an evenly covered neighbourhood U in X . This lifts to some $\tilde{U} \in \tilde{X}$. If we draw a teeny-tiny loop γ around x_0 inside U , we can lift this γ to $\tilde{\gamma}$ in \tilde{U} . But we know that \tilde{X} is simply connected. So $\tilde{\gamma}$ is homotopic to the constant path. Hence γ is also homotopic to the constant path. So all loops (contained in U) at x_0 are homotopic to the constant path.

It seems like for every $x_0 \in X$, there is some neighbourhood of x_0 that is simply connected. Except that's not what we just showed above. The homotopy from $\tilde{\gamma}$ to the constant path is a homotopy in \tilde{X} , and can pass through anything

in \tilde{X} , not just \tilde{U} . Hence the homotopy induced in X is also a homotopy in X , not a homotopy in U . So U itself need not be simply connected. What we have is a slightly weaker notion.

Definition (Locally simply connected). X is *locally simply connected* if for all $x_0 \in X$, there is some neighbourhood U of x_0 such that U is simply connected.

As we mentioned, what we actually want is a weaker condition.

Definition (Semi-locally simply connected). X is *semi-locally simply connected* if for all $x_0 \in X$, there is some neighbourhood U of x_0 such that any loop γ based at x_0 is homotopic to c_{x_0} as paths in X .

We have just argued that if a universal cover $p : \tilde{X} \rightarrow X$ exists, then X is semi-locally simply connected. This is really not interesting, since we don't care if a space is semi-locally simply connected. What is important is that this is the other direction. We still need one more additional condition:

Definition (Locally path connected). A space X is *locally path connected* if for any point x and any neighbourhood V of x , there is some open path connected $U \subseteq V$ such that $x \in U$.

It is important to note that a path connected space need not be locally path connected. It is an exercise for the reader to come up with a counterexample.

Theorem. If X is path connected, locally path connected and semi-locally simply connected, then X has a universal covering.

Note that we can alternatively define a universal covering as a covering space of X that is also a covering space of all other covers of X . If we use this definition, then we can prove this result easily using Zorn's lemma. However, that proof is not too helpful since it does not tell us where the universal covering comes from. Instead, we will provide a constructive proof (sketch) that will hopefully be more indicative of what universal coverings are like.

Proof. (idea) We pick a basepoint $x_0 \in X$ for ourselves. Suppose we have a universal covering \tilde{X} . Then this lifts to some \tilde{x}_0 in \tilde{X} . If we have any other point $\tilde{x} \in \tilde{X}$, since \tilde{X} should be path connected, there is a path $\tilde{\alpha} : \tilde{x}_0 \rightarrow \tilde{x}$. If we have another path, then since \tilde{X} is simply connected, then the paths are homotopic. Hence, we can identify each point in \tilde{X} with a path from \tilde{x}_0 , i.e.

$$\{\text{points of } \tilde{X}\} \longleftrightarrow \{\text{paths } \tilde{\alpha} \text{ from } \tilde{x}_0 \in \tilde{X}\} / \simeq.$$

This is not too helpful too, since we are defining \tilde{X} in terms of things in \tilde{X} . However, by path lifting, we know that paths $\tilde{\alpha}$ from \tilde{x}_0 in \tilde{X} biject with paths α from x_0 in X . Also, by homotopy lifting, homotopies of paths in X can be lifted to homotopies of paths in \tilde{X} . So we have

$$\{\text{points of } \tilde{X}\} \longleftrightarrow \{\text{paths } \alpha \text{ from } x_0 \in X\} / \simeq.$$

So we can produce our \tilde{X} by picking a basepoint $x_0 \in X$, and defining

$$\tilde{X} = \{\text{paths } \alpha : I \rightarrow X \text{ such that } \alpha(0) = x_0\} / \simeq.$$

The covering map $p : \tilde{X} \rightarrow X$ is given by $[\alpha] \mapsto \alpha(1)$.

One then has to work hard to define the topology, and then show this is simply connected. \square

3.4 The Galois correspondence

Recall that at the beginning, we wanted to establish a correspondence between covering spaces and fundamental groups. We have already established the result that covering maps are injective on π_1 . Therefore, given a (based) covering space $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$, we can give a subgroup $p_*(\pi_1(\tilde{X}, \tilde{x}_0)) \leq \pi_1(X, x_0)$. It turns out that as long as we define carefully what we mean for based covering spaces to be “the same”, this is a one-to-one correspondence — each subgroup corresponds to a covering space.


We can have the following table of correspondence:

Covering spaces		Fundamental group
(Based) covering spaces	\longleftrightarrow	Subgroups of π_1
Number of sheets	\longleftrightarrow	Index
Universal covers	\longleftrightarrow	Trivial subgroup

We now want to look at some of these correspondences.

Recall that we have shown that $\pi_1(X, x_0)$ acts on $p^{-1}(x_0)$. However, this is not a too interesting action, since $p^{-1}(x_0)$ is a discrete group with no structure. Having groups acting on a cube is fun because the cube has some structure. So we want something more “rich” for $\pi_1(X, x_0)$ to act on.

We note that we can make $\pi_1(X, x_0)$ “act on” the universal cover. How? Recall that in the torus example, each item in the fundamental group corresponds to translating the whole universal covering by some amount. In general, a point on \tilde{X} can be thought of as a path α on X starting from x_0 . Then it is easy to make a loop $\gamma : x_0 \rightsquigarrow x_0$ act on this: use the composition $\gamma \cdot \alpha : [\gamma] \cdot [\alpha] = [\gamma \cdot \alpha]$.



We will use this idea and return to the initial issue of making subgroups correspond with covering spaces. We want to show that this is surjective — every subgroup arises from some cover. We want to say “For any subgroup $H \leq \pi_1(X, x_0)$, there is a based covering map $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$ such that $p_*\pi_1(\tilde{X}, \tilde{x}_0) = H$ ”. Except, this cannot possibly be true, since by taking the trivial subgroup, this would imply that there is a universal covering for every space. So we need some additional assumptions.

Proposition. Let X be a path connected, locally path connected and semi-locally simply connected space. For any subgroup $H \leq \pi_1(X, x_0)$, there is a based covering map $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$ such that $p_*\pi_1(\tilde{X}, \tilde{x}_0) = H$.

Proof. Since X is path connected, locally path connected and semi-locally simply connected space, let \tilde{X} be a universal covering. We have an intermediate group H such that $\pi_1(\tilde{X}, \tilde{x}_0) = 1 \leq H \leq \pi_1(X, x_0)$. How can we obtain a corresponding covering space?

Note that if we have \tilde{X} and we want to recover X , we can quotient \tilde{X} by the action of $\pi_1(X, x_0)$. Since $\pi_1(X, x_0)$ acts on \tilde{X} , so does $H \leq \pi_1(X, x_0)$. Now we can define our covering space by taking quotients. We define \sim_H on \tilde{X} to be the orbit relation for the action of H , i.e. $\tilde{x} \sim_H \tilde{y}$ if there is some $h \in H$ such that $\tilde{y} = h\tilde{x}$. We then let \tilde{X} be the quotient space \tilde{X}/\sim_H .

We can now do the messy algebra to show that this is the covering space we want. \square

We have just showed that every subgroup comes from some covering space, i.e. the map from the set of covering spaces to the subgroups of π_1 is surjective. Now we want to prove injectivity. To do so, we need a generalization of the homotopy lifting lemma.

Suppose we have path-connected spaces (Y, y_0) , (X, x_0) and (\tilde{X}, \tilde{x}_0) , with $f : (Y, y_0) \rightarrow (X, x_0)$ a continuous map, $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$ a covering map. When does a lift of f to $\tilde{f} : (Y, y_0) \rightarrow (\tilde{X}, \tilde{x}_0)$ exist? The answer is given by the lifting criterion.

Lemma (Lifting criterion). Let $p : (\tilde{X}, \tilde{x}_0) \rightarrow (X, x_0)$ be a covering map of path-connected based spaces, and (Y, y_0) a path-connected locally path connected based space. If $f : (Y, y_0) \rightarrow (X, x_0)$ is a continuous map, then there is a (unique) lift $\tilde{f} : (Y, y_0) \rightarrow (\tilde{X}, \tilde{x}_0)$ such that the diagram below commutes (i.e. $p \circ \tilde{f} = f$):

$$\begin{array}{ccc} & & (\tilde{X}, \tilde{x}_0) \\ & \nearrow \tilde{f} & \downarrow p \\ (Y, y_0) & \xrightarrow{f} & (X, x_0) \end{array}$$

if and only if the following condition holds:

$$f_*\pi_1(Y, y_0) \leq p_*\pi_1(\tilde{X}, \tilde{x}_0).$$

Note that uniqueness comes from the uniqueness of lifts. So this lemma is really about existence.

Also, note that the condition holds trivially when Y is simply connected, e.g. when it is an interval (path lifting) or a square (homotopy lifting). So paths and homotopies can always be lifted.

Proof. One direction is easy: if \tilde{f} exists, then $f = p \circ \tilde{f}$. So $f_* = p_* \circ \tilde{f}_*$. So we know that $\text{im } f_* \subseteq \text{im } p_*$. So done.

In the other direction, uniqueness follows from the uniqueness of lifts. So we only need to prove existence. We define \tilde{f} as follows:

Given a $y \in Y$, there is some path $\alpha_y : y_0 \rightsquigarrow y$. Then f maps this to $\beta_y : x_0 \rightsquigarrow f(y)$ in X . By path lifting, this path lifts uniquely to $\tilde{\beta}_y$ in \tilde{X} . Then we set $\tilde{f}(y) = \tilde{\beta}_y(1)$. Note that if \tilde{f} exists, then this *must* be what \tilde{f} sends y to. What we need to show is that this is well-defined.

Suppose we picked a different path $\alpha'_y : y_0 \rightsquigarrow y$. Then this α'_y would have differed from α_y by a loop γ in Y .

Our condition that $f_*\pi_1(Y, y_0) \leq p_*\pi_1(\tilde{X}, \tilde{x}_0)$ says $f \circ \gamma$ is the image of a loop in \tilde{X} . So $\tilde{\beta}_y$ and $\tilde{\beta}'_y$ also differ by a loop in \tilde{X} , and hence have the same end point. So this shows that \tilde{f} is well-defined.

Finally, we show that \tilde{f} is continuous. First, observe that any open set $U \subseteq \tilde{X}$ can be written as a union of \tilde{V} such that $p|_{\tilde{V}} : \tilde{V} \rightarrow p(\tilde{V})$ is a homeomorphism. Thus, it suffices to show that if $p|_{\tilde{V}} : \tilde{V} \rightarrow p(\tilde{V}) = V$ is a homeomorphism, then $\tilde{f}^{-1}(\tilde{V})$ is open.

Let $y \in \tilde{f}^{-1}(\tilde{V})$, and let $x = f(y)$. Since $f^{-1}(V)$ is open and Y is locally path-connected, we can pick an open $W \subseteq f^{-1}(V)$ such that $y \in W$ and W is path connected. We claim that $W \subseteq \tilde{f}^{-1}(\tilde{V})$.

Indeed, if $z \in W$, then we can pick a path γ from y to z . Then f sends this to a path from x to $f(z)$. The lift of this path to \tilde{X} is given by $p|_{\tilde{V}}^{-1}(f(\gamma))$, whose end point is $p|_{\tilde{V}}^{-1}(f(z)) \in \tilde{V}$. So it follows that $\tilde{f}(z) = p|_{\tilde{V}}^{-1}(f(z)) \in \tilde{V}$. \square

Now we prove that every subgroup of π_1 comes from exactly one covering space. What this statement properly means is made precise in the following proposition:

Proposition. Let (X, x) , $(\tilde{X}_1, \tilde{x}_1)$, $(\tilde{X}_2, \tilde{x}_2)$ be path-connected based spaces, and $p_i : (\tilde{X}_i, \tilde{x}_i) \rightarrow (X, x)$ are covering maps. Then we have

$$p_{1*}\pi_1(\tilde{X}_1, \tilde{x}_1) = p_{2*}\pi_1(\tilde{X}_2, \tilde{x}_2)$$

if and only if there is some *homeomorphism* h such that the following diagram commutes:

$$\begin{array}{ccc} (\tilde{X}_1, \tilde{x}_1) & \xrightarrow{\quad h \quad} & (\tilde{X}_2, \tilde{x}_2) \\ & \searrow p_1 \quad \swarrow p_2 & \\ & (X, x) & \end{array}$$

i.e. $p_1 = p_2 \circ h$.

Note that this is a stronger statement than just saying the two covering spaces are homeomorphic. We are saying that we can find nice homeomorphism that works well with the covering map p .

Proof. If such a homeomorphism exists, then clearly the subgroups are equal. If the subgroups are equal, we rotate our diagram a bit:

$$\begin{array}{ccc} & & (\tilde{X}_2, \tilde{x}_2) \\ & \nearrow h=\tilde{p}_1 & \downarrow p_2 \\ (\tilde{X}_1, \tilde{x}_1) & \xrightarrow{\quad p_1 \quad} & (X, x) \end{array}$$

Then $h = \tilde{p}_1$ exists by the lifting criterion. By symmetry, we can get $h^{-1} = \tilde{p}_2$. To show \tilde{p}_2 is indeed the inverse of \tilde{p}_1 , note that $\tilde{p}_2 \circ \tilde{p}_1$ is a lift of $p_2 \circ \tilde{p}_1 = p_1$.

Since $\text{id}_{\tilde{X}_1}$ is also a lift, by the uniqueness of lifts, we know $\tilde{p}_2 \circ \tilde{p}_1$ is the identity map. Similarly, $\tilde{p}_1 \circ \tilde{p}_2$ is also the identity.

$$\begin{array}{ccccc}
 & & & (\tilde{X}_1, \tilde{x}_1) & \\
 & & \nearrow \tilde{p}_2 & \downarrow p_1 & \\
 (\tilde{X}_1, \tilde{x}_1) & \xrightarrow{\tilde{p}_1} & (\tilde{X}_2, \tilde{x}_2) & \xrightarrow{p_2} & (X, x_0) \\
 & \searrow p_1 & & &
 \end{array}$$

□

Now what we would like to do is to forget about the basepoints. What happens when we change base points? Recall that the effect of changing base points is that we will conjugate our group. This doesn't actually change the group itself, but if we are talking about subgroups, conjugation can send a subgroup into a different subgroup. Hence, if we do not specify the basepoint, we don't get a subgroup of π_1 , but a *conjugacy class* of subgroups.

Proposition. Unbased covering spaces correspond to conjugacy classes of subgroups.

4 Some group theory

Algebraic topology is about translating topology into group theory. Unfortunately, you don't know group theory. Well, maybe you do, but not the right group theory. So let's learn group theory!

4.1 Free groups and presentations

Recall that in IA Groups, we defined, say, the dihedral group to be

$$D_{2n} = \langle r, s \mid r^n = s^2 = e, srs = r^{-1} \rangle.$$

What does this expression actually mean? Can we formally assign a meaning to this expression?

We start with a simple case — the free group. This is, in some sense, the “freest” group we can have. These are defined in terms of alphabets and words.

Definition (Alphabet and words). We let $S = \{s_\alpha : \alpha \in \Lambda\}$ be our *alphabets*, and we have an extra set of symbols $S^{-1} = \{s_\alpha^{-1} : \alpha \in \Lambda\}$. We assume that $S \cap S^{-1} = \emptyset$. What do we do with alphabets? We write words with it!

We define S^* to be the set of *words* over $S \cup S^{-1}$, i.e. it contains n -tuples $x_1 \cdots x_n$ for any $0 \leq n < \infty$, where each $x_i \in S \cup S^{-1}$.

Example. Let $S = \{a, b\}$. Then words could be the empty word \emptyset , or a , or $aba^{-1}b^{-1}$, or $aa^{-1}aaaaabbbb$, etc. We are usually lazy and write $aa^{-1}aaaaabbbb$ as $aa^{-1}a^5b^4$.

When we see things like aa^{-1} , we would want to cancel them. This is called *elementary reduction*.

Definition (Elementary reduction). An *elementary reduction* takes a word $us_\alpha s_\alpha^{-1}v$ and gives uv , or turn $us_\alpha^{-1}s_\alpha v$ into uv .

Since each reduction shortens the word, and the word is finite in length, we cannot keep reducing for ever. Eventually, we reach a *reduced* state.

Definition (Reduced word). A word is *reduced* if it does not admit an elementary reduction.

Example. \emptyset , a , $aba^{-1}b^{-1}$ are reduced words, while $aa^{-1}aaaaabbbb$ is not.

Note that there is an inclusion map $S \rightarrow S^*$ that sends the alphabet s_α to the word s_α .

Definition (Free group). The *free group* on the set S , written $F(S)$, is the set of reduced words on S^* together with some operations:

- (i) Multiplication is given by concatenation followed by elementary reduction to get a reduced word. For example, $(aba^{-1}b^{-1}) \cdot (bab) = aba^{-1}b^{-1}bab = ab^2$
- (ii) The identity is the empty word \emptyset .
- (iii) The inverse of $x_1 \cdots x_n$ is $x_n^{-1} \cdots x_1^{-1}$, where, of course, $(s_\alpha^{-1})^{-1} = s_\alpha$.

Note that we have not showed that multiplication is well-defined — we might reduce the same word in different ways and reach two different reduced words. We will show that this is indeed well-defined later, using topology!

Some people like to define the free group in a different way. This is a cleaner way to define the free group without messing with alphabets and words, but is (for most people) less intuitive. This definition also does not make it clear that the free group $F(S)$ of any set S exists. We will state this definition as a lemma.

Lemma. If G is a group and $\phi : S \rightarrow G$ is a set map. Then there exists a unique homomorphism $f : F(S) \rightarrow G$ such that the following diagram commutes:

$$\begin{array}{ccc} & F(S) & \\ \uparrow & \searrow f & \\ S & \xrightarrow{\phi} & G \end{array}$$

where the arrow not labeled is the natural inclusion map that sends s_α (as an alphabet) to s_α (as a word).

Proof. Clearly if f exists, then f must send each s_α to $\phi(s_\alpha)$ and s_α^{-1} to $\phi(s_\alpha)^{-1}$. Then the values of f on all other elements must be determined by

$$f(x_1 \cdots x_n) = f(x_1) \cdots f(x_n)$$

since f is a homomorphism. So if f exists, it must be unique. So it suffices to show that this f is a well-defined homomorphism.

This is well-defined if we define $F(S)$ to be the set of all reduced words, since each reduced word has a unique representation (since it is *defined* to be the representation itself).

To show this is a homomorphism, suppose

$$x = x_1 \cdots x_n a_1 \cdots a_k, \quad y = a_k^{-1} \cdots a_1^{-1} y_1 \cdots y_m,$$

where $y_1 \neq x_n^{-1}$. Then

$$xy = x_1 \cdots x_n y_1 \cdots y_m.$$

Then we can compute

$$\begin{aligned} f(x)f(y) &= (\phi(x_1) \cdots \phi(x_n) \phi(a_1) \cdots \phi(a_k)) (\phi(a_k)^{-1} \cdots \phi(a_1)^{-1} \phi(y_1) \cdots \phi(y_m)) \\ &= \phi(x_1) \cdots \phi(x_n) \cdots \phi(y_1) \cdots \phi(y_m) \\ &= f(xy). \end{aligned}$$

So f is a homomorphism. □

We call this a “universal property” of $F(S)$. We can show that $F(S)$ is the unique group satisfying the conditions of this lemma (up to isomorphism), by taking $G = F(S)$ and using the uniqueness properties.

Definition (Presentation of a group). Let S be a set, and let $R \subseteq F(S)$ be any subset. We denote by $\langle\langle R \rangle\rangle$ the *normal closure* of R , i.e. the smallest normal subgroup of $F(S)$ containing R . This can be given explicitly by

$$\langle\langle R \rangle\rangle = \left\{ \prod_{i=1}^n g_i r_i g_i^{-1} : n \in \mathbb{N}, r_i \in R, g_i \in F(S) \right\}.$$

Then we write

$$\langle S \mid R \rangle = F(S)/\langle\langle R \rangle\rangle.$$

This is just the usual notation we have for groups. For example, we can write

$$D_{2n} = \langle r, s \mid r^n, s^2, sr sr \rangle.$$

Again, we can define this with a universal property.

Lemma. If G is a group and $\phi : S \rightarrow G$ is a set map such that $f(r) = 1$ for all $r \in R$ (i.e. if $r = s_1^{\pm 1} s_2^{\pm 1} \cdots s_m^{\pm 1}$, then $\phi(r) = \phi(s_1)^{\pm 1} \phi(s_2)^{\pm 1} \cdots \phi(s_m)^{\pm 1} = 1$). Then there exists a unique homomorphism $f : \langle S \mid R \rangle \rightarrow G$ such that the following triangle commutes:

$$\begin{array}{ccc} & \langle S \mid R \rangle & \\ \uparrow & \searrow f & \\ S & \xrightarrow{\phi} & G \end{array}$$

Proof is similar to the previous one.

In some sense, this says that all the group $\langle S \mid R \rangle$ does is it satisfies the relations in R , and nothing else.

Example (The stupid canonical presentation). Let G be a group. We can view the group as a set, and hence obtain a free group $F(G)$. There is also an obvious surjection $G \rightarrow G$. Then by the universal property of the free group, there is a surjection $f : F(G) \rightarrow G$. Let $R = \ker f = \langle\langle R \rangle\rangle$. Then $\langle G \mid R \rangle$ is a presentation for G , since the first isomorphism theorem says

$$G \cong F(G)/\ker f.$$

This is a really stupid example. For example, even the simplest non-trivial group $\mathbb{Z}/2$ will be written as a quotient of a free group with two generators. However, this tells us that every group has a presentation.


Example. $\langle a, b \mid b \rangle \cong \langle a \rangle \cong \mathbb{Z}$.

Example. With a bit of work, we can show that $\langle ab \mid ab^{-3}, ba^{-2} \rangle = \mathbb{Z}/5$.

4.2 Another view of free groups


Recall that we have not yet properly defined free groups, since we did not show that multiplication is well-defined. We are now going to do this using topology.

Again let S be a set. For the following illustration, we will just assume $S = \langle a, b \rangle$, but what we will do works for any set S . We define X by




We call this a “rose with 2 petals”. This is a cell complex, with one 0-cell and $|S|$ 1-cells. For each $s \in S$, we have one 1-cell, e_s , and we fix a path $\gamma_s : [0, 1] \rightarrow e_s$ that goes around the 1-cell once. We will call the 0-cells and 1-cells *vertices* and *edges*, and call the whole thing a *graph*.

What’s the universal cover of X ? Since we are just lifting a 1-complex, the result should be a 1-complex, i.e. a graph. Moreover, this graph is connected and simply connected, i.e. it’s a tree. We also know that every vertex in the universal cover is a copy of the vertex in our original graph. So it must have 4 edges attached to it. So it has to look like something this:




In X , we know that at each vertex, there should be an edge labeled a going in; an edge labeled a going out; an edge labeled b going in; an edge labeled b going out. This should be the case in \tilde{X} as well. So \tilde{X} looks like this:



The projection map is then obvious — we send all the vertices in \tilde{X} to $x_0 \in X$, and then the edges according to the labels they have, in a way that respects the direction of the arrow. It is easy to show this is really a covering map.


We are now going to show that this tree “is” the free group. Notice that every word $w \in S^*$ denotes a unique “edge path” in \tilde{X} starting at \tilde{x}_0 , where an edge path is a sequence of oriented edges $\tilde{e}_1, \dots, \tilde{e}_n$ such that the “origin” of \tilde{e}_{i+1} is equal to the “terminus” of \tilde{e}_i .

For example, the following path corresponds to $w = abb^{-1}b^{-1}ba^{-1}b^{-1}$.




We can note a few things:

- (i) \tilde{X} is connected. So for all $\tilde{x} \in p^{-1}(x_0)$, there is an edge-path $\tilde{\gamma} : \tilde{x}_0 \rightarrow \tilde{x}$.
- (ii) If an edge path $\tilde{\gamma}$ fails to be locally injective, we can simplify it. How can an edge fail to be locally injective? It is fine if we just walk along a path, since we are just tracing out a line. So it fails to be locally injective if two consecutive edges in the path are the same edge with opposite orientations:



We can just remove the redundant lines and get



This reminds us of two things — homotopy of paths *and* elementary reduction of words.

- (iii) Each point $\tilde{x} \in p^{-1}(x_0)$ is joined to \tilde{x}_0 by a *unique* locally injective edge-path.
- (iv) For any $w \in S^*$, from (ii), we know that $\tilde{\gamma}$ is locally injective if and only if w is reduced.

We can thus conclude that there are bijections

$$F(S) \longleftarrow p^{-1}(x_0) \longrightarrow \pi_1(X, x_0)$$

that send \tilde{x} to the word $w \in F(S)$ such that $\tilde{\gamma}_w$ is a locally injective edge path to $\tilde{x}_0 \rightsquigarrow \tilde{x}$; and \tilde{x} to $[\gamma] \in \pi_1(X, x_0)$ such that $\tilde{x}_0 \cdot [\gamma] = \tilde{x}$.

So there is a bijection between $F(S)$ and $\pi_1(X, x_0)$. It is easy to see that the operation on $F(S)$ and $\pi_1(X, x_0)$ are the same, since they are just concatenating words or paths. So this bijection identifies the two group structures. So this induces an isomorphism $F(S) \cong \pi_1(X, x_0)$.

4.3 Free products with amalgamation

We managed to compute the fundamental group of the circle. But we want to find the fundamental group of more things. Recall that at the beginning, we defined cell complexes, and said these are the things we want to work with. Cell complexes are formed by gluing things together. So we want to know what happens when we glue things together.

Suppose a space X is constructed from two spaces A, B by gluing (i.e. $X = A \cup B$). We would like to describe $\pi_1(X)$ in terms of $\pi_1(A)$ and $\pi_1(B)$. To understand this, we need to understand how to “glue” groups together.

Definition (Free product). Suppose we have groups $G_1 = \langle S_1 \mid R_1 \rangle, G_2 = \langle S_2 \mid R_2 \rangle$, where we assume $S_1 \cap S_2 = \emptyset$. The *free product* $G_1 * G_2$ is defined to be

$$G_1 * G_2 = \langle S_1 \cup S_2 \mid R_1 \cup R_2 \rangle.$$

This is not a really satisfactory definition. A group can have many different presentations, and it is not clear this is well-defined. However, it is clear that this group exists. Note that there are natural maps $j_i : G_i \rightarrow G_1 * G_2$ that send generators to generators. Then we can have the following universal property of the free product:

Lemma. $G_1 * G_2$ is the group such that for any group K and homomorphisms $\phi_i : G_i \rightarrow K$, there exists a unique homomorphism $f : G_1 * G_2 \rightarrow K$ such that the following diagram commutes:

$$\begin{array}{ccccc}
 & & G_2 & & \\
 & & \downarrow j_2 & \searrow \phi_2 & \\
 G_1 & \xrightarrow{j_1} & G_1 * G_2 & \xrightarrow{f} & K \\
 & \searrow \phi_1 & & \nearrow & \\
 & & & &
 \end{array}$$

Proof. It is immediate from universal property of the definition of presentations. \square

Corollary. The free product is well-defined.

Proof. The conclusion of the universal property can be seen to characterize $G_1 * G_2$ up to isomorphism. \square

Again, we have a definition in terms of a concrete construction of the group, without making it clear this is well-defined; then we have a universal property that makes it clear this is well-defined, but not clear that the object actually exists. Combining the two would give everything we want.

However, this free product is not exactly what we want, since there is little interaction between G_1 and G_2 . In terms of gluing spaces, this corresponds to gluing A and B when $A \cap B$ are trivial (i.e. simply connected). What we really need is the free product with amalgamation, as you might guess have guessed from the title.

Definition (Free product with amalgamation). Suppose we have groups G_1 , G_2 and H , with the following maps:

$$\begin{array}{ccc} H & \xrightarrow{i_2} & G_2 \\ \downarrow i_1 & & \\ G_1 & & \end{array}$$

The *free product with amalgamation* is defined to be

$$G_1 *_H G_2 = G_1 * G_2 / \langle\langle (j_2 \circ i_2(h))^{-1}(j_1 \circ i_1)(h) : h \in H \rangle\rangle.$$

Here we are attempting to identify things “in H ” as the same, but we need to use the maps j_k and i_k to map the things from H to $G_1 * G_2$. So we want to say “for any h , $j_1 \circ i_1(h) = j_2 \circ i_2(h)$ ”, or $(j_2 \circ i_2(h))^{-1}(j_1 \circ i_1)(h) = e$. So we quotient by the (normal closure) of these things.

We have the universal property

Lemma. $G_1 *_H G_2$ is the group such that for any group K and homomorphisms $\phi_i : G_i \rightarrow K$, there exists a unique homomorphism $G_1 *_H G_2 \rightarrow K$ such that the following diagram commutes:

$$\begin{array}{ccccc} H & \xrightarrow{i_2} & G_2 & & \\ \downarrow i_1 & & \downarrow j_2 & \searrow \phi_2 & \\ G_1 & \xrightarrow{j_1} & G_1 *_H G_2 & \xrightarrow{\quad} & K \\ & \searrow \phi_1 & \nearrow f & & \end{array}$$


This is the language we will need to compute fundamental groups. These definitions will (hopefully) become more concrete as we see more examples.

5 Seifert-van Kampen theorem

5.1 Seifert-van Kampen theorem

The Seifert-van Kampen theorem is the theorem that tells us what happens when we glue spaces together.

Here we let $X = A \cup B$, where $A, B, A \cap B$ are path-connected.



We pick a basepoint $x_0 \in A \cap B$ for convenience. Since we like diagrams, we can write this as a commutative diagram:

$$\begin{array}{ccc} A \cap B & \hookrightarrow & B \\ \downarrow & & \downarrow \\ A & \hookrightarrow & X \end{array}$$

where all arrows are inclusion (i.e. injective) maps. We can consider what happens when we take the fundamental groups of each space. Then we have the induced homomorphisms

$$\begin{array}{ccc} \pi_1(A \cap B, x_0) & \longrightarrow & \pi_1(B, x_0) \\ \downarrow & & \downarrow \\ \pi_1(A, x_0) & \longrightarrow & \pi_1(X, x_0) \end{array}$$

We might guess that $\pi_1(X, x_0)$ is just the free product with amalgamation

$$\pi_1(X, x_0) = \pi_1(A, x_0) \underset{\pi_1(A \cap B, x_0)}{*} \pi_1(B, x_0).$$

The Seifert-van Kampen theorem says that, under mild hypotheses, this guess is correct.

Theorem (Seifert-van Kampen theorem). Let A, B be open subspaces of X such that $X = A \cup B$, and $A, B, A \cap B$ are path-connected. Then for any $x_0 \in A \cap B$, we have

$$\pi_1(X, x_0) = \pi_1(A, x_0) \underset{\pi_1(A \cap B, x_0)}{*} \pi_1(B, x_0).$$

Note that by the universal property of the free product with amalgamation, we by definition know that there is a unique map $\pi_1(A, x_0) \underset{\pi_1(A \cap B, x_0)}{*} \pi_1(B, x_0) \rightarrow \pi_1(X, x_0)$. The theorem asserts that *this* map is an isomorphism.


Proof is omitted because time is short.

Example. Consider a higher-dimensional sphere $S^n = \{\mathbf{v} \in \mathbb{R}^{n+1} : |\mathbf{v}| = 1\}$ for $n \geq 2$. We want to find $\pi_1(S^n)$.

The idea is to write S^n as a union of two open sets. We let $n = \mathbf{e}_1 \in S^n \subseteq \mathbb{R}^{n+1}$ be the North pole, and $s = -\mathbf{e}_1$ be the South pole. We let $A = S^n \setminus \{n\}$,

and $B = S^n \setminus \{s\}$. By stereographic projection, we know that $A, B \cong \mathbb{R}^n$. The hard part is to understand the intersection.

To do so, we can draw a cylinder $S^{n-1} \times (-1, 1)$, and project our $A \cap B$ onto the cylinder. We can similarly project the cylinder onto $A \cap B$. So $A \cap B \cong S^{n-1} \times (-1, 1) \simeq S^{n-1}$, since $(-1, 1)$ is contractible.



We can now apply the Seifert-van Kampen theorem. Note that this works only if S^{n-1} is path-connected, i.e. $n \geq 2$. Then this tells us that

$$\pi_1(S^n) \cong \pi_1(\mathbb{R}^n) *_{\pi_1(S^{n-1})} \pi_1(\mathbb{R}^n) \cong 1 *_{\pi_1(S^{n-1})} 1$$

It is easy to see this is the trivial group. We can see this directly from the universal property of the amalgamated free product, or note that it is the quotient of $1 * 1$, which is 1.

So for $n \geq 2$, $\pi_1(S^n) \cong 1$.

We have found yet another simply connected space. However, this is unlike our previous examples. Our previous spaces were simply connected because they were contractible. However, we will later see that S^n is not contractible. So this is genuinely a new, interesting example.

Why did we go through all this work to prove that $\pi_1(S^n) \cong 1$? It feels like we can just prove this directly — pick a point that is not in the curve as the North pole, project stereographically to \mathbb{R}^n , and contract it there. However, the problem is that space-filling curves exist. We cannot guarantee that we can pick a point not on the curve! It is indeed possible to prove directly that given any curve on S^n (with $n \geq 1$), we can deform it slightly so that it is no longer surjective. Then the above proof strategy works. However, using Seifert-van Kampen is a much neater.

Example (\mathbb{RP}^n). Recall that $\mathbb{RP}^n \cong S^n / \{\pm \text{id}\}$, and the quotient map $S^n \rightarrow \mathbb{RP}^n$ is a covering map. Now that we have proved that S^n is simply connected, we know that S^n is a universal cover of \mathbb{RP}^n .


For any $x_0 \in \mathbb{RP}^n$, we have a bijection

$$\pi_1(\mathbb{RP}^n, x_0) \leftrightarrow p^{-1}(x_0).$$


Since $p^{-1}(x_0)$ has two elements by definition, we know that $|\pi_1(\mathbb{RP}^n, x_0)| = 2$. So $\pi_1(\mathbb{RP}^n, x_0) = \mathbb{Z}/2$.

You will prove a generalization of this in example sheet 2.


Example (Wedge of two circles). We are going to consider the operation of *wedging*. Suppose we have two topological spaces, and we want to join them together. The natural way to join them is to take the disjoint union. What if we have based spaces? If we have (X, x_0) and (Y, y_0) , we cannot just take the disjoint union, since we will lose our base point. What we do is take the *wedge sum*, where we take the disjoint union and then identify the base points:



Suppose we take the wedge sum of two circles $S^1 \vee S^1$. We would like to pick A, B to be each of the circles, but we cannot, since A and B have to be open. So we take slightly more, and get the following:



Each of A and B now look like this:



We see that both A and B retract to the circle. So $\pi_1(A) \cong \pi_1(B) \cong \mathbb{Z}$, while $A \cap B$ is a cross, which retracts to a point. So $\pi_1(A \cap B) = 1$.

Hence by the Seifert-van Kampen theorem, we get

$$\pi_1(S^1 \vee S^1) = \pi_1(A, x_0) *_{\pi_1(A \cap B, x_0)} \pi_1(B, x_0) \cong \mathbb{Z} * \mathbb{Z} \cong \mathbb{Z} * \mathbb{Z} \cong F_2,$$

where F_2 is just $F(S)$ for $|S| = 2$. We can see that $\mathbb{Z} * \mathbb{Z} \cong F_2$ by showing that they satisfy the same universal property.

Note that we have already figured this out when we studied the free group, where we realized F_2 is the fundamental group of this thing.

More generally, as long as x_0, y_0 in X and Y are “reasonable”, $\pi_1(X \vee Y) \cong \pi_1(X) * \pi_1(Y)$.

Next, we would exhibit some nice examples of the covering spaces of $S^1 \wedge S^1$, i.e. the “rose with 2 petals”.

Recall that $\pi_1(S^1 \wedge S^1, x_0) \cong F_2 \cong \langle a, b \rangle$.


Example. Consider the map $\phi : F_2 \rightarrow \mathbb{Z}/3$ which sends $a \mapsto 1, b \mapsto 1$. Note that 1 is *not* the identity, since this is an abelian group, and the identity is 0. This exists since the universal property tells us we just have to say where the generators go, and the map exists (and is unique).

Now $\ker \phi$ is a subgroup of F_2 . So there is a based covering space of $S^1 \wedge S^1$ corresponding to it, say, \tilde{X} . Let’s work out what it is.


First, we want to know how many sheets it has, i.e. how many copies of x_0 we have. There are three, since we know that the number of sheets is the index of the subgroup, and the index of $\ker \phi$ is $|\mathbb{Z}/3| = 3$ by the first isomorphism theorem.




Let’s try to lift the loop a at \tilde{x}_0 . Since $a \notin \ker \phi = \pi_1(\tilde{X}, \tilde{x}_0)$, a does not lift to a loop. So it goes to another vertex.




Similarly, $a^2 \notin \ker \phi = \pi_1(\tilde{X}, \tilde{x}_0)$. So we get the following



Since $a^3 \in \ker \phi$, we get a loop labelled a .



Note that $ab^{-1} \in \ker \phi$. So ab^{-1} gives a loop. So b goes in the same direction at b :



This is our covering space.


This is a fun game to play at home:

- (i) Pick a group G (finite groups are recommended).
- (ii) Then pick $\alpha, \beta \in G$ and let $\phi : F_2 \rightarrow G$ send $a \rightarrow \alpha, b \rightarrow \beta$.
- (iii) Compute the covering space corresponding to ϕ .

5.2 The effect on π_1 of attaching cells

We have started the course by talking about cell complexes, but largely ignored them afterwards. Finally, we are getting back to them.

The process of attaching cells is as follows: we start with a space X , get a function $S^{n-1} \rightarrow X$, and attach D^n to X by gluing along the image of f to get $X \cup_f D^n$:



Since we are attaching stuff, we can use the Seifert-van Kampen theorem to analyse this.

Theorem. If $n \geq 3$, then $\pi_1(X \cup_f D^n) \cong \pi_1(X)$. More precisely, the map $\pi_1(X, x_0) \rightarrow \pi_1(X \cup_f D^n, x_0)$ induced by inclusion is an isomorphism, where x_0 is a point on the image of f .

This tells us that attaching a high-dimensional disk does not do anything to the fundamental group.

Proof. Again, the difficulty of applying Seifert-van Kampen theorem is that we need to work with open sets.

Let $0 \in D^n$ be any point in the interior of D^n . We let $A = X \cup_f (D^n \setminus \{0\})$. Note that $D^n \setminus \{0\}$ deformation retracts to the boundary S^{n-1} . So A deformation retracts to X . Let $B = \mathring{D}$, the interior of D^n . Then

$$A \cap B = \mathring{D}^n \setminus 0 \cong S^{n-1} \times (-1, 1)$$

We cannot use y_0 as our basepoint, since this point is not in $A \cap B$. Instead, pick an arbitrary $y_1 \in A \cap B$. Since D^n is path connected, we have a path $\gamma : y_1 \rightsquigarrow y_0$, and we can use this to recover the fundamental groups based y_0 .

Now Seifert-van Kampen theorem says

$$\pi_1(X \cup_f D^n, y_1) \cong \pi_1(A, y_1) \underset{\pi_1(A \cap B, y_1)}{*} \pi_1(B, y_1).$$

Since B is just a disk, and $A \cap B$ is simply connected ($n \geq 3$ implies S^{n-1} is simply connected), their fundamental groups are trivial. So we get

$$\pi_1(X \cup_f D^n, y_1) \cong \pi_1(A, y_1).$$

We can now use γ to change base points from y_1 to y_0 . So

$$\pi_1(X \cup_f D^n, y_0) \cong \pi_1(A, y_0) \cong \pi_1(X, y_0).$$

□

The more interesting case is when we have smaller dimensions.

Theorem. If $n = 2$, then the natural map $\pi_1(X, X_0) \rightarrow \pi_1(X \cup_f D^n, x_0)$ is *surjective*, and the kernel is $\langle\langle [f] \rangle\rangle$. Note that this statement makes sense, since S^{n-1} is a circle, and $f : S^{n-1} \rightarrow X$ is a loop in X .

This is what we would expect, since if we attach a disk onto the loop given by f , this loop just dies.

Proof. As before, we get

$$\pi_1(X \cup_f D^n, y_1) \cong \pi_1(A, y_1) \underset{\pi_1(A \cap B, y_1)}{*} \pi_1(B, y_1).$$

Again, B is contractible, and $\pi_1(B, y_1) \cong 1$. However, $\pi_1(A \cap B, y_1) \cong \mathbb{Z}$. Since $\pi_1(A \cap B, y_1)$ is just (homotopic to) the loop induced by f , it follows that

$$\pi_1(A, y_1) \underset{\pi_1(A \cap B, y_1)}{*} 1 = (\pi_1(A, y_1) * 1) / \langle\langle \pi_1(A \cap B, y_1) \rangle\rangle \cong \pi_1(X, x_0) / \langle\langle f \rangle\rangle.$$


□

In summary, we have

$$\pi_1(X \cup_f D^n) = \begin{cases} \pi_1(X) & n \geq 3 \\ \pi_1(X) / \langle\langle f \rangle\rangle & n = 2 \end{cases}$$

This is a useful result, since this is how we build up cell complexes. If we want to compute the fundamental groups, we can just work up to the two-cells, and know that the higher-dimensional cells do not have any effect. Moreover, whenever X is a cell complex, we should be able to write down the presentation of $\pi_1(X)$.

Example. Let X be the 2-torus. Possibly, our favorite picture of the torus is (not a doughnut):




This is already a description of the torus as a cell complex!


We start with our zero complex $X^{(0)}$:

•

We then add our 1-cells to get $X^{(1)}$:



We now glue our square to the cell complex to get $X = X^{(2)}$:



matching up the colors and directions of arrows.

So we have our torus as a cell complex. What is its fundamental group? There are many ways we can do this computation, but this time we want to do it as a cell complex.

We start with $X^{(0)}$. This is a single point. So its fundamental group is $\pi_1(X^{(0)}) = 1$.

When we add our two 1-cells, we get $\pi_1(X^{(1)}) = F_2 \cong \langle a, b \rangle$.

Finally, to get $\pi_1(X)$, we have to quotient out by the boundary of our square, which is just $aba^{-1}b^{-1}$. So we have

$$\pi_1(X^{(2)}) = F_2 / \langle \langle aba^{-1}b^{-1} \rangle \rangle = \langle a, b \mid aba^{-1}b^{-1} \rangle \cong \mathbb{Z}^2.$$

We have the last congruence since we have two generators, and then we make they commute by quotienting the commutator out.

This procedure can be reversed — given a presentation of a group, we can just add the right edges and squares to produce a cell complex with that presentation.

Corollary. For any (finite) group presentation $\langle S \mid R \rangle$, there exist a (finite) cell complex (of dimension 2) X such that $\pi_1(X) \cong \langle S \mid R \rangle$.

There really isn't anything that requires that finiteness in this proof, but finiteness makes us feel more comfortable.

Proof. Let $S = \{a_1, \dots, a_m\}$ and $R = \{r_1, \dots, r_n\}$. We start with a single point, and get our $X^{(1)}$ by adding a loop about the point for each $a_i \in S$. We then get our 2-cells e_j^2 for $j = 1, \dots, n$, and attaching them to $X^{(1)}$ by $f_i : S^1 \rightarrow X^{(1)}$ given by a based loop representing $r_i \in F(S)$. \square

Since all groups have presentations, this tells us that all groups are fundamental groups of some spaces (at least those with finite presentations).

5.3 A refinement of the Seifert-van Kampen theorem

We are going to make a refinement of the theorem so that we don't have to worry about that openness problem. We first start with a definition.


Definition (Neighbourhood deformation retract). A subset $A \subseteq X$ is a *neighbourhood deformation retract* if there is an open set $A \subseteq U \subseteq X$ such that A is a strong deformation retract of U , i.e. there exists a retract $r : U \rightarrow A$ and $r \simeq \text{id}_U \text{ rel } A$.

This is something that is true most of the time, in sufficiently sane spaces.

Example. If Y is a subcomplex of a cell complex, then Y is a neighbourhood deformation retract.

Theorem. Let X be a space, $A, B \subseteq X$ closed subspaces. Suppose that A , B and $A \cap B$ are path connected, and $A \cap B$ is a neighbourhood deformation retract of A and B . Then for any $x_0 \in A \cap B$.

$$\pi_1(X, x_0) = \pi_1(A, x_0) *_{\pi_1(A \cap B, x_0)} \pi_1(B, x_0).$$



This is just like Seifert-van Kampen theorem, but usually easier to apply, since we no longer have to “fatten up” our A and B to make them open.

Proof. Pick open neighbourhoods $A \cap B \subseteq U \subseteq A$ and $A \cap B \subseteq V \subseteq B$ that strongly deformation retracts to $A \cap B$. Let U be such that U retracts to $A \cap B$. Since U retracts to A , it follows that U is path connected since path-connectedness is preserved by homotopies.

Let $A' = A \cup V$ and $B' = B \cup U$. Since $A' = (X \setminus B) \cup V$, and $B' = (X \setminus A) \cup U$, it follows that A' and B' are open.


Since U and V retract to $A \cap B$, we know $A' \simeq A$ and $B' \simeq B$. Also, $A' \cap B' = (A \cup V) \cap (B \cup U) = U \cup V \simeq A \cap B$. In particular, it is path connected. So by Seifert van-Kampen, we get

$$\pi_1(A \cup B) = \pi_1(A', x_0) *_{\pi_1(A' \cap B', x_0)} \pi_1(B', x_0) = \pi_1(A, x_0) *_{\pi_1(A \cap B, x_0)} \pi_1(B, x_0).$$

□

This is basically what we’ve done all the time when we enlarge our A and B to become open.

Example. Let $X = S^1 \wedge S^1$, the rose with two petals. Let $A, B \cong S^1$ be the circles.



Then since $\{x_0\} = A \cap B$ is a neighbourhood deformation retract of A and B , we know that

$$\pi_1 X \cong \pi_1 S^1 * \pi_1 S^1.$$

5.4 The fundamental group of all surfaces

We have found that the torus has fundamental group \mathbb{Z}^2 , but we already knew this, since the torus is just $S^1 \times S^1$, and the fundamental group of a product is the product of the fundamental groups, as you have shown in the example sheet. So we want to look at something more interesting. We look at *all* surfaces.

We start by defining what a surface is. It is surprisingly difficult to get mathematicians to agree on how we can define a surface. Here we adopt the following definition:


Definition (Surface). A *surface* is a Hausdorff topological space such that every point has a neighbourhood U that is homeomorphic to \mathbb{R}^2 .

Some people like \mathbb{C} more than \mathbb{R} , and sometimes they put \mathbb{C}^2 instead of \mathbb{R}^2 in the definition, which is confusing, since that would have two complex dimensions and hence *four* real dimensions. Needless to say, the actual surfaces will also be different and have different homotopy groups. We will just focus on surfaces with two real dimensions.


To find the fundamental group of all surfaces, we rely on the following theorem that tells us what surfaces there are.

Theorem (Classification of compact surfaces). If X is a compact surface, then X is homeomorphic to a space in one of the following two families:

- (i) The *orientable surface of genus g* , Σ_g includes the following (please excuse my drawing skills):




A more formal definition of this family is the following: we start with a 2-sphere, and remove a few discs from it and get $S^2 \setminus \bigcup_{i=1}^g D^2$. Then we take g tori with an open disc removed, and attach them to the circles.




- (ii) The *non-orientable surface of genus n* , $E_n = \{\mathbb{RP}^2, K, \dots\}$ (where K is the Klein bottle). This has a similar construction as above: we start with the sphere S^2 , make a few holes, and then glue Möbius strips to them.

It would be nice to be able to compute fundamental groups of these guys. To do so, we need to write them as polygons with identification.

Example. To obtain surface of genus two, written Σ_2 , we start with what we had for a torus:



If we just wanted a torus, we are done (after closing the loop), but now we want a surface with genus 2, so we add another torus:



To visualize how this works, imagine cutting this apart along the dashed line. This would give two tori with a hole, where the boundary of the holes are just the dashed line. Then gluing back the dashed lines would give back our orientable surface with genus 2.



In general, to produce Σ_g , we produce a polygon with $4g$ sides. Then we get

$$\pi_1 \Sigma_g = \langle a_1, b_1, \dots, a_g, b_g \mid a_1 b_1 a_1^{-1} b_1^{-1} \dots a_g b_g a_g^{-1} b_g^{-1} \rangle.$$

We do we care? The classification theorem tells us that each surface is homeomorphic to *some* of these orientable and non-orientable surfaces, but it doesn't tell us there is no overlap. It might be that $\Sigma_6 \cong \Sigma_{241}$, via some weird homeomorphism that destroys some holes.

However, this result lets us know that all these orientable surfaces are genuinely different. While it is difficult to stare at this fundamental group and say that $\pi_1 \Sigma_g \not\cong \pi_1 \Sigma_{g'}$ for $g \neq g'$, we can perform a little trick. We can take the abelianization of the group $\pi_1 \Sigma_g$, where we further quotient by all commutators. Then the abelianized fundamental group of Σ_g will simply be \mathbb{Z}^{2g} . These are clearly distinct for different values of g . So all these surfaces are distinct. Moreover, they are not even homotopy equivalent.

The fundamental groups of the non-orientable surfaces is left as an exercise for the reader.

6 Simplicial complexes

So far, we have taken a space X , and assigned some things to it. The first was easy — $\pi_0(X)$. It was easy to calculate and understand. We then spent a lot of time talking about $\pi_1(X)$. What are they good for? A question we motivated ourselves with was to prove $\mathbb{R}^m \cong \mathbb{R}^n$ implies $n = m$. π_0 was pretty good for the case when $n = 1$. If $\mathbb{R}^m \cong \mathbb{R}$, then we would have $\mathbb{R}^m \setminus \{0\} \cong \mathbb{R} \setminus \{0\} \simeq S^0$. We know that $|\pi_0(S^0)| = 2$, while $|\pi_0(\mathbb{R}^m \setminus \{0\})| = 1$ for $m \neq 1$. This is just a fancy way of saying that $\mathbb{R} \setminus \{0\}$ is disconnected while $\mathbb{R}^m \setminus \{0\}$ is not for $m \neq 1$.

We can just add 1 to n , and add 1 to our subscript. If $\mathbb{R}^m \cong \mathbb{R}^2$, then we have $\mathbb{R}^m \setminus \{0\} \cong \mathbb{R}^2 \setminus \{0\} \simeq S^1$. We know that $\pi_1(S^1) \cong \mathbb{Z}$, while $\pi_1(\mathbb{R}^m \setminus \{0\}) \cong \pi_1(S^{m-1}) \cong 1$ unless $m = 2$.

The obvious thing to do is to create some $\pi_n(X)$. But as you noticed, π_1 took us quite a long time to define, and was *really* hard to compute. As we would expect, this only gets harder as we get to higher dimensions. This is indeed possible, but will only be done in Part III courses.

The problem is that π_n works with groups, and groups are hard. There are too many groups out there. We want to do some easier algebra, and a good choice is linear algebra. Linear algebra is easy. In the rest of the course, we will have things like $H_0(X)$ and $H_1(X)$, instead of π_n , which are more closely related to linear algebra.

Another way to motivate the abandoning of groups is as follows: recall last time we saw that if X is a finite cell, reasonably explicitly defined, then we can write down a presentation $\langle S \mid R \rangle$ for $\pi_1(X)$. This sounds easy, except that we don't understand presentations. In fact there is a theorem that says there is no algorithm that decides if a particular group presentation is actually the trivial group.

So even though we can compute the fundamental group in terms of presentations, this is not necessarily helpful. On the other hand, linear algebra is easy. Computers can do them. So we will move on to working with linear algebra instead.

This is where *homology theory* comes in. It takes a while for us to define it, but after we finish developing the machinery, things are easy.

6.1 Simplicial complexes

There are many ways of formulating homology theory, and these are all equivalent at least for sufficiently sane spaces (such as cell complexes). In this course, we will be using *simplicial homology*, which is relatively more intuitive and can be computed directly. The drawback is that we will have to restrict to a particular kind of space, known as *simplicial complexes*. This is not a very serious restriction *per se*, since many spaces like spheres are indeed simplicial complexes. However, the definition of simplicial homology is based on exactly how we view our space as a simplicial complex, and it will take us quite a lot of work to show that the simplicial homology is indeed a property of the space itself, and not how we represent it as a simplicial complex.


We now start by defining simplicial complexes, and developing some general theory of simplicial complexes that will become useful later on.

Definition (Affine independence). A finite set of points $\{a_1, \dots, a_n\} \subseteq \mathbb{R}^m$ is


affinely independent iff

$$\sum_{i=1}^n t_i a_i = 0 \text{ with } \sum_{i=1}^n t_i = 0 \Leftrightarrow t_i = 0 \text{ for all } i.$$

Example. When $n = 3$, the following points are affinely independent:



The following are not:



The proper way of understanding this definition is via the following lemma:

Lemma. $a_0, \dots, a_n \in \mathbb{R}^m$ are affinely independent if and only if $a_1 - a_0, \dots, a_n - a_0$ are linearly independent.

Alternatively, $n + 1$ affinely independent points span an n -dimensional thing.

Proof. Suppose a_0, \dots, a_n are affinely independent. Suppose

$$\sum_{i=1}^n \lambda_i (a_i - a_0) = 0.$$

Then we can rewrite this as

$$\left(-\sum_{i=1}^n \lambda_i \right) a_0 + \lambda_1 a_1 + \dots + \lambda_n a_n = 0.$$

Now the sum of the coefficients is 0. So affine independence implies that all coefficients are 0. So $a_1 - a_0, \dots, a_n - a_0$ are linearly independent.

On the other hand, suppose $a_1 - a_0, \dots, a_n - a_0$ are linearly independent. Now suppose

$$\sum_{i=0}^n t_i a_i = 0, \quad \sum_{i=0}^n t_i = 0.$$

Then we can write

$$t_0 = -\sum_{i=1}^n t_i.$$

Then the first equation reads

$$0 = \left(-\sum_{i=1}^n t_i \right) a_0 + t_1 a_1 + \dots + t_n a_n = \sum_{i=1}^n t_i (a_i - a_0).$$

So linear independence implies all $t_i = 0$. □

The relevance is that these can be used to define simplexes (which are simple, as opposed to complexes).

Definition (n -simplex). An n -simplex is the convex hull of $(n + 1)$ affinely independent points $a_0, \dots, a_n \in \mathbb{R}^m$, i.e. the set

$$\sigma = \langle a_0, \dots, a_n \rangle = \left\{ \sum_{i=0}^n t_i a_i : \sum_{i=0}^n t_i = 1, t_i \geq 0 \right\}.$$

The points a_0, \dots, a_n are the *vertices*, and are said to *span* σ . The $(n + 1)$ tuples (t_0, \dots, t_n) are called the *barycentric coordinates* for the point $\sum t_i a_i$.


Example. When $n = 0$, then our 0-simplex is just a point:




When $n = 1$, then we get a line:



When $n = 2$, we get a triangle:



When $n = 3$, we get a tetrahedron:




The key motivation of this is that simplices are determined by their vertices. Unlike arbitrary subspaces of \mathbb{R}^n , they can be specified by a finite amount of data. We can also easily extract the faces of the simplices.

Definition (Face, boundary and interior). A *face* of a simplex is a subset (or subsimplex) spanned by a subset of the vertices. The *boundary* is the union of the proper faces, and the *interior* is the complement of the boundary.

The boundary of σ is usually denoted by $\partial\sigma$, while the interior is denoted by $\mathring{\sigma}$, and we write $\tau \leq \sigma$ when τ is a face of σ .

In particular, the interior of a vertex is the vertex itself. Note that this notion of interior and boundary is distinct from the topological notion of boundary.

Example. The *standard n -simplex* is spanned by the basis vectors $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ in \mathbb{R}^{n+1} . For example, when $n = 2$, we get the following:




We will now glue simplices together to build *complexes*, or *simplicial complexes*.

Definition. A (geometric) *simplicial complex* is a finite set K of simplices in \mathbb{R}^n such that


- (i) If $\sigma \in K$ and τ is a face of σ , then $\tau \in K$.
- (ii) If $\sigma, \tau \in K$, then $\sigma \cap \tau$ is either empty or a face of both σ and τ .

Definition (Vertices). The *vertices* of K are the zero simplices of K , denoted V_K .

Example. This is a simplicial complex:



These are not:



Technically, a simplicial complex is defined to be a set of simplices, which is just collections of points. It is not a subspace of \mathbb{R}^n . Hence we have the following definition:

Definition (Polyhedron). The *polyhedron* defined by K is the union of the simplices in K , and denoted by $|K|$.

We make this distinction because distinct simplicial complexes may have the same polyhedron, such as the following:



Just as in cell complexes, we can define things like dimensions.

Definition (Dimension and skeleton). The *dimension* of K is the highest dimension of a simplex of K . The d -*skeleton* $K^{(d)}$ of K is the union of the n -simplices in K for $n \leq d$.

Note that since these are finite and live inside \mathbb{R}^n , we know that $|K|$ is always compact and Hausdorff.

Usually, when we are given a space, say S^n , it is not defined to be a simplicial complex. We can “make” them simplicial complexes by a *triangulation*.


Definition (Triangulation). A *triangulation* of a space X is a homeomorphism $h : |K| \rightarrow X$, where K is some simplicial complex.

Example. Let σ be the standard n -simplex. The boundary $\partial\sigma$ is homeomorphic to S^{n-1} (e.g. the boundary of a (solid) triangle is the boundary of the triangle, which is also a circle). This is called the *simplicial* $(n-1)$ *sphere*.

We can also triangulate our S^n in a different way:

Example. In \mathbb{R}^{n+1} , consider the simplices $\langle \pm \mathbf{e}_0, \dots, \pm \mathbf{e}_n \rangle$ for each possible combination of signs. So we have 2^{n+1} simplices in total. Then their union defines a simplicial complex K , and

$$|K| \cong S^n.$$



The nice thing about this triangulation is that the simplicial complex is invariant under the antipodal map. So not only can we think of this as a triangulation of the sphere, but a triangulation of \mathbb{RP}^n as well.


As always, we don't just look at objects themselves, but also maps between them.

Definition (Simplicial map). A *simplicial map* $f : K \rightarrow L$ is a function $f : V_K \rightarrow V_L$ such that if $\langle a_0, \dots, a_n \rangle$ is a simplex in K , then $\{f(a_0), \dots, f(a_n)\}$ spans a simplex of L .

The nice thing about simplicial maps is that we only have to specify where the vertices go, and there are only finitely many vertices. So we can completely specify a simplicial map by writing down a finite amount of information.

It is important to note that we say $\{f(a_0), \dots, f(a_n)\}$ as a set span a simplex of L . In particular, they are allowed to have repeats.

Example. Suppose we have the standard 2-simplex K as follows:




The following does *not* define a simplicial map because $\langle a_1, a_2 \rangle$ is a simplex in K , but $\{f(a_1), f(a_2)\}$ does not span a simplex:

$$\begin{array}{ccc} f(a_0), f(a_1) & & f(a_2) \\ \bullet & & \bullet \end{array}$$

On the other hand, the following *is* a simplicial map, because now $\{f(a_1), f(a_2)\}$ spans a simplex, and note that $\{f(a_0), f(a_1), f(a_2)\}$ also spans a 1-simplex because we are treating the collection of three vertices as a set, and not a simplex.

$$\begin{array}{ccc} f(a_0), f(a_1) & & f(a_2) \\ \bullet & \text{---} & \bullet \end{array}$$

Finally, we can also do the following map:



The following lemma is obvious, but we will need it later on.

Lemma. If K is a simplicial complex, then every point $x \in |K|$ lies in the interior of a *unique* simplex.

As we said, simplicial maps are nice, but they are not exactly what we want. We want to have maps between *spaces*.

Lemma. A simplicial map $f : K \rightarrow L$ induces a continuous map $|f| : |K| \rightarrow |L|$, and furthermore, we have

$$|f \circ g| = |f| \circ |g|.$$

There is an obvious way to define this map. We know how to map vertices, and then just extend everything linearly.

Proof. For any point in a simplex $\sigma = \langle a_0, \dots, a_n \rangle$, we define

$$|f| \left(\sum_{i=0}^n t_i a_i \right) = \sum_{i=0}^n t_i f(a_i).$$

The result is in L because $\{f(a_i)\}$ spans a simplex. It is not difficult to see this is well-defined when the point lies on the boundary of a simplex. This is clearly continuous on σ , and is hence continuous on $|K|$ by the gluing lemma.

The final property is obviously true by definition. \square

6.2 Simplicial approximation

This is all very well, but we are really interested in *continuous maps*. So given a continuous map f , we would like to find a related simplicial map g . In particular, we want to find a simplicial map g that “approximates” f in some sense. The definition we will write down is slightly awkward, but it turns out this is the most useful definition.

Definition (Open star and link). Let $x \in |K|$. The *open star* of x is the union of all the interiors of the simplices that contain x , i.e.

$$\text{St}_K(x) = \bigcup_{x \in \sigma \in K} \overset{\circ}{\sigma}.$$

The *link* of x , written $\text{Lk}_K(x)$, is the union of all those simplices that do not contain x , but are faces of a simplex that do contain x .

Definition (Simplicial approximation). Let $f : |K| \rightarrow |L|$ be a continuous map between the polyhedra. A function $g : V_K \rightarrow V_L$ is a *simplicial approximation* to f if for all $v \in V_K$,

$$f(\text{St}_K(v)) \subseteq \text{St}_L(g(v)). \quad (*)$$

The following lemma tells us why this is a good definition:

Lemma. If $f : |K| \rightarrow |L|$ is a map between polyhedra, and $g : V_K \rightarrow V_L$ is a simplicial approximation to f , then g is a simplicial map, and $|g| \simeq f$. Furthermore, if f is already simplicial on some subcomplex $M \subseteq K$, then we get $g|_M = f|_M$, and the homotopy can be made rel M .

Proof. First we want to check g is really a simplicial map if it satisfies $(*)$. Let $\sigma = \langle a_0, \dots, a_n \rangle$ span a simplex in K . We want to show that $\{g(a_0), \dots, g(a_n)\}$ spans a simplex in L .

Pick arbitrary $x \in \overset{\circ}{\sigma}$. Since σ contains each a_i , we know that $x \in \text{St}_K(a_i)$ for all i . Hence we know that

$$f(x) \in \bigcap_{i=0}^n f(\text{St}_K(a_i)) \subseteq \bigcap_{i=0}^n \text{St}_L(g(a_i)).$$

Hence we know that there is one simplex, say, τ that contains all $g(a_i)$ whose interior contains $f(x)$. Since each $g(a_i)$ is a vertex in L , each $g(a_i)$ must be a vertex of τ . So they span a face of τ , as required.


We now want to prove that $|g| \simeq f$. We let $H : |K| \times I \rightarrow |L| \subseteq \mathbb{R}^m$ be defined by

$$(x, t) \mapsto t|g|(x) + (1 - t)f(x).$$


This is clearly continuous. So we need to check that $\text{im } H \subseteq |L|$. But we know that both $|g|(x)$ and $f(x)$ live in τ and τ is convex. It thus follows that $H(x \times I) \subseteq \tau \subseteq |L|$.

To prove the last part, it suffices to show that every simplicial approximation to a simplicial map must be the map itself. Then the homotopy is rel M by construction above. This is easily seen to be true — if g is a simplicial approximation to f , then $f(v) \in f(\text{St}_K(v)) \subseteq \text{St}_L(g(v))$. Since $f(v)$ is a vertex and $g(v)$ is the only vertex in $\text{St}_L(g(v))$, we must have $f(v) = g(v)$. So done. \square


What's the best thing we might hope for at this point? It would be great if every map is homotopic to a simplicial map. Is this possible? Let's take a nice example. Let's consider the following K :



How many homotopy classes of continuous maps are there $K \rightarrow K$? Countably many, one for each winding number. However, there can only be at most $3^3 = 27$ simplicial maps. The problem is that we don't have enough vertices to realize all those interesting maps. The idea is to refine our simplicial complexes. Suppose we have the following simplex:



What we do is to add a point in the center of each simplex, and join them up:



This is known as the barycentric subdivision. After we subdivide it once, we can realize more homotopy classes of maps. We will show that for any map, as long as we are willing to barycentrically subdivide the simplex many times, we can find a simplicial approximation to it.

Definition (Barycenter). The *barycenter* of $\sigma = \langle a_0, \dots, a_n \rangle$ is

$$\hat{\sigma} = \sum_{i=0}^n \frac{1}{n+1} a_i.$$

Definition (Barycentric subdivision). The *(first) barycentric subdivision* K' of K is the simplicial complex:

$$K' = \{ \langle \hat{\sigma}_0, \dots, \hat{\sigma}_n \rangle : \sigma_i \in K \text{ and } \sigma_0 < \sigma_1 < \dots < \sigma_n \}.$$

If you stare at this long enough, you will realize this is exactly what we have drawn above.

The r th barycentric subdivision $K^{(r)}$ is defined inductively as the barycentric subdivision of the $r-1$ th barycentric subdivision, i.e.

$$K^{(r)} = (K^{(r-1)})'.$$

Proposition. $|K| = |K'|$ and K' really is a simplicial complex.

Proof. Too boring to be included in lectures. □

We now have a slight problem. Even though $|K'|$ and $|K|$ are equal, the identity map from $|K'|$ to $|K|$ is not a simplicial map.

To solve this problem, we can choose any function $K \rightarrow V_K$ by $\sigma \mapsto v_\sigma$ with $v_\sigma \in \sigma$, i.e. a function that sends any simplex to any of its vertices. Then we can define $g : K' \rightarrow K$ by sending $\hat{\sigma} \mapsto v_\sigma$. Then this is a simplicial map, and indeed a simplicial approximation to the identity map $|K'| \rightarrow |K|$.

The key theorem is that as long as we are willing to perform barycentric subdivisions, then we can always find a simplicial approximation.

Theorem (Simplicial approximation theorem). Let K and L be simplicial complexes, and $f : |K| \rightarrow |L|$ a continuous map. Then there exists an r and a simplicial map $g : K^{(r)} \rightarrow L$ such that g is a simplicial approximation of f . Furthermore, if f is already simplicial on $M \subseteq K$, then we can choose g such that $|g|_M = f|_M$.

The first thing we have to figure out is how far we are going to subdivide. To do this, we want to quantify how “fine” our subdivisions are.

Definition (Mesh). Let K be a simplicial complex. The *mesh* of K is

$$\mu(K) = \max\{\|v_0 - v_1\| : \langle v_0, v_1 \rangle \in K\}.$$

We have the following lemma that tells us how large our mesh is:

Lemma. Let $\dim K \leq n$, then

$$\mu(K^{(r)}) = \left(\frac{n}{n+1}\right)^r \mu(K).$$

The key point is that as $r \rightarrow \infty$, the mesh goes to zero. So indeed we can make our barycentric subdivisions finer and finer. The proof is purely technical and omitted.

Now we can prove the simplicial approximation theorem.

Proof of simplicial approximation theorem. Suppose we are given the map $f : |K| \rightarrow |L|$. We have a natural cover of $|L|$, namely the open stars of all vertices. We can use f to pull these back to $|K|$ to obtain a cover of $|K|$:

$$\{f^{-1}(\text{St}_L(w)) : w \in V_L\}.$$

The idea is to barycentrically subdivide our K such that each open star of K is contained in one of these things.

By the Lebesgue number lemma, there exists some δ , the *Lebesgue number* of the cover, such that for each $x \in |K|$, $B_\delta(x)$ is contained in *some* element of the cover. By the previous lemma, there is an r such that $\mu(K^{(r)}) < \delta$.

Now since the mesh $\mu(K^{(r)})$ is the smallest distance between any two vertices, the radius of every open star $\text{St}_{K^{(r)}}(x)$ is at most $\mu(K^{(r)})$. Hence it follows that $\text{St}_{K^{(r)}}(x) \subseteq B_\delta(x)$ for all vertices $x \in V_{K^{(r)}}$. Therefore, for all $x \in V_{K^{(r)}}$, there is some $w \in V_L$ such that

$$\text{St}_{K^{(r)}}(x) \subseteq B_\delta(x) \subseteq f^{-1}(\text{St}_L(w)).$$

Therefore defining $g(x) = w$, we get

$$f(\text{St}_{K^{(r)}}(x)) \subseteq \text{St}_L(g(x)).$$

So g is a simplicial approximation of f .

The last part follows from the observation that if f is a simplicial map, then it maps vertices to vertices. So we can pick $g(v) = f(v)$. \square

7 Simplicial homology

7.1 Simplicial homology

For now, we will forget about simplicial approximations and related fluff, but just note that it is fine to assume everything can be considered to be simplicial. Instead, we are going to use this framework to move on and define some new invariants of simplicial complexes K , known as $H_n(K)$. These are analogous to π_0, π_1, \dots , but only use linear algebra in the definitions, and is thus much simpler. The drawback, however, is that the definitions are slightly less intuitive at first sight.

Despite saying “linear algebra”, we won’t be working with vector spaces most of the time. Instead, we will be using abelian groups, which really should be thought of as \mathbb{Z} -modules. At the end, we will come up with an analogous theory using \mathbb{Q} -modules, i.e. \mathbb{Q} -vector spaces, and most of the theory will carry over. Using \mathbb{Q} makes some of our work easier, but as a result we would have lost some information. In the most general case, we can replace \mathbb{Z} with any abelian group, but we will not be considering any of these in this course.

Recall that we defined an n -simplex as the collection of $n + 1$ vertices that span a simplex. As a simplex, when we permute the vertices, we still get the same simplex. What we want to do is to remember the orientation of the simplex. In particular, we want think of the simplices (a, b) and (b, a) as different:


$$a \bullet \longrightarrow \bullet b \qquad a \bullet \longleftarrow \bullet b$$

Hence we define *oriented simplices*.

Definition (Oriented n -simplex). An *oriented n -simplex* in a simplicial complex K is an $n + 1$ -tuple (a_0, \dots, a_n) of vertices $a_i \in V_k$ such that $\langle a_0, \dots, a_n \rangle \in K$, where we think of two $n + 1$ tuples (a_0, \dots, a_n) and $(a_{\pi(0)}, \dots, a_{\pi(n)})$ as the same *oriented* simplex if $\pi \in S^n$ is an *even* permutation.

We often denote an oriented simplex as σ , and then $\bar{\sigma}$ denotes the same simplex with the opposite orientation.

Example. As oriented 2-simplices, (v_0, v_1, v_2) and (v_1, v_2, v_0) are equal, but they are different from (v_2, v_1, v_0) . We can imagine the two different orientations of the simplices as follows:



One and two dimensions are the dimensions where we can easily visualize the orientation. This is substantially harder in higher dimensions, and often we just work with the definition instead.

Definition (Chain group $C_n(K)$). Let K be a simplicial complex. For each $n \geq 0$, we define $C_n(K)$ as follows:

Let $\{\sigma_1, \dots, \sigma_\ell\}$ be the set of n simplices of k . For each i , choose an orientation on σ_i . That is, choose an order for the vertices (up to an even

permutation). This choice is not important, but we need to make it. Now when we say σ_i , we mean the oriented simplex with this particular orientation.

Now let $C_n(K)$ be the free abelian group with basis $\{\sigma_1, \dots, \sigma_\ell\}$, i.e. $C_n(K) \cong \mathbb{Z}^\ell$. So an element in $C_n(K)$ might look like

$$\sigma_3 - 7\sigma_1 + 52\sigma_{64} - 28\sigma_{1000000}.$$

In other words, an element of $C_n(K)$ is just a formal sum of n -simplices.


For convenience, we define $C_\ell(K) = 0$ for $\ell < 0$. This will save us from making exceptions for $n = 0$ cases later.

For each oriented simplex, we identify $-\sigma_i$ with $\bar{\sigma}_i$, at least when $n \geq 1$.

In this definition, we have to choose a particular orientation for each of our simplices. If you don't like making arbitrary choices, we could instead define $C_n(K)$ as some quotient, but it is slightly more complicated.

Note that if there are no n simplices (e.g. when $n = -1$), we can still meaningfully talk about $C_n(K)$, but it's just 0.

Example. We can think of elements in the chain group $C_1(X)$ as “paths” in X . For example, we might have the following simplex:



Then the path $v_0 \rightsquigarrow v_1 \rightsquigarrow v_2 \rightsquigarrow v_0 \rightsquigarrow v_1$ around the left triangle is represented by the member

$$\sigma_1 - \sigma_2 + \sigma_3 + \sigma_1 = 2\sigma_1 - \sigma_2 + \sigma_3.$$

Of course, with this setup, we can do more random things, like adding 57 copies of σ_6 to it, and this is also allowed. So we could think of these as disjoint union of paths instead.

When defining fundamental groups, we had homotopies that allows us to “move around”. Somehow, we would like to say two of these paths are “equivalent” under certain conditions. To do so, we need to define the *boundary homomorphisms*.

Definition (Boundary homomorphisms). We define *boundary homomorphisms*

$$d_n : C_n(K) \rightarrow C_{n-1}(K)$$

by


$$(a_0, \dots, a_n) \mapsto \sum_{i=0}^n (-1)^i (a_0, \dots, \hat{a}_i, \dots, a_n),$$

where $(a_0, \dots, \hat{a}_i, \dots, a_n) = (a_0, \dots, a_{i-1}, a_{i+1}, \dots, a_n)$ is the simplex with a_i removed.

This means we remove each vertex in turn and add them up. This is clear if we draw some pictures in low dimensions:

$$v_0 \longrightarrow v_1 \quad \longmapsto \quad \begin{array}{c} -v_0 \\ \bullet \end{array} \quad \begin{array}{c} v_1 \\ \bullet \end{array}$$

If we take a triangle, we get



An important property of the boundary map is that the boundary of a boundary is empty:

Lemma. $d_{n-1} \circ d_n = 0$.

In other words, $\text{im } d_{n+1} \subseteq \ker d_n$.

Proof. This just involves expanding the definition and working through the mess. \square

With this in mind, we will define the homology groups as follows:

Definition (Simplicial homology group $H_n(K)$). The n th simplicial homology group $H_n(K)$ is defined as


$$H_n(K) = \frac{\ker d_n}{\text{im } d_{n+1}}.$$

This is a nice, clean definition, but what does this mean geometrically?

Somehow, $H_k(K)$ describes all the “ k -dimensional holes” in $|K|$. Since we are going to draw pictures, we are going to start with the easy case of $k = 1$. Our $H_1(K)$ is made from the kernel of d_1 and the image of d_2 . First, we give these things names.

Definition (Boundary and cycles). The k -cycles of K is $\ker d_k$. The k -boundaries is $\text{im } d_{k+1}$.

Suppose we have some $c \in \ker d_k$. In other words, $dc = 0$. If we interpret c as a “path”, if it has no boundary, then it represents some sort of loop, i.e. a cycle. For example, if we have the following cycle:



We have


$$c = (e_0, e_1) + (e_1, e_2) + (e_2, e_0).$$

We can then compute the boundary as

$$dc = (e_1 - e_0) + (e_2 - e_1) + (e_0 - e_2) = 0.$$


So this c is indeed a cycle.

Now if $c \in \text{im } d_2$, then $c = db$ for some 2-chain b , i.e. c is the boundary of some two-dimensional thing. This is why we call this a 1-boundary. For example, suppose we have our cycle as above, but is itself a boundary of a 2-chain.



We see that a cycle that have a boundary has been “filled in”. Hence the “holes” are, roughly, the cycles that haven’t been filled in. Hence we define the homology group as the cycles quotiented by the boundaries, and we interpret this as k -dimensional “holes”.

Example. Let K be the standard simplicial 1-sphere, ie, we have the following in \mathbb{R}^3 .



Our simplices are thus

$$K = \{\langle e_0 \rangle, \langle e_1 \rangle, \langle e_2 \rangle, \langle e_0, e_1 \rangle, \langle e_1, e_2 \rangle, \langle e_2, e_0 \rangle\}.$$

Our chain groups are

$$C_0(K) = \langle (e_0), (e_1), (e_2) \rangle \cong \mathbb{Z}^3$$

$$C_1(K) = \langle (e_0, e_1), (e_1, e_2), (e_2, e_0) \rangle \cong \mathbb{Z}^3.$$

All other chain groups are zero. Note that our notation is slightly confusing here, since the brackets $\langle \cdot \rangle$ can mean the simplex spanned by the vertices, or the group generated by certain elements. However, you are probably clueful enough to distinguish the two use cases.

Hence, the only non-zero boundary map is

$$d_1 : C_1(K) \rightarrow C_0(K).$$

We can write down its matrix with respect to the given basis.

$$\begin{pmatrix} -1 & 0 & 1 \\ 1 & -1 & 0 \\ 0 & 1 & -1 \end{pmatrix}$$

We have now everything we need to know about the homology groups, and we just need to do some linear algebra to figure out the image and kernel, and thus the homology groups. We have

$$H_0(K) = \frac{\ker(d_0 : C_0(K) \rightarrow C_{-1}(K))}{\operatorname{im}(d_1 : C_1(K) \rightarrow C_0(K))} \cong \frac{C_0(K)}{\operatorname{im} d_1} \cong \frac{\mathbb{Z}^3}{\operatorname{im} d_1}.$$

After doing some row operations with our matrix, we see that the image of d_1 is a two-dimensional subspace generated by the image of two of the edges. Hence we have

$$H_0(K) = \mathbb{Z}.$$

What does this $H_0(\mathbb{Z})$ represent? We initially said that $H_k(K)$ should represent the k -dimensional holes, but when $k = 0$, this is simpler. As for π_0 , H_0 just represents the path components of K . We interpret this to mean K has one path component. In general, if K has r path components, then we expect $H_0(K)$ to be \mathbb{Z}^r .

Similarly, we have

$$H_1(K) = \frac{\ker d_1}{\operatorname{im} d_2} \cong \ker d_1.$$

It is easy to see that in fact we have


$$\ker d_1 = \langle (e_0, e_1) + (e_1, e_2) + (e_2, e_0) \rangle \cong \mathbb{Z}.$$

So we also have

$$H_1(K) \cong \mathbb{Z}.$$

We see that this $H_1(K)$ is generated by precisely the single loop in the triangle. The fact that $H_1(K)$ is non-trivial means that we do indeed have a hole in the middle of the circle.

Example. Let L be the standard 2-simplex (and all its faces) in \mathbb{R}^3 .



Now our chain groups are

$$\begin{aligned} C_0(L) &= C_0(K) \cong \mathbb{Z}^3 \cong \langle (e_0), (e_1), (e_2) \rangle \\ C_1(L) &= C_1(K) \cong \mathbb{Z}^3 \cong \langle (e_0, e_1), (e_1, e_2), (e_2, e_0) \rangle \\ C_2(L) &\cong \mathbb{Z} = \langle (e_0, e_1, e_2) \rangle. \end{aligned}$$

Since d_1 is the same as before, the only new interesting boundary map is d_2 . We compute

$$d_2((e_0, e_1, e_2)) = (e_0, e_1) + (e_1, e_2) + (e_2, e_0).$$

We know that $H_0(L)$ depends only on d_0 and d_1 , which are the same as for K . So

$$H_0(L) \cong \mathbb{Z}.$$

Again, the interpretation of this is that L is path-connected. The first homology group is

$$H_1(L) = \frac{\ker d_1}{\operatorname{im} d_2} = \frac{\langle (e_0, e_1) + (e_1, e_2) + (e_2, e_0) \rangle}{\langle (e_0, e_1) + (e_1, e_2) + (e_2, e_0) \rangle} \cong 0.$$

This precisely illustrates the fact that the “hole” is now filled in in L .

Finally, we have

$$H_2(L) = \frac{\ker d_2}{\operatorname{im} d_3} = \ker d_2 \cong 0.$$

This is zero since there aren’t any two-dimensional holes in L .

We have hopefully gained some intuition on what these homology groups mean. We are now going to spend a lot of time developing formalism.

7.2 Some homological algebra

We will develop some formalism to help us compute homology groups $H_k(K)$ in lots of examples. Firstly, we axiomatize the setup we had above.

Definition (Chain complex and differentials). A *chain complex* C_\bullet is a sequence of abelian groups C_0, C_1, C_2, \dots equipped with maps $d_n : C_n \rightarrow C_{n-1}$ such that $d_{n-1} \circ d_n = 0$ for all n . We call these maps the *differentials* of C_\bullet .

$$0 \xleftarrow{d_0} C_0 \xleftarrow{d_1} C_1 \xleftarrow{d_2} C_2 \xleftarrow{d_3} \dots$$

Whenever we have some of these things, we can define homology in exactly how we defined them for simplicial complexes.

Definition (Cycles and boundaries). The *n -cycles* is

$$Z_n(C) = \ker d_n.$$

The *n -boundaries* is

$$B_n(C) = \operatorname{im} d_n.$$

Definition (Homology group). The *n -th homology group* of C_\bullet is defined to be

$$H_n(C) = \frac{\ker d_n}{\ker d_{n+1}} = \frac{Z_n(C)}{B_n(C)}.$$

In mathematics, when we have objects, we don’t just talk about the objects themselves, but also functions between them. Suppose we have two chain complexes. For the sake of simplicity, we write the chain maps of both as d_n .

In general, we want to have maps between these two sequences. This would correspond to having a map $f_i : C_i \rightarrow D_i$ for each i , satisfying some commutativity relations.

Definition (Chain map). A chain map $f_{\bullet} : C_{\bullet} \rightarrow D_{\bullet}$ is a sequence of homomorphisms $f_n : C_n \rightarrow D_n$ such that

$$f_{n-1} \circ d_n = d_n \circ f_n$$

for all n . In other words, the following diagram commutes:

$$\begin{array}{ccccccc} 0 & \xleftarrow{d_0} & C_0 & \xleftarrow{d_1} & C_1 & \xleftarrow{d_2} & C_2 \xleftarrow{d_3} \cdots \\ & & \downarrow f_0 & & \downarrow f_1 & & \downarrow f_2 \\ 0 & \xleftarrow{d_0} & D_0 & \xleftarrow{d_1} & D_1 & \xleftarrow{d_2} & D_2 \xleftarrow{d_3} \cdots \end{array}$$

We want to have homotopies. So far, chain complexes seem to be rather rigid, but homotopies themselves are rather floppy. How can we define homotopies for chain complexes? It turns out we can have a completely algebraic definition for *chain homotopies*.

Definition (Chain homotopy). A *chain homotopy* between chain maps $f_{\bullet}, g_{\bullet} : C_{\bullet} \rightarrow D_{\bullet}$ is a sequence of homomorphisms $h_n : C_n \rightarrow D_{n+1}$ such that

$$g_n - f_n = d_{n+1} \circ h_n + h_{n-1} \circ d_n.$$

We write $f_{\bullet} \simeq g_{\bullet}$ if there is a chain homotopy between f_{\bullet} and g_{\bullet} .


The arrows can be put in the following diagram, which is *not* commutative:

$$\begin{array}{ccccc} C_{n-1} & & \xleftarrow{d_n} & C_n & \\ & \searrow h_{n-1} & & \downarrow g_n, f_n & \swarrow h_n \\ & & D_n & \xleftarrow{d_{n+1}} & D_{n+1} \end{array}$$

The intuition behind this definition is as follows: suppose $C_{\bullet} = C_{\bullet}(K)$ and $D_{\bullet} = C_{\bullet}(L)$ for K, L simplicial complexes, and f_{\bullet} and g_{\bullet} are “induced” by simplicial maps $f, g : K \rightarrow L$. How can we detect if f and g are homotopic via the homotopy groups?

Suppose $H : |K| \times I \rightarrow |L|$ is a homotopy from f to g . We further suppose that H actually comes from a simplicial map $K \times I \rightarrow L$ (we’ll skim over the technical issue of how we can make $K \times I$ a simplicial complex. Instead, you are supposed to figure this out yourself in example sheet 3).

Let σ be an n -simplex of K , and here is a picture of $H(\sigma \times I)$:



Let

$$h_n(\sigma) = H(\sigma \times I).$$

We think of this as an $(n+1)$ -chain. What is its boundary? We've got the vertical sides plus the top and bottom. The bottom should just be $f(\sigma)$, and the top is just $g(\sigma)$, since H is an homotopy from f to g . How about the sides? They are what we get when we pull the boundary $\partial\sigma$ up with the homotopy, i.e. $H(\partial\sigma \times I) = h_{n-1} \circ d_n(\sigma)$. Now note that $f(\sigma)$ and $g(\sigma)$ have opposite orientations, so we get the result

$$d_{n+1} \circ h_n(\sigma) = h_{n-1} \circ d_n(\sigma) + g_n(\sigma) - f_n(\sigma).$$

Rearranging and dropping the σ s, we get

$$d_{n+1} \circ h_n - h_{n-1} \circ d_n = g_n - f_n.$$

This looks remarkably like our definition for chain homotopies of maps, with the signs a bit wrong. So in reality, we will have to fiddle with the sign of h_n a bit to get it right, but you get the idea.

Lemma. A chain map $f_* : C_* \rightarrow D_*$ induces a homomorphism:

$$\begin{aligned} f_* : H_n(C) &\rightarrow H_n(D) \\ [c] &\mapsto [f(c)] \end{aligned}$$

Furthermore, if f_* and g_* are chain homotopic, then $f_* = g_*$.

Proof. Since the homology groups are defined as the cycles quotiented by the boundaries, to show that f_* defines a homomorphism, we need to show f sends cycles to cycles and boundaries to boundaries. This is an easy check. If $d_n(\sigma) = 0$, then

$$d_n(f_n(\sigma)) = f_n(d_n(\sigma)) = f_n(0) = 0.$$

So $f_n(\sigma) \in Z_n(D)$.

Similarly, if σ is a boundary, say $\sigma = d_n(\tau)$, then

$$f_n(\sigma) = f_n(d_n(\tau)) = d_n(f_n(\tau)).$$

So $f_n(\sigma)$ is a boundary. It thus follows that f_* is well-defined.

Now suppose h_n is a chain homotopy between f and g . For any $c \in Z_n(C)$, we have

$$g_n(c) - f_n(c) = d_{n+1} \circ h_n(c) + h_{n-1} \circ d_n(c)$$

Since $c \in Z_n(C)$, we know that $d_n(c) = 0$. So

$$g_n(c) - f_n(c) = d_{n+1} \circ h_n(c) \in B_n(D).$$

Hence $g_n(c) - f_n(c)$ differ by a boundary. So $[g_n(c)] - [f_n(c)] = 0$ in $H_n(D)$, i.e. $f_*(c) = g_*(c)$. \square

The following statements are easy to check:

Proposition.

- (i) Being chain-homotopic is an equivalence relation of chain maps.
- (ii) If $a_0 : A_0 \rightarrow C_0$ is a chain map and $f_* \simeq g_*$, then $f_* \circ a_* \simeq g_* \circ a_*$.

(iii) If $f : C \rightarrow D$ and $g : D \rightarrow A$ are chain maps, then

$$g_* \circ f_* = (f \circ g)_*.$$

(iv) $(\text{id}_C)_* = \text{id}_{H_*(C)}$.

The last two statements can be summarized in fancy languages by saying that H_n is a functor.

Definition (Chain homotopy equivalence). Chain complexes C and D are *chain homotopy equivalent* if there exists $f : C \rightarrow D$ and $g : D \rightarrow C$ such that

$$f \circ g \simeq \text{id}_D, \quad g \circ f \simeq \text{id}_C.$$

We should think of this in exactly the same way as we think of homotopy equivalences of spaces. The chain complexes themselves are not necessarily the same, but the induced homology groups will be.

Lemma. Let $f : C \rightarrow D$ is a *chain homotopy equivalence*, then $f_* : H_n(C) \rightarrow H_n(D)$ is an isomorphism for all n .

Proof. Let g be the homotopy inverse. Since $f \circ g \simeq \text{id}_D$, we know $f_* \circ g_* = \text{id}_{H_*(D)}$. Similarly, $g_* \circ f_* = \text{id}_{H_*(C)}$. So we get isomorphisms between $H_n(C)$ and $H_n(D)$. \square

7.3 Homology calculations

We'll get back to topology and compute some homologies.

Here, K is always a simplicial complex, and $C_* = C_*(K)$.

Lemma. Let $f : K \rightarrow L$ be a simplicial map. Then f induces a chain map $f_* : C_*(K) \rightarrow C_*(L)$. Hence it also induces $f_* : H_n(K) \rightarrow H_n(L)$.

Proof. This is fairly obvious, except that simplicial maps are allowed to “squash” simplices, so f might send an n -simplex to an $(n-1)$ -simplex, which is not in $D_n(L)$. We solve these problem by just killing these troublesome simplices.

Let σ be an oriented n -simplex in K , corresponding to a basis element of $C_n(K)$. Then we define

$$f_n(\sigma) = \begin{cases} f(\sigma) & f(\sigma) \text{ is an } n\text{-simplex} \\ 0 & f(\sigma) \text{ is a } k\text{-simplex for } k < n \end{cases}.$$

More precisely, if $\sigma = (a_0, \dots, a_n)$, then


$$f_n(\sigma) = \begin{cases} (f(a_0), \dots, f(a_n)) & (f(a_0), \dots, f(a_n)) \text{ spans an } n\text{-simplex} \\ 0 & \text{otherwise} \end{cases}.$$

We then extend f_n linearly to obtain $f_n : C_n(K) \rightarrow C_n(L)$.

It is immediate from this that this satisfies the chain map condition, i.e. f_* commutes with the boundary operators. \square

Definition (Cone). A simplicial complex is a *cone* if, for some $v_0 \in V_k$,

$$|K| = \text{St}_K(v_0) \cup \text{Lk}_K(v_0).$$



We see that a cone ought to be contractible — we can just squash it to the point v_0 . This is what the next lemma tells us.

Lemma. If K is a cone with cone point v_0 , then inclusion $i : \{v_0\} \rightarrow |K|$ induces a chain homotopy equivalence $i_* : C_n(\{v_0\}) \rightarrow C_n(K)$. Therefore

$$H_n(K) = \begin{cases} \mathbb{Z} & n = 0 \\ 0 & n > 0 \end{cases}$$

The homotopy inverse to i_* is given by r_* , where $r : k \rightarrow v_0$ is the only map. It is clear that $r_* \circ i_* = \text{id}$, and we need to show that $i_* \circ r_* \simeq \text{id}$. This chain homotopy can be defined by $h_n : C_n(K) \rightarrow C_{n+1}(K)$, where h_n associates to any simplex σ in $\text{Lk}_K(v_0)$ to the simplex spanned by σ and v_0 . Details are left to the reader.

Corollary. If Δ^n is the standard n simplex, and L consists of Δ^n and all its faces, then

$$H_k(L) = \begin{cases} \mathbb{Z} & k = 0 \\ 0 & k > 0 \end{cases}$$

Proof. K is a cone (on any vertex). □

What we would really like is an example of a non-trivial homology groups. Moreover, we want them in higher dimensions, and not just the examples we got for fundamental groups. An obvious candidate is the standard n sphere.

Corollary. Let K be the standard $(n-1)$ sphere (i.e. the proper faces of L from above). Then for $n \geq 2$, we have

$$H_k(K) = \begin{cases} \mathbb{Z} & k = 0 \\ 0 & 0 < k < n-1 \\ \mathbb{Z} & k = n-1 \end{cases}$$

Proof. We write down the chain groups for K and L .

$$\begin{array}{ccccccc} 0 & \longleftarrow & C_0(L) & \xleftarrow{d_1^L} & C_1(L) & \longleftarrow & \cdots \xleftarrow{d_{n-1}^L} C_{n-1}(L) \xleftarrow{d_n^L} C_n(L) \\ & & \parallel & & \parallel & & \parallel \\ 0 & \longleftarrow & C_0(K) & \xleftarrow{d_1^K} & C_1(K) & \longleftarrow & \cdots \xleftarrow{d_{n-1}^K} C_{n-1}(K) \longleftarrow C_n(K) = 0 \end{array}$$

For $k < n - 1$, we have $C_k(K) = C_k(L)$ and $C_{k+1}(K) = C_{k+1}(L)$. Also, the boundary maps are equal. So

$$H_k(K) = H_k(L) = 0.$$

We now need to compute

$$H_{n-1}(K) = \ker d_{n-1}^K = \ker d_{n-1}^L = \operatorname{im} d_n^L.$$

We get the last equality since

$$\frac{\ker d_{n-1}^L}{\operatorname{im} d_n^L} = H_{n-1}(L) = 0.$$

We also know that $C_n(L)$ is generated by just one simplex (e_0, \dots, e_n) . So $C_n(L) \cong \mathbb{Z}$. Also d_n^L is injective since it does not kill the generator (e_0, \dots, e_n) . So

$$H_{n-1}(K) \cong \operatorname{im} d_n^L \cong \mathbb{Z}.$$

□

This is very exciting, because at last, we have a suggestion for a non-trivial invariant of S^{n-1} for $n > 2$. We say this is just a “suggestion”, since the simplicial homology is defined for simplicial complexes, and we are not guaranteed that if we put a different simplicial complex on S^{n-1} , we will get the same homology groups. So the major technical obstacle we still need to overcome is to see that H_k are invariants of the polyhedron $|K|$, not just K , and similarly for maps. But this will take some time.

We will quickly say something we’ve alluded to already:

Lemma (Interpretation of H_0). $H_0(K) \cong \mathbb{Z}^d$, where d is the number of path components of K .


Proof. Let K be our simplicial complex and $v, w \in V_k$. We note that by definition, v, w represent the same homology class in $H_0(K)$ if and only if there is some c such that $d_1 c = w - v$. The requirement that $d_1 c = w - v$ is equivalent to saying c is a path from v to w . So $[v] = [w]$ if and only if v and w are in the same path component of K . □

Most of the time, we only care about path-connected spaces, so $H_0(K) \cong \mathbb{Z}$.

7.4 Mayer-Vietoris sequence

The Mayer-Vietoris sequence is exactly like the Seifert-van Kampen theorem for fundamental groups, which tells us what happens when we glue two spaces together.

Suppose we have a space $K = M \cup N$, where $M \cap N$ are path-connected.



We will learn how to compute the homology of the union $M \cup N$ in terms of those of M, N and $M \cap N$.

Recall that to state the Seifert-van Kampen theorem, we need to learn some new group-theoretic notion of free products with amalgamation. The situation is somewhat similar here. We will need to learn some algebra in order to state the Mayer-Vietoris sequence. These are known as *exact sequences*.

Definition (Exact sequence). A pair of homomorphisms of abelian groups

$$A \xrightarrow{f} B \xrightarrow{g} C$$

is *exact (at B)* if

$$\text{im } f = \ker g.$$

A collection of homomorphisms

$$\cdots \xrightarrow{f_{i-1}} A_i \xrightarrow{f_i} A_{i+1} \xrightarrow{f_{i+1}} A_{i+2} \xrightarrow{f_{i+2}} \cdots$$

is *exact at A_i* if

$$\ker f_i = \text{im } f_{i-1}.$$

We say it is *exact* if it is exact at every A_i .

Recall that if we seen something similar before. When we defined the chain complexes, we had $d^2 = 0$, i.e. $\text{im } d \subseteq \ker d$. Here we are requiring exact equivalence, which is something even better.

Algebraically, we can think of an exact sequence as chain complexes with trivial homology groups. Alternatively, we see the homology groups as measuring the failure of a sequence to be exact.

There is a particular type of exact sequences that is important.

Definition (Short exact sequence). A *short exact sequence* is an exact sequence of the form

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow 0$$

What does this mean?

- The kernel of f is equal to the image of the zero map, i.e. $\{0\}$. So f is injective.
- The image of g is the kernel of the zero map, which is everything. So g is surjective.
- $\text{im } f = \ker g$.

Since we like chain complexes, we can produce short exact sequences of chain complexes.

Definition (Short exact sequence of chain complexes). A *short exact sequence of chain complexes* is a pair of *chain maps* i_\bullet and j_\bullet .

$$0 \longrightarrow A_\bullet \xrightarrow{i_\bullet} B_\bullet \xrightarrow{j_\bullet} C_\bullet \longrightarrow 0$$

such that for each k ,

$$0 \longrightarrow A_k \xrightarrow{i_k} B_k \xrightarrow{j_k} C_k \longrightarrow 0$$

is exact.

Note that by requiring the maps to be chain maps, we imply that i_k and j_k commute with the boundary maps the chain complexes.

The reason why we care about short exact sequences of chain complexes (despite it having such a long name) is the following result:

Theorem (Snake lemma). If we have a short exact sequence of complexes

$$0 \longrightarrow A. \xrightarrow{i.} B. \xrightarrow{j.} C. \longrightarrow 0$$

then a miracle happens to their homology groups. In particular, there is a long exact sequence (i.e. an exact sequence that is not short)

$$\begin{array}{ccccccc} \cdots & \longrightarrow & H_n(A) & \xrightarrow{i_*} & H_n(B) & \xrightarrow{j_*} & H_n(C) \longrightarrow \\ & & & & \partial_* & & \\ & \longleftarrow & H_{n-1}(A) & \xrightarrow{i_*} & H_{n-1}(B) & \xrightarrow{j_*} & H_{n-1}(C) \longrightarrow \cdots \end{array}$$

where i_* and j_* are induced by $i.$ and $j.$, and ∂_* is a map we will define in the proof.

Having an exact sequence is good, since if we know most of the terms in an exact sequence, we can figure out the remaining ones. To do this, we also need to understand the maps i_* , j_* and ∂_* , but we don't need to understand *all*, since we can deduce some of them with exactness. Yet, we still need to know some of them, and since they are defined in the proof, you need to remember the proof.

Note, however, if we replace \mathbb{Z} in the definition of chain groups by a field (e.g. \mathbb{Q}), then all the groups become vector spaces. Then everything boils down to the rank-nullity theorem. Of course, this does not get us the right answer in exams, since we want to have homology groups over \mathbb{Z} , and not \mathbb{Q} , but this helps us to understand the exact sequences somewhat. If, at any point, homology groups confuses you, then you can try to work with homology groups over \mathbb{Q} and get a feel for what homology groups are like, since this is easier.

We are first going to apply this result to obtain the Mayer-Vietoris theorem.

Theorem (Mayer-Vietoris theorem). Let K, L, M, N be simplicial complexes with $K = M \cup N$ and $L = M \cap N$. We have the following inclusions and maps:

$$\begin{array}{ccc} L & \xhookrightarrow{i} & M \\ \downarrow j & & \downarrow k \\ N & \xhookrightarrow{\ell} & K. \end{array}$$

Then there exists some natural homomorphism $\partial_* : H_n(K) \rightarrow H_{n-1}(L)$ that gives the following long exact sequence:

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\partial_*} & H_n(L) & \xrightarrow{i_*+j_*} & H_n(M) \oplus H_n(N) & \xrightarrow{k_*-\ell_*} & H_n(K) \longrightarrow \\ & & & & \partial_* & & \\ & \longleftarrow & H_{n-1}(L) & \xrightarrow{i_*+j_*} & H_{n-1}(M) \oplus H_{n-1}(N) & \xrightarrow{k_*-\ell_*} & H_{n-1}(K) \longrightarrow \cdots \\ & & & & & & \\ \cdots & \longrightarrow & H_0(M) \oplus H_0(N) & \xrightarrow{k_*-\ell_*} & H_0(K) & \longrightarrow & 0 \end{array}$$

Note that unlike the Seifert-van Kampen theorem, this does not require the intersection $L = M \cap N$ to be (path) connected. This restriction was needed for Seifert-van Kampen since the fundamental group is unable to see things outside the path component of the basepoint, and hence it does not like non-path connected spaces well. However, homology groups don't have these problems.

$$0 \longrightarrow C_n(L) \xrightarrow{i_n+j_n} C_n(M) \oplus C_n(N) \xrightarrow{k_n-\ell_n} C_n(K) \longrightarrow 0$$

It is easy to see this is a short exact exact sequence of chain complexes. The image of $i_n + j_n$ is the set of all elements of the form (x, x) , and the kernel of $k_n - \ell_n$ is also these. It is also easy to see that $i_n + j_n$ is injective and $k_n - \ell_n$ is surjective. \square

Yet, to properly apply the Mayer-Vietoris sequence, we need to understand the map ∂_* . To do so, we need to prove the snake lemma.

$$0 \longrightarrow A. \xrightarrow{i.} B. \xrightarrow{j.} C. \longrightarrow 0$$
$$\begin{array}{ccccccc} \cdots & \longrightarrow & H_n(A) & \xrightarrow{i_*} & H_n(B) & \xrightarrow{j_*} & H_n(C) & \longrightarrow & \cdots \\ & & & & \partial_* & & & & \\ & \longleftarrow & H_{n-1}(A) & \xrightarrow{i_*} & H_{n-1}(B) & \xrightarrow{j_*} & H_{n-1}(C) & \longrightarrow & \cdots \end{array}$$

The method of proving this is sometimes known as “diagram chasing”, where we just “chase” around commutative diagrams to find the elements we need. The idea of the proof is as follows — in the short exact sequence, we can think of A as a subgroup of B , and C as the quotient B/A , by the first isomorphism theorem. So any element of C can be represented by an element of B . We apply the boundary map to this representative, and then exactness shows that this must come from some element of A . We then check carefully that these are well-defined, i.e. does not depend on the representatives chosen.

Proof. The proof of this is in general not hard. It just involves a lot of checking of the details, such as making sure the homomorphisms are well-defined, are actually homomorphisms, are exact at all the places etc. The only important and non-trivial part is just the construction of the map ∂_* .

First we look at the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & A_n & \xrightarrow{i_n} & B_n & \xrightarrow{j_n} & C_n \longrightarrow 0 \\ & & \downarrow d_n & & \downarrow d_n & & \downarrow d_n \\ 0 & \longrightarrow & A_{n-1} & \xrightarrow{i_{n-1}} & B_{n-1} & \xrightarrow{j_{n-1}} & C_{n-1} \longrightarrow 0 \end{array}$$

To construct $\partial_* : H_n(C) \rightarrow H_{n-1}(A)$, let $[x] \in H_n(C)$ be a class represented by $x \in Z_n(C)$. We need to find a cycle $z \in A_{n-1}$. By exactness, we know the map $j_n : B_n \rightarrow C_n$ is surjective. So there is a $y \in B_n$ such that $j_n(y) = x$. Since our target is A_{n-1} , we want to move down to the next level. So consider $d_n(y) \in B_{n-1}$. We would be done if $d_n(y)$ is in the image of i_{n-1} . By exactness, this is equivalent saying $d_n(y)$ is in the kernel of j_{n-1} . Since the diagram is commutative, we know

$$j_{n-1} \circ d_n(y) = d_n \circ j_n(y) = d_n(x) = 0,$$

using the fact that x is a cycle. So $d_n(y) \in \ker j_{n-1} = \text{im } i_{n-1}$. Moreover, by exactness again, i_{n-1} is injective. So there is a unique $z \in A_{n-1}$ such that $i_{n-1}(z) = d_n(y)$. We have now produced our z .

We are not done. We have $\partial_*[x] = [z]$ as our candidate definition, but we need to check many things:

- (i) We need to make sure ∂_* is indeed a homomorphism.
- (ii) We need $d_{n-1}(z) = 0$ so that $[z] \in H_{n-1}(A)$;
- (iii) We need to check $[z]$ is well-defined, i.e. it does not depend on our choice of y and x for the homology class $[x]$.
- (iv) We need to check the exactness of the resulting sequence.

We now check them one by one:

- (i) Since everything involved in defining ∂_* are homomorphisms, it follows that ∂_* is also a homomorphism.
- (ii) We check $d_{n-1}(z) = 0$. To do so, we need to add an additional layer.

$$\begin{array}{ccccccc} 0 & \longrightarrow & A_n & \xrightarrow{i_n} & B_n & \xrightarrow{j_n} & C_n \longrightarrow 0 \\ & & \downarrow d_n & & \downarrow d_n & & \downarrow d_n \\ 0 & \longrightarrow & A_{n-1} & \xrightarrow{i_{n-1}} & B_{n-1} & \xrightarrow{j_{n-1}} & C_{n-1} \longrightarrow 0 \\ & & \downarrow d_{n-1} & & \downarrow d_{n-1} & & \downarrow d_{n-1} \\ 0 & \longrightarrow & A_{n-2} & \xrightarrow{i_{n-2}} & B_{n-2} & \xrightarrow{j_{n-2}} & C_{n-2} \longrightarrow 0 \end{array}$$

We want to check that $d_{n-1}(z) = 0$. We will use the commutativity of the diagram. In particular, we know

$$i_{n-2} \circ d_{n-1}(z) = d_{n-1} \circ i_{n-1}(z) = d_{n-1} \circ d_n(y) = 0.$$

By exactness at A_{n-2} , we know i_{n-2} is injective. So we must have $d_{n-1}(z) = 0$.

- (iii) (a) First, in the proof, suppose we picked a different y' such that $j_n(y') = j_n(y) = x$. Then $j_n(y' - y) = 0$. So $y' - y \in \ker j_n = \text{im } i_n$. Let $a \in A_n$ be such that $i_n(a) = y' - y$. Then

$$\begin{aligned} d_n(y') &= d_n(y' - y) + d_n(y) \\ &= d_n \circ i_n(a) + d_n(y) \\ &= i_{n-1} \circ d_n(a) + d_n(y). \end{aligned}$$

Hence when we pull back $d_n(y')$ and $d_n(y)$ to A_{n-1} , the results differ by the boundary $d_n(a)$, and hence produce the same homology class.

- (b) Suppose $[x'] = [x]$. We want to show that $\partial_*[x] = \partial_*[x']$. This time, we add a layer above.

$$\begin{array}{ccccccc} 0 & \longrightarrow & A_{n+1} & \xrightarrow{i_{n+1}} & B_{n+1} & \xrightarrow{j_{n+1}} & C_{n+1} \longrightarrow 0 \\ & & \downarrow d_{n+1} & & \downarrow d_{n+1} & & \downarrow d_{n+1} \\ 0 & \longrightarrow & A_n & \xrightarrow{i_n} & B_n & \xrightarrow{j_n} & C_n \longrightarrow 0 \\ & & \downarrow d_n & & \downarrow d_n & & \downarrow d_n \\ 0 & \longrightarrow & A_{n-1} & \xrightarrow{i_{n-1}} & B_{n-1} & \xrightarrow{j_{n-1}} & C_{n-1} \longrightarrow 0 \end{array}$$

By definition, since $[x'] = [x]$, there is some $c \in C_{n+1}$ such that

$$x' = x + d_{n+1}(c).$$

By surjectivity of j_{n+1} , we can write $c = j_{n+1}(b)$ for some $b \in B_{n+1}$. By commutativity of the squares, we know

$$x' = x + j_n \circ d_{n+1}(b).$$

The next step of the proof is to find some y such that $j_n(y) = x$. Then

$$j_n(y + d_{n+1}(b)) = x'.$$

So the corresponding y' is $y' = y + d_{n+1}(b)$. So $d_n(y) = d_n(y')$, and hence $\partial_*[x] = \partial_*[x']$.

- (iv) This is yet another standard diagram chasing argument. When reading this, it is helpful to look at a diagram and see how the elements are chased along. It is even more beneficial to attempt to prove this yourself.

- (a) $\text{im } i_* \subseteq \ker j_*$: This follows from the assumption that $i_n \circ j_n = 0$.
 (b) $\ker j_* \subseteq \text{im } i_*$: Let $[b] \in H_n(B)$. Suppose $j_*([b]) = 0$. Then there is some $c \in C_{n+1}$ such that $j_n(b) = d_{n+1}(c)$. By surjectivity of j_{n+1} , there is some $b' \in B_{n+1}$ such that $j_{n+1}(b') = c$. By commutativity, we know $j_n(b) = j_n \circ d_{n+1}(b')$, i.e.

$$j_n(b - d_{n+1}(b')) = 0.$$

By exactness of the sequence, we know there is some $a \in A_n$ such that

$$i_n(a) = b - d_{n+1}(b').$$

Moreover,

$$i_{n-1} \circ d_n(a) = d_n \circ i_n(a) = d_n(b - d_{n+1}(b')) = 0,$$

using the fact that b is a cycle. Since i_{n-1} is injective, it follows that $d_n(a) = 0$. So $[a] \in H_n(A)$. Then

$$i_*([a]) = [b] - [d_{n+1}(b')] = [b].$$

So $[b] \in \text{im } i_*$.

- (c) $\text{im } j_* \subseteq \ker \partial_*$: Let $[b] \in H_n(B)$. To compute $\partial_*(j_*([b]))$, we first pull back $j_n(b)$ to $b \in B_n$. Then we compute $d_n(b)$ and then pull it back to A_{n+1} . However, we know $d_n(b) = 0$ since b is a cycle. So $\partial_*(j_*([b])) = 0$, i.e. $\partial_* \circ j_* = 0$.
- (d) $\ker \partial_* \subseteq \text{im } j_*$: Let $[c] \in H_n(C)$ and suppose $\partial_*([c]) = 0$. Let $b \in B_n$ be such that $j_n(b) = c$, and $a \in A_{n-1}$ such that $i_{n-1}(a) = d_n(b)$. By assumption, $\partial_*([c]) = [a] = 0$. So we know a is a boundary, say $a = d_n(a')$ for some $a' \in A_n$. Then by commutativity we know $d_n(b) = d_n \circ i_n(a')$. In other words,

$$d_n(b - i_n(a')) = 0.$$

So $[b - i_n(a')] \in H_n(B)$. Moreover,

$$j_*([b - i_n(a')]) = [j_n(b) - j_n \circ i_n(a')] = [c].$$

So $[c] \in \text{im } j_*$.

- (e) $\text{im } \partial_* \subseteq \ker i_*$: Let $[c] \in H_n(C)$. Let $b \in B_n$ be such that $j_n(b) = c$, and $a \in A_{n-1}$ be such that $i_n(a) = d_n(b)$. Then $\partial_*([c]) = [a]$. Then

$$i_*([a]) = [i_n(a)] = [d_n(b)] = 0.$$

So $i_* \circ \partial_* = 0$.

- (f) $\ker i_* \subseteq \text{im } \partial_*$: Let $[a] \in H_n(A)$ and suppose $i_*([a]) = 0$. So we can find some $b \in B_{n+1}$ such that $i_n(a) = d_{n+1}(b)$. Let $c = j_{n+1}(b)$. Then

$$d_{n+1}(c) = d_{n+1} \circ j_{n+1}(b) = j_n \circ d_{n+1}(b) = j_n \circ i_n(a) = 0.$$

So $[c] \in H_n(C)$. Then $[a] = \partial_*([c])$ by definition of ∂_* . So $[a] \in \text{im } \partial_*$.


□

7.5 Continuous maps and homotopy invariance

This is the most technical part of the homology section of the course. The goal is to see that the homology groups $H_*(K)$ depends only on the polyhedron, and not the simplicial structure on it. Moreover, we will show that they are *homotopy invariants* of the space, and that homotopic maps $f \simeq g : |K| \rightarrow |L|$ induce equal maps $H_*(K) \rightarrow H_*(L)$. Note that this is a lot to prove. At this point, we don't even know arbitrary continuous maps induce *any* map on the homology. We only know simplicial maps do.

We start with a funny definition.

Definition (Contiguous maps). Simplicial maps $f, g : K \rightarrow L$ are *contiguous* if for each $\sigma \in K$, the simplices $f(\sigma)$ and $g(\sigma)$ (i.e. the simplices spanned by the image of the vertices of σ) are faces of some simplex $\tau \in L$.



The significance of this definition comes in two parts: simplicial approximations of the same map are contiguous, and contiguous maps induce the same maps on homology.

Lemma. If $f, g : K \rightarrow L$ are simplicial approximations to the same map F , then f and g are contiguous.

Proof. Let $\sigma \in K$, and pick some $s \in \sigma$. Then $F(s) \in \tau$ for some $\tau \in L$. Then the definition of simplicial approximation implies that for any simplicial approximation f to F , $f(\sigma)$ spans a face of τ . \square

Lemma. If $f, g : K \rightarrow L$ are contiguous simplicial maps, then

$$f_* = g_* : H_n(K) \rightarrow H_n(L)$$

for all n .

Geometrically, the homotopy is obvious. If $f(\sigma)$ and $g(\sigma)$ are both faces of τ , then we just pick the homotopy as τ . The algebraic definition is less inspiring, and it takes some staring to see it is indeed what we think it should be.


Proof. We will write down a chain homotopy between f and g :

$$h_n((a_0, \dots, a_n)) = \sum_{i=0}^n (-1)^i [f(a_0), \dots, f(a_i), g(a_i), \dots, g(a_n)],$$

where the square brackets means the corresponding oriented simplex if there are no repeats, 0 otherwise.

We can now check by direct computation that this is indeed a chain homotopy. \square

Now we know that if $f, g : K \rightarrow L$ are both simplicial approximations to the same F , then they induce equal maps on homology. However, we not yet know that continuous maps induce well-defined maps on homologies, since to produce simplicial approximations, we needed to perform barycentric subdivision, and we need to make sure this does not change the homology.



Given a barycentric subdivision, we need to choose a simplicial approximation to the identity map $a : K' \rightarrow K$. It turns out this is easy to do, and we can do it almost arbitrarily.

Lemma. Each vertex $\hat{\sigma} \in K'$ is a barycenter of some $\sigma \in K$. Then we choose $a(\hat{\sigma})$ to be an arbitrary vertex of σ . This defines a function $a : V_{K'} \rightarrow V_K$. This a is a simplicial approximation to the identity. Moreover, *every* simplicial approximation to the identity is of this form.

Proof. Omitted. \square

Next, we need to show this gives an isomorphism on the homologies.

Proposition. Let K' be the barycentric subdivision of K , and $a : K' \rightarrow K$ a simplicial approximation to the identity map. Then the induced map $a_* : H_n(K') \rightarrow H_n(K)$ is an isomorphism for all n .

Proof. We first deal with K being a simplex σ and its faces. Now that K is just a cone (with any vertex as the cone vertex), and K' is also a cone (with the barycenter as the cone vertex). Therefore

$$H_n(K) \cong H_n(K') \cong \begin{cases} \mathbb{Z} & n = 0 \\ 0 & n > 0 \end{cases}$$

So only $n = 0$ is (a little) interesting, but it is easy to check that a_* is an isomorphism in this case, since it just maps a vertex to a vertex, and all vertices in each simplex are in the same homology class..

To finish the proof, note that K is built up by gluing up simplices, and K' is built by gluing up subdivided simplices. So to understand their homology groups, we use the Mayer-Vietoris sequence.

Given a complicated simplicial complex K , let σ be a maximal dimensional simplex of K . We let $L = K \setminus \{\sigma\}$ (note that L includes the boundary of σ). We let $S = \{\sigma \text{ and all its faces}\} \subseteq K$ and $T = L \cap S$.

We can do similarly for K' , and let L', S', T' be the corresponding barycentric subdivisions. We have $K = L \cup S$ and $K' = L' \cup S'$ (and $L' \cap S' = T'$). By the previous lemma, we see our construction of a gives $a(L') \subseteq L$, $a(S') \subseteq S$ and $a(T') \subseteq T$. So these induce maps of the corresponding homology groups

$$\begin{array}{ccccccccc} H_n(T') & \rightarrow & H_n(S') \oplus H_n(L') & \rightarrow & H_n(K') & \rightarrow & H_{n-1}(T') & \rightarrow & H_{n-1}(S') \oplus H_{n-1}(L') \\ \downarrow a_* & & \downarrow a_* \oplus a_* & & \downarrow a_* & & \downarrow a_* & & \downarrow a_* \oplus a_* \\ H_n(T) & \rightarrow & H_n(S) \oplus H_n(L) & \rightarrow & H_n(K) & \rightarrow & H_{n-1}(T) & \rightarrow & H_{n-1}(S) \oplus H_{n-1}(L) \end{array}$$

By induction, we can assume all but the middle maps are isomorphisms. By the five lemma, this implies the middle map is an isomorphism, where the five lemmas is as follows: \square

Lemma (Five lemma). Consider the following commutative diagram:

$$\begin{array}{ccccccccc} A_1 & \longrightarrow & B_1 & \longrightarrow & C_1 & \longrightarrow & D_1 & \longrightarrow & E_1 \\ \downarrow a & & \downarrow b & & \downarrow c & & \downarrow d & & \downarrow e \\ A_2 & \longrightarrow & B_2 & \longrightarrow & C_2 & \longrightarrow & D_2 & \longrightarrow & E_2 \end{array}$$

If the top and bottom rows are exact, and a, b, d, e are isomorphisms, then c is also an isomorphism.

Proof. Exercise in example sheet. \square

We now have everything we need to move from simplicial maps to continuous maps. Putting everything we have proved so far together, we obtain the following result:

Proposition. To each continuous map $f : |K| \rightarrow |L|$, there is an associated map $f_* : H_n(K) \rightarrow H_n(L)$ (for all n) given by

$$f_* = s_* \circ \nu_{K,r}^{-1},$$

where $s : K^{(r)} \rightarrow L$ is a simplicial approximation to f , and $\nu_{K,r} : H_n(K^{(r)}) \rightarrow H_n(K)$ is the isomorphism given by composing maps $H_n(K^{(i)}) \rightarrow H_n(K^{(i-1)})$ induced by simplicial approximations to the identity.

Furthermore:

- (i) f_* does not depend on the choice of r or s .
- (ii) If $g : |M| \rightarrow |K|$ is another continuous map, then

$$(f \circ g)_* = f_* \circ g_*.$$

Corollary. If $f : |K| \rightarrow |L|$ is a homeomorphism, then $F_* : H_n(K) \rightarrow H_n(L)$ is an isomorphism for all n .

Proof. Immediate from (ii) of previous proposition. \square

This is good. We know now that homology groups is a property of the space itself, not simplicial complexes. For example, we have computed the homology groups of a particular triangulation of the n -sphere, and we know it is true for *all* triangulations of the n -sphere.

We're not exactly there yet. We have just seen that homology groups are invariant under homeomorphisms. What we would like to know is that they are invariant under *homotopy*. In other words, we want to know homotopy equivalent maps induce equal maps on the homology groups.

The strategy is:

- (i) Show that “small” homotopies don't change the maps on H_n
- (ii) Note that all homotopies can be decomposed into “small” homotopies.

Lemma. Let L be a simplicial complex (with $|L| \subseteq \mathbb{R}^n$). Then there is an $\varepsilon = \varepsilon(L) > 0$ such that if $f, g : |K| \rightarrow |L|$ satisfying $\|f(x) - g(x)\| < \varepsilon$, then $f_* = g_* : H_n(K) \rightarrow H_n(L)$ for all n .

The idea of the proof is that if $|f(x) - g(x)|$ is small enough, we can barycentrically subdivide L such that we get a simplicial approximation to both f and g .

Proof. By the Lebesgue number lemma, there is an $\varepsilon > 0$ such that each ball of radius 2ε in $|L|$ lies in some star $\text{St}_L(w)$.

Now apply the Lebesgue number lemma again to $\{f^{-1}(B_\varepsilon(y))\}_{y \in |L|}$, an open cover of $|K|$, and get $\delta > 0$ such that for all $x \in |K|$, we have

$$f(B_\delta(x)) \subseteq B_\varepsilon(y) \subseteq B_{2\varepsilon}(y) \subseteq \text{St}_L(w)$$

for some $y \in |L|$ and $\text{St}_L(w)$. Now since g and f differ by at most ε , we know

$$g(B_\delta(x)) \subseteq B_{2\varepsilon}(y) \subseteq \text{St}_L(w).$$

Now subdivide r times so that $\mu(K^{(r)}) < \frac{1}{2}\delta$. So for all $v \in V_K(R)$, we know

$$\text{St}_{K^{(r)}}(v) \subseteq B_\delta(V).$$

This gets mapped by *both* f and g to $\text{St}_L(w)$ for the same $w \in V_L$. We define $s : V_{K^{(r)}} \rightarrow V_L$ sending $v \mapsto w$. \square

Theorem. Let $f \simeq g : |K| \rightarrow |L|$. Then $f_* = g_*$.

Proof. Let $H : |K| \times I \rightarrow |L|$. Since $|K| \times I$ is compact, we know H is uniformly continuous. Pick $\varepsilon = \varepsilon(L)$ as in the previous lemma. Then there is some δ such that $|s - t| < \delta$ implies $|H(x, s) - H(x, t)| < \varepsilon$ for all $x \in |K|$.

Now choose $0 = t_0 < t_1 < \dots < t_n = 1$ such that $t_i - t_{i-1} < \delta$ for all i . Define $f_i : |K| \rightarrow |L|$ by $f_i(x) = H(x, t_i)$. Then we know $\|f_i - f_{i-1}\| < \varepsilon$ for all i . Hence $(f_i)_* = (f_{i-1})_*$. Therefore $(f_0)_* = (f_n)_*$, i.e. $f_* = g_*$. \square

This is good, since we know we can not only deal with spaces that are homeomorphic to complexes, but spaces that are homotopic to complexes. This is important, since all complexes are compact, but we would like to talk about non-compact spaces such as open balls as well.

Definition (*h-triangulation and homology groups*). An h -triangulation of a space X is a simplicial complex K and a homotopy equivalence $g : |K| \rightarrow X$. We define $H_n(X) = H_n(K)$ for all n .

Lemma. $H_n(X)$ is well-defined, i.e. it does not depend on the choice of K .

Proof. Clear from previous theorem. \square


We have spent a lot of time and effort developing all the technical results and machinery of homology groups. We will now use them to *do stuff*.

Digression — historical motivation


At first, we motivated the study of algebraic topology by saying we wanted to show that \mathbb{R}^n and \mathbb{R}^m are not homeomorphic. However, historically, this is not why people studied algebraic topology, since there are other ways to prove this is true.

Initially, people were interested in studying knots. These can be thought of as embeddings $S \hookrightarrow S^3$. We really should just think of S^3 as $\mathbb{R}^3 \cup \{\infty\}$, where the point at infinity is just there for convenience.

The most interesting knot we know is the *unknot* U :



A less interesting but still interesting knot is the *trefoil* T .



One question people at the time asked was whether the trefoil knot is just the unknot in disguise. It obviously isn't, but can we prove it? In general, given two knots, is there any way we can distinguish if they are the same?

The idea is to study the fundamental groups of the knots. It would obviously be silly to compute the fundamental groups of U and T themselves, since they are both isomorphic to S^1 and the groups are just \mathbb{Z} . Instead, we look at the fundamental groups of the complements. It is not difficult to see that

$$\pi_1(S^3 \setminus U) \cong \mathbb{Z},$$

and with some suitable machinery, we find

$$\pi_1(S^3 \setminus T) \cong \langle a, b \mid a^3 b^{-2} \rangle.$$

Staring at it hard enough, we can construct a surjection $\pi_1(S^3 \setminus T) \rightarrow S_3$. This tells us $\pi_1(S^3 \setminus T)$ is non-abelian, and is certainly not \mathbb{Z} . So we know U and T are genuinely different knots.

7.6 Homology of spheres and applications

Lemma. The sphere S^{n-1} is triangulable, and

$$H_k(S^{n-1}) \cong \begin{cases} \mathbb{Z} & k = 0, n-1 \\ 0 & \text{otherwise} \end{cases}$$

Proof. We already did this computation for the standard $(n-1)$ -sphere $\partial\Delta^n$, where Δ^n is the standard n -simplex. \square

We immediately have a few applications


Proposition. $\mathbb{R}^n \not\cong \mathbb{R}^m$ for $m \neq n$.

Proof. See example sheet 4. \square

Theorem (Brouwer's fixed point theorem (in all dimensions)). There is no retraction D^n onto $\partial D^n \cong S^{n-1}$. So every continuous map $f : D^n \rightarrow D^n$ has a fixed point.

Proof. The proof is exactly the same as the two-dimensional case, with homology groups instead of the fundamental group.

We first show the second part from the first. Suppose $f : D^n \rightarrow D^n$ has no fixed point. Then the following $g : D^n \rightarrow \partial D^n$ is a continuous retraction.



So we now show no such continuous retraction can exist. Suppose $r : D^n \rightarrow \partial D^n$ is a retraction, i.e. $r \circ i \simeq \text{id} : \partial D^n \rightarrow D^n$.

$$S^{n-1} \xrightarrow{i} D^n \xrightarrow{r} S^{n-1}$$

We now take the $(n-1)$ th homology groups to obtain

$$H_{n-1}(S^{n-1}) \xrightarrow{i_*} H_{n-1}(D^n) \xrightarrow{r_*} H_{n-1}(S^{n-1}).$$

Since $r \circ i$ is homotopic to the identity, this composition must also be the identity, but this is clearly nonsense, since $H_{n-1}(S^{n-1}) \cong \mathbb{Z}$ while $H_{n-1}(D^n) \cong 0$. So such a continuous retraction cannot exist. \square

Note it is important that we can work with continuous maps directly, and not just their simplicial approximations. Otherwise, here we can only show that every simplicial approximation of f has a fixed point, but this does not automatically entail that f itself does not have a fixed point.


For the next application, recall from the first example sheet that if n is odd, then the antipodal map $a : S^n \rightarrow S^n$ is homotopic to the identity. What if n is even?

The idea is to consider the effect on the homology group:

$$a_* : H_n(S^n) \rightarrow H_n(S^n).$$

By our previous calculation, we know a_* is a map $a_* : \mathbb{Z} \rightarrow \mathbb{Z}$. If a is homotopic to the identity, then a_* should be homotopic to the identity map. We will now compute a_* and show that it is multiplication by -1 when n is even.

To do this, we want to use a triangulation that is compatible with the antipodal map. The standard triangulation clearly doesn't work. Instead, we use the following triangulation $h : |K| \rightarrow S^n$:



The vertices of K are given by

$$V_k = \{\pm \mathbf{e}_0, \pm \mathbf{e}_1, \dots, \pm \mathbf{e}_n\}.$$

This triangulation works nicely with the antipodal map, since this maps a vertex to a vertex. To understand the fundamental group better, we need the following lemma:

Lemma. In the triangulation of S^n given by vertices $V_k = \{\pm \mathbf{e}_0, \pm \mathbf{e}_1, \dots, \pm \mathbf{e}_n\}$, the element

$$x = \sum_{\epsilon \in \{\pm 1\}^{n+1}} \epsilon_0 \cdots \epsilon_n (\epsilon_0 \mathbf{e}_0, \dots, \epsilon_n \mathbf{e}_n)$$

is a cycle and generates $H_n(S^n)$.

Proof. By direct computation, we see that $dx = 0$. So x is a cycle. To show it generates $H_n(S^n)$, we note that everything in $H_n(S^n) \cong \mathbb{Z}$ is a multiple of the generator, and since x has coefficients ± 1 , it cannot be a multiple of anything else (apart from $-x$). So x is indeed a generator. \square

Now we can prove our original proposition.


Proposition. If n is even, then the antipodal map $a \neq \text{id}$.

Proof. We can directly compute that $a_*x = (-1)^{n+1}x$. If n is even, then $a_* = -1$, but $\text{id}_* = 1$. So $a \neq \text{id}$. \square


7.7 Homology of surfaces

We want to study compact surfaces and their homology groups. To work with the simplicial homology, we need to assume they are triangulable. We will not prove this fact, and just assume it to be true (it really is).

Recall we have classified compact surfaces, and have found the following orientable surfaces Σ_g .



We also have non-compact versions of these, known as F_g , where we take the above ones and cut out a hole:




We want to compute the homology groups of these Σ_g . One way to do so is to come up with specific triangulations of the space, and then compute the homology groups directly. However, there is a better way to do it. Given a Σ_g , we can slice it apart along a circle and write it as


$$\Sigma_g = F_{g-1} \cup F_1.$$




To do so, we need to compute $H_*(F_g)$. Fortunately, this is not too hard, since it turns out F_g is homotopic to a relatively simple space. recall we produced Σ_g by starting with a $4g$ -gon and gluing edges:




Now what is F_g ? F_g is Σ_g with a hole cut out of it. We'll cut the hole out at the center, and be left with



We can now expand the hole to the boundary, and get a deformation retraction from F_g to its boundary:




$$H_n(F_g) \cong H_n(X_{2g}) \cong \begin{cases} \mathbb{Z} & n = 0 \\ \mathbb{Z}^{2g} & n = 1 \\ 0 & n > 1 \end{cases}$$
$$\begin{array}{ccccccc}
0 & \longrightarrow & H_2(S^1) & \longrightarrow & H_2(F_{g-1}) \oplus H_2(F_1) & \longrightarrow & H_2(\Sigma_g) \\
& & & & & & \searrow \\
& & & & & & \rightarrow H_1(S^1) \longrightarrow H_1(F_{g-1}) \oplus H_1(F_1) \longrightarrow H_1(\Sigma_g) \\
& & & & & & \searrow \\
& & & & & & \rightarrow H_0(S^1) \longrightarrow H_0(F_{g-1}) \oplus H_0(F_1) \longrightarrow H_0(\Sigma_g) \longrightarrow 0
\end{array}$$
$$0 \rightarrow H_2(\Sigma_g) \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}^{2g} \rightarrow H_1(\Sigma_g) \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}^2 \rightarrow \mathbb{Z} \rightarrow 0$$
$$0 \longrightarrow \mathbb{Z}^{2g} \longrightarrow H_1(\Sigma_g) \longrightarrow \ker(\mathbb{Z} \rightarrow \mathbb{Z}^2) \longrightarrow 0.$$
$$H_1(\Sigma_q) \cong \mathbb{Z}^{2g}.$$

91

7.8 Rational homology, Euler and Lefschetz numbers

So far, we have been working with chains with integral coefficients. It turns out we can use rational coefficients instead. In the past, $C_n(K)$ was an abelian group, or a \mathbb{Z} -module. If we use rational coefficients, since \mathbb{Q} is a field, this becomes a vector space, and we can use a lot of nice theorems about vector spaces, such as the rank-nullity theorem. Moreover, we can reason talk about things like the dimensions of these homology groups.

Definition (Rational homology group). For a simplicial complex K , we can define the *rational n -chain* $C_n(K, \mathbb{Q})$ in the same way as $C_n(K) = C_n(K, \mathbb{Z})$. That is, $C_n(K, \mathbb{Q})$ is the vector space over \mathbb{Q} with basis the n -simplices of K (with a choice of orientation).

We can define d_n, Z_n, B_n as before, and the *rational n th homology group* is

$$H_n(K; \mathbb{Q}) \cong \frac{Z_n(K; \mathbb{Q})}{B_n(K; \mathbb{Q})}.$$

Now our homology group is a vector space, and it is much easier to work with. However, the consequence is that we will lose some information. Fortunately, the way in which we lose information is very well-understood and rather simple.

Lemma. If $H_n(K) \cong \mathbb{Z}^k \oplus F$ for F a finite group, then $H_n(K; \mathbb{Q}) \cong \mathbb{Q}^k$.

Proof. Exercise. □

Hence, when passing to rational homology groups, we lose all information about the torsion part. In some cases this information loss is not very significant, but in certain cases, it can be. For example, in \mathbb{RP}^2 , we have

$$H_n(\mathbb{RP}^2) \cong \begin{cases} \mathbb{Z} & n = 0 \\ \mathbb{Z}/2 & n = 1 \\ 0 & n > 1 \end{cases}$$

If we pass on to rational coefficients, then we have lost everything in $H_1(\mathbb{RP}^2)$, and \mathbb{RP}^2 looks just like a point:

$$H_n(\mathbb{RP}^2; \mathbb{Q}) \cong H_n(*; \mathbb{Q}).$$

Example. We have

$$H_n(S^n, \mathbb{Q}) \cong \begin{cases} \mathbb{Q} & k = 0, n \\ 0 & \text{otherwise} \end{cases}.$$

We also have

$$H_n(\Sigma_g, \mathbb{Q}) \cong \begin{cases} \mathbb{Q} & k = 0, 2 \\ \mathbb{Q}^{2g} & k = 1 \\ 0 & \text{otherwise} \end{cases}.$$

In this case, we have not lost any information because there was no torsion part of the homology groups.

However, for the non-orientable surfaces, we have

$$H_k(E_n, \mathbb{Q}) = \begin{cases} \mathbb{Q} & k = 0 \\ \mathbb{Q}^{n-1} & k = 1 \\ 0 & \text{otherwise} \end{cases},$$

This time, this is different from the integral coefficient case, where we have an extra \mathbb{Z}_2 term in H_1 .

The advantage of this is that the homology “groups” are in fact vector spaces, and we can talk about things like dimensions. Also, maps on homology groups are simply linear maps, i.e. matrices, and we can study their properties with proper linear algebra.

Recall that the Euler characteristic of a polyhedron is defined as “faces – edges + vertices”. This works for two-dimensional surfaces, and we would like to extend this to higher dimensions. There is an obvious way to do so, by counting higher-dimensional surfaces and putting the right signs. However, if we define it this way, it is not clear that this is a property of the space itself, and not just a property of the triangulation. Hence, we define it as follows:

Definition (Euler characteristic). The *Euler characteristic* of a triangulable space X is

$$\chi(X) = \sum_{i \geq 0} (-1)^i \dim_{\mathbb{Q}} H_i(X; \mathbb{Q}).$$

This clearly depends only on the homotopy type of X , and not the triangulation. We will later show this is equivalent to what we used to have.

More generally, we can define the *Lefschetz number*.

Definition (Lefschetz number). Given any map $f : X \rightarrow X$, we define the *Lefschetz number* of f as

$$L(f) = \sum_{i \geq 0} (-1)^i \operatorname{tr}(f_* : H_i(X; \mathbb{Q}) \rightarrow H_i(X; \mathbb{Q})).$$

Why is this a generalization of the Euler characteristic. Just note that the trace of the identity map is the number of dimensions. So we have

$$\chi(X) = L(\operatorname{id}).$$

Example. We have

$$\chi(S^n) = \begin{cases} 2 & n \text{ even} \\ 0 & n \text{ odd} \end{cases}$$

We also have

$$\chi(\Sigma_g) = 2 - 2g, \quad \chi(E_n) = 2 - n.$$

Example. if $\alpha : S^n \rightarrow S^n$ is the antipodal map, we saw that $\alpha_* : H_n(S^n) \rightarrow H_n(S^n)$ is multiplication by $(-1)^{n+1}$. So

$$L(\alpha) = 1 + (-1)^n (-1)^{n+1} = 1 - 1 = 0.$$

We see that even though the antipodal map has different behaviour for different dimensions, the Lefschetz number ends up being zero all the time. We will soon see why this is the case.

Why do we want to care about the Lefschetz number? The important thing is that we will use this to prove a really powerful generalization of Brouwer's fixed point theorem that allows us to talk about things that are not balls.

Before that, we want to understand the Lefschetz number first. To define the Lefschetz number, we take the trace of f_* , and this is a map of the homology groups. However, we would like to understand the Lefschetz number in terms of the chain groups, since these are easier to comprehend. Recall that the homology groups are defined as quotients of the chain group, so we would like to know what happens to the trace when we take quotients.

Lemma. Let V be a finite-dimensional vector space and $W \leq V$ a subspace. Let $A : V \rightarrow V$ be a linear map such that $A(W) \subseteq W$. Let $B = A|_W : W \rightarrow W$ and $C : V/W \rightarrow V/W$ the induced map on the quotient. Then

$$\text{tr}(A) = \text{tr}(B) + \text{tr}(C).$$

Proof. In the right basis,

$$A = \begin{pmatrix} B & A' \\ 0 & C \end{pmatrix}.$$

□

What this allows us to do is to not look at the induced maps on homology, but just the maps on chain complexes. This makes our life much easier when it comes to computation.

Corollary. Let $f_* : C_*(K, \mathbb{Q}) \rightarrow C_*(K, \mathbb{Q})$ be a chain map. Then

$$\sum_{i \geq 0} (-1)^i \text{tr}(f_i : C_i(K) \rightarrow C_i(K)) = \sum_{i \geq 0} (-1)^i \text{tr}(f_* : H_i(K) \rightarrow H_i(K)),$$

with homology groups understood to be over \mathbb{Q} .

This is a great corollary. The thing on the right is the conceptually right thing to have — homology groups are nice and are properties of the space itself, not the triangulation. However, to actually do computations, we want to work with the chain groups and actually calculate with chain groups.

Proof. There is an exact sequence

$$0 \longrightarrow B_i(K; \mathbb{Q}) \longrightarrow Z_i(K; \mathbb{Q}) \longrightarrow H_i(K; \mathbb{Q}) \longrightarrow 0$$

This is since $H_i(K, \mathbb{Q})$ is defined as the quotient of Z_i over B_i . We also have the exact sequence

$$0 \longrightarrow Z_i(K; \mathbb{Q}) \longrightarrow C_i(K; \mathbb{Q}) \xrightarrow{d_i} B_{i-1}(K; \mathbb{Q}) \longrightarrow 0$$

This is true by definition of B_{i-1} and Z_i . Let $f_i^H, f_i^B, f_i^Z, f_i^C$ be the various maps induced by f on the corresponding groups. Then we have

$$\begin{aligned} L(|f|) &= \sum_{i \geq 0} (-1)^i \operatorname{tr}(f_i^H) \\ &= \sum_{i \geq 0} (-1)^i (\operatorname{tr}(f_i^Z) - \operatorname{tr}(f_i^B)) \\ &= \sum_{i \geq 0} (-1)^i (\operatorname{tr}(f_i^C) - \operatorname{tr}(f_{i-1}^B) - \operatorname{tr}(f_i^B)). \end{aligned}$$

Because of the alternating signs in dimension, each f_i^B appears twice in the sum with opposite signs. So all f_i^B cancel out, and we are left with

$$L(|f|) = \sum_{i \geq 0} (-1)^i \operatorname{tr}(f_i^C).$$

□

Since $\operatorname{tr}(\operatorname{id}|_{C_i(K; \mathbb{Q})}) = \dim_{\mathbb{Q}} C_i(K; \mathbb{Q})$, which is just the number of i simplices, this tells us the Euler characteristic we just defined is the usual Euler characteristic, ie,

$$\chi(X) = \sum_{i \geq 0} (-1)^i \text{number of } i \text{ simplices.}$$

Finally, we get to the important theorem of the section.

Theorem (Lefschetz fixed point theorem). Let $f : X \rightarrow X$ be a continuous map from a triangulable space to itself. If $L(f) \neq 0$, then f has a fixed point.

Proof. We prove the contrapositive. Suppose f has no fixed point. We will show that $L(f) = 0$. Let

$$\delta = \inf\{|x - f(x)| : x \in X\}$$

thinking X as a subset of \mathbb{R}^n . We know this is non-zero, since f has no fixed point, and X is compact (and hence the infimum point is achieved by some x).

Choose a triangulation $L : |K| \rightarrow X$ such that $\mu(K) < \frac{\delta}{2}$. We now let

$$g : K^{(r)} \rightarrow K$$

be a simplicial approximation to f . Since we picked our triangulation to be so fine, for $x \in \sigma \in K$, we have

$$|f(x) - g(x)| < \frac{\delta}{2}$$

since the mesh is already less than $\frac{\delta}{2}$. Also, we know

$$|f(x) - x| \geq \delta.$$

So we have

$$|g(x) - x| > \frac{\delta}{2}.$$

So we must have $g(x) \notin \sigma$. The conclusion is that for any $\sigma \in K$, we must have

$$g(\sigma) \cap \sigma = \emptyset.$$

Now we compute $L(f) = L(|g|)$. The only complication here is that g is a map from $K^{(r)}$ to K , and the domains and codomains are different. So we need to compose it with $s_i : C_i(K; \mathbb{Q}) \rightarrow C_i(K^{(r)}; \mathbb{Q})$ induced by inverses of simplicial approximations to the identity map. Then we have

$$\begin{aligned} L(|g|) &= \sum_{i \geq 0} (-1)^i \operatorname{tr}(g_* : H_i(X; \mathbb{Q}) \rightarrow H_i(X; \mathbb{Q})) \\ &= \sum_{i \geq 0} (-1)^i \operatorname{tr}(g_i \circ s_i : C_i(K; \mathbb{Q}) \rightarrow C_i(K; \mathbb{Q})) \end{aligned}$$

Now note that s_i takes simplices of σ to sums of subsimplices of σ . So $g_i \circ s_i$ takes every simplex off itself. So each diagonal terms of the matrix of $g_i \circ s_i$ is 0! Hence the trace is

$$L(|g|) = 0$$

□

Example. If X is *any* contractible polyhedron (e.g. a disk), then $L(f) = 1$ for any $f : X \rightarrow X$, which is obvious once we contract the space to a point. So f has a fixed point.

Example. Suppose G is a path-connected topological group, i.e. X is a group and a topological space, and inverse and multiplication are continuous maps.

If $g \neq 1$, then the map

$$\begin{aligned} r_g : G &\rightarrow G \\ \gamma &\mapsto \gamma g \end{aligned}$$

has no fixed point. This implies $L(r_g) = 0$. However, since the space is path connected, $r_g \simeq \operatorname{id}$, where the homotopy is obtained by multiplying elements along the path from e to r_g . So

$$\chi(G) = L(\operatorname{id}_G) = L(r_g) = 0.$$

So if $G \neq 1$, then in fact $\chi(G) = 0$.

This is quite fun. We have worked with many surfaces. Which can be topological groups? The torus is, since it is just $S^1 \times S^1$, and S^1 is a topological group. The Klein bottle? Maybe. However, other surfaces cannot since they don't have Euler characteristic 0.