

Part III – Introduction to Discrete Analysis (Ongoing course, rough)

Based on lectures by Professor W. T. Gowers

Notes taken by Bhavik Mehta

Michaelmas 2018

Contents

1	The discrete Fourier transform	2
1.1	Roth's Theorem	4
1.2	Bogolyubov's method	6

1 The discrete Fourier transform

Let N be some fixed positive integer. Write ω for $e^{\frac{2\pi i}{N}}$, and \mathbb{Z}_N for $\mathbb{Z}/N\mathbb{Z}$.

Definition (Discrete Fourier transform). Let $f : \mathbb{Z}_N \rightarrow \mathbb{C}$. Given $r \in \mathbb{Z}_N$, define $\hat{f}(r)$ to be

$$\frac{1}{N} \sum_{x \in \mathbb{Z}_N} f(x) \omega^{-rx}.$$

Notation. From now on, we shall use notation $\mathbb{E}_{x \in \mathbb{Z}_N}$ for $\frac{1}{N} \sum_{x \in \mathbb{Z}_N}$, where the subscript is omitted when it is clear from context.

Notice we can write

$$\hat{f}(r) = \mathbb{E}_x f(x) e^{-\frac{2\pi i r x}{N}},$$

highlighting the similarity with the usual Fourier transform.

If we write ω_r for the function $x \mapsto \omega^{rx}$, and $\langle f, g \rangle$ for $\mathbb{E}_x f(x) \overline{g(x)}$, then $\hat{f}(r) = \langle f, \omega_r \rangle$. Let us write $\|f\|_p$ for $(\mathbb{E}_x |f(x)|^p)^{\frac{1}{p}}$ and call the resulting space $L_p(\mathbb{Z}_N)$.

Important convention. We use *averages* for the ‘original functions’ in ‘physical space’ and *sums* for their Fourier transforms in ‘frequency space’.

Lemma 1 (Parseval’s identity). If $f, g : \mathbb{Z}_N \rightarrow \mathbb{C}$, then $\langle \hat{f}, \hat{g} \rangle = \langle f, g \rangle$.

Proof.

$$\begin{aligned} \langle \hat{f}, \hat{g} \rangle &= \sum_r \hat{f}(r) \overline{\hat{g}(r)} \\ &= \sum_r (\mathbb{E}_x f(x) \omega^{-rx}) \overline{(\mathbb{E}_y g(y) \omega^{-ry})} \\ &= \mathbb{E}_x \mathbb{E}_y f(x) \overline{g(y)} \sum_r \omega^{-r(x-y)} \\ &= \mathbb{E}_x \mathbb{E}_y f(x) \overline{g(y)} \Delta_{xy} \\ &= \mathbb{E}_x f(x) \mathbb{E}_y \overline{g(y)} \Delta_{xy} \\ &= \mathbb{E}_x f(x) \overline{g(x)} = \langle f, g \rangle \end{aligned}$$

where

$$\Delta_{xy} = \begin{cases} N & x = y \\ 0 & x \neq y. \end{cases}$$

□

Definition (Convolution). The convolution $\widehat{f * g}(x)$ is defined to be

$$\mathbb{E}_{y+z=x} f(y) g(z) = \mathbb{E}_y f(y) g(x-y).$$

Lemma 2 (Convolution identity).

$$\widehat{f * g}(r) = \hat{f}(r)\hat{g}(r).$$

Proof.

$$\begin{aligned}\widehat{f * g}(r) &= \mathbb{E}_x f * g(x) \omega^{-rx} \\ &= \mathbb{E}_x \mathbb{E}_{y+z=x} f(y)g(z) \omega^{-rx} \\ &= \mathbb{E}_x \mathbb{E}_{y+z=x} f(y)g(z) \omega^{-ry} \omega^{-rz} \\ &= \mathbb{E}_y f(y) \omega^{-ry} \mathbb{E}_z g(z) \omega^{-rz} = \hat{f}(r)\hat{g}(r).\end{aligned}\quad \square$$

Lemma 3 (Inversion formula).

$$f(x) = \sum_r \hat{f}(r) \omega^{rx}$$

Proof.

$$\begin{aligned}\sum_r \hat{f}(r) \omega^{rx} &= \sum_r \mathbb{E}_y f(y) \omega^{r(x-y)} \\ &= \mathbb{E}_y f(y) \sum_r \omega^{r(x-y)} \\ &= \mathbb{E}_y f(y) \Delta_{xy} = f(x).\end{aligned}\quad \square$$

Further observations:

- If f is real-valued, then $\hat{f}(-r) = \mathbb{E}_x f(x) \omega^{rx} = \overline{\mathbb{E}_x f(x) \omega^{-rx}} = \overline{\hat{f}(r)}$.
- If $A \subset \mathbb{Z}_n$, write A (instead of $\mathbb{1}_A$ or χ_A) for the characteristic function of A . Then $\hat{A}(0) = \mathbb{E}_x A(x) = \frac{|A|}{N}$, the density of A .
- Also, $\|\hat{A}\|_2^2 = \langle \hat{A}, \hat{A} \rangle = \langle A, A \rangle = \mathbb{E}_x A(x)^2 = \mathbb{E}_x A(x) = \frac{|A|}{N}$.

Let $f : \mathbb{Z}_N \rightarrow \mathbb{C}$. Given $\mu \in \mathbb{Z}_N$ with $(\mu, N) = 1$, define $f_\mu(x)$ to be $f(\mu^{-1}x)$. Then

$$\begin{aligned}\hat{f}_\mu(r) &= \mathbb{E}_x f_\mu(x) \omega^{-rx} \\ &= \mathbb{E}_x f(x/\mu) \omega^{-rx} \\ &= \mathbb{E}_x f(x) \omega^{-r\mu x} \\ &= \hat{f}(\mu r).\end{aligned}$$

1.1 Roth's Theorem

Theorem 4. For every $\delta > 0$, there exists N such that if $A \subseteq \{1, \dots, N\}$ is a set of size at least δN then A must contain an arithmetic progression of length 3.

This is the $k = 3$ case of Szemerédi's theorem.

Basic strategy: show that if A has density $\geq \delta$ and no arithmetic progression of length 3, then there is a long arithmetic progression $P \subseteq \{1, \dots, N\}$ such that

$$|A \cap P| \geq (\delta + c(\delta))|P|.$$

In particular, we have that $|P| \rightarrow \infty$ as $N \rightarrow \infty$.

The proof we give will produce a bound $\delta \geq \frac{C}{\log \log N}$, but this is not the best known. If the bound was reduced to $\frac{1}{\log N}$, this produces a combinatorial proof of the fact that there are arbitrarily long arithmetic progressions in the primes. The best known bound is $\frac{(\log \log N)^4}{\log N}$ by Thomas Bloom. In the other direction, we know $e^{-\sqrt{\log N}}$ does not work.

Lemma 5. Let $A, B, C \subset \mathbb{Z}_N$ have densities α, β, γ , for N odd. If $\max_{r \neq 0} |\hat{A}(r)| \leq \frac{\alpha(\beta\gamma)^{\frac{1}{2}}}{2}$ and $\frac{\alpha\beta\gamma}{2} > \frac{1}{N}$ then there exists $x, d \in \mathbb{Z}_N$ with $d \neq 0$ such that $(x, x+d, x+2d) \in A \times B \times C$.

Proof.

$$\begin{aligned} \mathbb{E}_{x,d} A(x)B(x+d)C(x+2d) &= \mathbb{E}_{x+z=2y} A(x)B(y)C(z) \\ &= \mathbb{E}_u \left(\mathbb{E}_{x+z=u} A(x)C(z) \right) \mathbb{E}_{2y=u} B(y) \\ &= \mathbb{E}_u (A * C)(u) B_2(u) = \langle A * C, B_2 \rangle \\ &= \langle \widehat{A * C}, \hat{B}_2 \rangle \\ &= \langle \hat{A} \hat{C}, \hat{B}_2 \rangle \\ &= \sum_r \hat{A}(r) \hat{C}(r) \hat{B}(-2r) \\ &= \alpha\beta\gamma + \sum_{r \neq 0} \hat{A}(r) \hat{C}(r) \hat{B}(-2r). \end{aligned}$$

We have a lower bound on the left term, so focus on the right.

$$\begin{aligned} \left| \sum_{r \neq 0} \hat{A}(r) \hat{B}(-2r) \hat{C}(r) \right| &\leq \frac{\alpha(\beta\gamma)^{\frac{1}{2}}}{2} \sum_{r \neq 0} |\hat{B}(-2r) \hat{C}(r)| \\ &\leq \frac{\alpha(\beta\gamma)^{\frac{1}{2}}}{2} \left(\sum_r |\hat{B}(-2r)|^2 \right)^{\frac{1}{2}} \left(\sum_r |\hat{C}(r)|^2 \right)^{\frac{1}{2}} \\ &= \frac{\alpha(\beta\gamma)^{\frac{1}{2}}}{2} \|\hat{B}\|_2 \|\hat{C}\|_2 = \frac{\alpha(\beta\gamma)^{\frac{1}{2}}}{2} \|B\|_2 \|C\|_2 \\ &= \frac{\alpha\beta\gamma}{2}. \end{aligned}$$

The contribution to $\mathbb{E}_{x,d} A(x)B(x+d)C(x+2d)$ from $d=0$ is at most $\frac{1}{N}$, so if $\frac{\alpha\beta\gamma}{2} > \frac{1}{N}$, we are done. \square

Now let A be a subset of $\{1, \dots, N\}$ of density $\geq \delta$ and let $B = C = A \cap (\frac{N}{3}, \frac{2N}{3}]$. If B has density $< \frac{\delta}{5}$, then either $A \cap [1, \frac{N}{3}]$ or $A \cap [\frac{2N}{3}, N]$ has density at least $\frac{2\delta}{5}$. So in that case we find an AP P of length about $\frac{N}{3}$ such that $\frac{|A \cap P|}{|P|} \geq \frac{6\delta}{5}$.

Otherwise, we find that if $\max_{r \neq 0} |\hat{A}(r)| \leq \frac{\delta^2}{10}$ and $\frac{\delta^3}{50} > \frac{1}{N}$ then $A \times B \times C$ contains a 3AP $\implies A$ contains a 3AP. So if A does not contain a 3AP, then either we find P of length about $\frac{N}{3}$ with $\frac{|A \cap P|}{|P|} \geq \frac{6\delta}{5}$ or $\exists r \neq 0$ such that $|\hat{A}(r)| \geq \frac{\delta^2}{10}$.

Definition. If X is a finite set and $f : X \rightarrow \mathbb{C}$, $Y \subseteq X$, write $\text{osc}(f|_Y)$ to mean $\max_{y_1, y_2 \in Y} |f(y_1) - f(y_2)|$.

Lemma 6. Let $r \in \hat{\mathbb{Z}}_n$ and let $\epsilon > 0$. Then there is a partition of $\{1, 2, \dots, N\}$ into arithmetic progressions P_i of length at least $c(\epsilon)\sqrt{N}$ such that $\text{osc}(\omega_r|_{P_i}) \leq \epsilon$ for each i .

Proof. Let $t = \lfloor \sqrt{N} \rfloor$. Of the numbers $1, \omega^r, \omega^{2r}, \dots, \omega^{tr}$ there must be two that differ by at most $\frac{2\pi}{t}$. If $|\omega^{ar} - \omega^{br}| \leq \frac{2\pi}{t}$ with $a < b$, then $|1 - \omega^{dr}| \leq \frac{2\pi}{t}$ where $d = b - a$. Now, by the triangle inequality, if $u < v$, then

$$|\omega^{urd} - \omega^{vrd}| \leq |\omega^{urd} - \omega^{(u+1)rd}| + |\omega^{urd} - \omega^{(u+1)rd}| + \dots + |\omega^{urd} - \omega^{(u+1)rd}| \leq \frac{2\pi}{t}(v - u).$$

So if P is a progression with common difference d and length l , then $\text{osc}(\omega_r|_P) \leq \frac{2\pi l}{t}$. So divide up $\{1, \dots, N\}$ into residue classes mod d and partition each residue class into parts of length between $\frac{\epsilon t}{4\pi}$ and $\frac{\epsilon t}{2\pi}$ (possible, since $d \leq t \leq \sqrt{N}$). We are done, with $c(\epsilon) = \frac{\epsilon}{16}$. \square

Now let us use the information that $r \neq 0$ and $|\hat{A}(r)| \geq \frac{\delta^2}{10}$. Define the balanced function f of A by $f(x) = A(x) - \frac{|A|}{N}$ for each x .

Note that $\hat{f}(0) = 0$ and $\hat{f}(r) = \hat{A}(r)$ for all $r \neq 0$. Now let P_1, \dots, P_m be given by [Lemma 6](#) with $\epsilon = \frac{\delta^2}{20}$. Then

$$\begin{aligned} \frac{\delta^2}{10} &\leq \frac{1}{N} \left| \sum_x f(x) \omega^{-rx} \right| \leq \frac{1}{N} \sum_{i=1}^m \left| \sum_{x \in P_i} f(x) \omega^{-rx} \right| \\ &\leq \frac{1}{N} \sum_{i=1}^m \left[\left| \sum_{x \in P_i} f(x) \omega^{-rx_i} \right| + \left| \sum_{x \in P_i} f(x) (\omega^{-rx} - \omega^{-rx_i}) \right| \right] \end{aligned}$$

where $x_i \in P_i$ is arbitrary

$$\leq \frac{1}{N} \sum_{i=1}^m \left| \sum_{x \in P_i} f(x) \right| + \frac{\delta^2}{20}$$

So

$$\sum_{i=1}^N \left| \sum_{x \in P_i} f(x) \right| \geq \frac{\delta^2 N}{20}.$$

Also,

$$\sum_{i=1}^m \sum_{x \in P_i} f(x) = 0.$$

So

$$\sum_{i=1}^m \left(\left| \sum_{x \in P_i} f(x) \right| + \sum_{x \in P_i} f(x) \right) \geq \frac{\delta^2}{20} \sum_{i=1}^m |P_i|$$

Therefore, $\exists i$ such that

$$\begin{aligned} \left| \sum_{x \in P_i} f(x) \right| + \sum_{x \in P_i} f(x) &\geq \frac{\delta^2}{20} |P_i| \\ \implies \sum_{x \in P_i} f(x) &\geq \frac{\delta}{40} |P_i| \\ \implies |A \cap P_i| &\geq \left(\delta + \frac{\delta^2}{40} \right) |P_i| \end{aligned}$$

So now, either

1. A contains a $3AP$
2. N is even
3. $\exists P \subset \{1, \dots, N\}$, $|P| \geq \frac{N}{3}$ such that $|A \cap P| \geq \frac{6\delta}{5} |P|$
4. $\exists P \subset \{1, \dots, N\}$, $|P| \geq \frac{\delta^2}{320} \sqrt{N}$ such that $|A \cap P| \geq \left(\delta + \frac{\delta^2}{40} \right) |P|$

If 2 holds, write $N = N_1 + N_2$ with N_1, N_2 odd, $N_1 + N_2 \approx \frac{N}{2}$. Then A has density at least δ in one of $\{1, \dots, N_1\}$ or $\{N_1 + 1, \dots, N_1 + N_2\}$.

If 4 holds (note $3 \Rightarrow 4$) then we pass to P and start again. After $\frac{40}{\delta}$ iterations, the density at least doubles. So the total number of iterations we can have is $\leq \frac{40}{\delta} + \frac{40}{2\delta} + \frac{40}{4\delta} + \dots \leq \frac{80}{\delta}$.

If $\frac{\delta^2}{320} \sqrt{N} \geq N^{\frac{1}{3}}$ at each iteration, and $\frac{\delta^3}{25} > N^{-1}$ (which follows from the first condition) then after $\frac{80}{\delta}$ iterations we have $N \geq N^{\left(\frac{1}{3}\right)^{\frac{80}{\delta}}}$. So the argument works provided

$$\begin{aligned} N^{\left(\frac{1}{3}\right)^{\frac{80}{\delta}}} &\geq \left(\frac{320}{\delta^2} \right)^6 \iff \left(\frac{1}{3} \right)^{\frac{80}{\delta}} \log N \geq 6 \left(\log 320 + 2 \log \frac{1}{\delta} \right) \\ &\iff -\frac{80}{\delta} \log 3 + \log \log N \geq \log 6 + \log \left(\log 320 + 2 \log \frac{1}{\delta} \right) \\ &\iff \log \log N \geq \frac{160}{\delta} \iff \delta \geq \frac{160}{\log \log N}. \end{aligned}$$

1.2 Bogolyubov's method

Let $K \subset \hat{\mathbb{Z}}_N$ and let $\delta > 0$.

Definition (Bohr set). The **Bohr set** $B(K, \delta)$ has two definitions.

1. $B(K, \delta) = \{x \in \mathbb{Z}_N \mid rx \in [-\delta N, \delta N] \ \forall r \in K\}$ (arc-length definition)
2. $B(K, \delta) = \{x \in \mathbb{Z}_N \mid |1 - \omega^{rx}| < \delta \ \forall r \in K\}$ (chord-length definition)

Definition. Let G be an abelian group and let A, B be subsets of G . Then

$$\begin{aligned} A + B &= \{a + b \mid a \in A, b \in B\} \\ A - B &= \{a - b \mid a \in A, b \in B\} \\ rA &= \{a_1 + \dots + a_r \mid a_1, \dots, a_r \in A\} \end{aligned}$$

Lemma 7. Let $A \subset \mathbb{Z}_N$ be a set of density α . Then $2A - 2A$ contains a [Bohr set](#) (arc) with $|K| \geq \alpha^{-2}$.

Proof. Observe that $x \in 2A - 2A$ iff $A * A * (-A) * (-A)(x) \neq 0$. But

$$\begin{aligned} A * A * (-A) * (-A)(x) &= \sum_r \overline{A * A * (-A) * (-A)(r)} \omega^{rx} \\ &= \sum_r |\hat{A}(r)|^4 \omega^{rx}. \end{aligned}$$

Let $K = \{r \mid |\hat{A}(r)| \geq \alpha^{\frac{3}{2}}\}$. Then $\alpha = \|\hat{A}\|_2^2 = \sum_r |\hat{A}(r)|^2 \geq \alpha^3 |K|$ So $|K| \leq \alpha^{-2}$.

Now suppose that $x \in B(K, \frac{1}{4})$. Then

$$\sum_r |\hat{A}(r)|^4 \omega^{rx} = \alpha^4 + \sum_{r \in K \setminus \{0\}} |\hat{A}(r)|^4 \omega^{rx} + \sum_{r \notin K} |\hat{A}(r)|^4 \omega^{rx}.$$

The real part of the second term is non-negative, since $rx \in [-\frac{N}{4}, \frac{N}{4}]$ when $r \in K$. Also

$$\left| \sum_{r \notin K} |\hat{A}(r)|^4 \omega^{rx} \right| \leq \sum_{r \notin K} |\hat{A}(r)|^4 < \alpha^3 \sum_{r \notin K} |\hat{A}(r)|^2 \leq \alpha^4.$$

It follows that $\operatorname{Re} \left(\sum_r |\hat{A}(r)|^4 \omega^{rx} \right) > 0$, so $x \in 2A - 2A$. □

Lemma 8. Let $K \subset \mathbb{Z}_N$ and let $\delta > 0$. Then

- (i) $B(K, \delta)$ (arc) has density at least $\delta^{|K|}$
- (ii) $B(K, \delta)$ contains a mod- N arithmetic progression of length $\geq \delta N^{\frac{1}{|K|}}$

Proof.

- (i) Let $K = \{r_1, \dots, r_k\}$. Consider the N k -tuples $(r_1 x, r_2 x, \dots, r_k x) \in \mathbb{Z}_N^k$. If we intersect this set of k -tuples with a random ‘box’ $[t_1, t_1 + \delta N] \times \dots \times [t_k, t_k + \delta N]$ then the expected number of the k -tuples in the box is $\delta^k N$ (since each one has a probability δ^k). But if $(r_1 x, \dots, r_k x)$ and $(r_1 y, \dots, r_k y)$ belong to this box, then $x - y \in B(K, \delta)$.

(ii) If we take $\eta > N^{\frac{1}{2}}$, then by (i) we get that $|B(K, \eta)| > 1$, so $\exists x \in B(K, \eta)$ such that $x \neq 0$. But then $dx \in B(K, d\eta)$ for every d . So if $d\eta \leq \delta$ then $dx \in B(K, \delta)$. That gives us an AP of length at least $\frac{\delta}{\eta}$. So we get one of length at least $\delta N^{\frac{1}{k}}$.

□