

Part III – Model Theory (Ongoing course, rough)

Based on lectures by Dr. S. Barbina

Notes taken by Bhavik Mehta

Michaelmas 2018

Contents

0	Introduction	2
1	Languages and structures	2
2	Review: Terms, formulae and their interpretations	4
3	Theories and elementarity	7
	Index	8

0 Introduction

Lecture 1 Model theory is a part of logic that began by looking at algebraic objects such as groups and combinatorial objects such like graphs, described in formal language. The basic question in model theory is: ‘how powerful is our description of these objects to pin them down’? In Logic and Set Theory, the focus was on what was provable from a theory and language, but here we focus on whether or not a model exists.

1 Languages and structures

Definition 1.1 (Language). A **language** L consists of

- (i) a set \mathcal{F} of function symbols, and for each $f \in \mathcal{F}$ a positive integer m_f the **arity** of f .
- (ii) a set \mathcal{R} of relation symbols, and for each $R \in \mathcal{R}$, a positive integer m_R .
- (iii) a set \mathcal{C} of constant symbols.

Note: each of \mathcal{F} , \mathcal{R} and \mathcal{C} can be empty.

Example. Take $L = \{\{\cdot, {}^{-1}\}, \{1\}\}$, for \cdot a binary function and ${}^{-1}$ an unary function, 1 a constant. This is the **language** of groups, call it L_{gp} . Also, $L_{lo} = \{<\}$ a single binary relation, for linear orders.

Definition 1.2 (L -structure). Given a **language** L , say, an **L -structure** consists of

- (i) a set M , the **domain**
- (ii) for each $f \in \mathcal{F}$, a function $f^{\mathcal{M}} : M^{m_f} \rightarrow M$.
- (iii) for each $R \in \mathcal{R}$, a relation $R^{\mathcal{M}} \subseteq M^{m_R}$.
- (iv) for each $c \in \mathcal{C}$, an element $c^{\mathcal{M}} \in M$.

$f^{\mathcal{M}}, R^{\mathcal{M}}, c^{\mathcal{M}}$ are the **interpretations** of f, R, c respectively.

Remark 1.3. We often fail to distinguish between the **symbols** in L and their **interpretations** in a **structure**, if the interpretations are clear from the context.

We may write $\mathcal{M} = \langle M, \mathcal{F}, \mathcal{R}, \mathcal{C} \rangle$.


Example 1.4.

- (a) $\mathcal{R} = \langle \mathbb{R}^+, \{\cdot, {}^{-1}\}, 1 \rangle$ is an L_{gp} -**structure**.
- (b) $\mathcal{Z} = \langle \mathbb{Z}, \{+, -\}, 0 \rangle$ is an L_{gp} -**structure**.
- (c) $\mathcal{Q} = \langle \mathbb{Q}, < \rangle$ is an L_{lo} -**structure**.

Definition 1.5 (Embedding). Let L be a **language**, let \mathcal{M}, \mathcal{N} be **L -structures**. An **embedding** of \mathcal{M} into \mathcal{N} is a one-to-one mapping $\alpha : M \rightarrow N$ such that

- (i) for all $f \in \mathcal{F}$, and $a_1, \dots, a_{m_f} \in M$,

$$\alpha(f^{\mathcal{M}}(a_1, \dots, a_{m_f})) = f^{\mathcal{N}}(\alpha(a_1), \dots, \alpha(a_{m_f}))$$



(ii) for all $R \in \mathcal{R}$, and $a_1, \dots, a_{n_R} \in M$

$$(a_1, \dots, a_{n_R}) \in R^{\mathcal{M}} \iff (\alpha(a_1), \dots, \alpha(a_{n_R})) \in R^{\mathcal{N}}$$

(iii) for all $c \in \mathcal{C}$, $\alpha(c^{\mathcal{M}}) = c^{\mathcal{N}}$.

An **isomorphism** of \mathcal{M} into \mathcal{N} is a surjective embedding (onto).

Exercise 1.6. Let G_1, G_2 be groups, regarded as L_{gp} -structures. Check that $G_1 \simeq G_2$ in the usual algebra sense if and only if there is an **isomorphism** $\alpha : G_1 \rightarrow G_2$ in the sense of [Definition 1.5](#).

2 Review: Terms, formulae and their interpretations

In addition to the [symbols](#) of L , we also have

- (i) infinitely many variables $\{x_i\}_{i \in I}$
- (ii) logical connectives \wedge, \neg (also expresses \vee, \implies, \iff)
- (iii) quantifier \exists (also expresses \forall)
- (iv) $(\ , \)$
- (v) equality symbol $=$

Definition 2.1 (L -terms). L -terms are defined recursively as follows:

- any variable x_i is a term
- any constant symbol is a term
- for any $f \in \mathcal{F}$, $f(t_1, \dots, t_{m_f})$ for any terms t_1, \dots, t_{m_f} is a term
- nothing else is a term

Notation: we write $t(x_1, \dots, x_m)$ to mean that the variables appearing in t are among x_1, \dots, x_m .

Lecture 2 **Example.** Take $\mathcal{R} = \langle \mathbb{R}^*, \{\cdot, ^{-1}\}, 1 \rangle$. Then $\cdot((x_1, x_2), x_3)$ is a [term](#), usually written $(x_1 \cdot x_2) \cdot x_3$. Also, $(\cdot(1, x_1))^{-1}$ is a [term](#), written $(1 \cdot x)^{-1}$

Definition 2.2. If \mathcal{M} is an L -structure, to each L -term $t(x_1, \dots, x_k)$ we assign a function a function $t^{\mathcal{M}} : M^k \rightarrow M$ defined as follows:

- (i) If $t = x_i$, $t^{\mathcal{M}}[a_1, \dots, a_k] = a_i$
- (ii) If $t = c$, $t^{\mathcal{M}}[a_1, \dots, a_k] = c^{\mathcal{M}}$.
- (iii) If $t = f(t_1(x_1, \dots, x_k), \dots, t_{m_f}(x_1, \dots, x_k))$,

$$t^{\mathcal{M}}(a_1, \dots, a_k) = f^{\mathcal{M}}(t_1^{\mathcal{M}}(a_1, \dots, a_k), \dots, t_{m_f}^{\mathcal{M}}(a_1, \dots, a_k))$$

Notice in L_{gp} , the term $x_2 \cdot x_3$ can be described as $t_1(x_1, x_2, x_3)$ or $t_2(x_1, x_2, x_3, x_4)$, or infinitely many other ways. Then t_1 is assigned to $t_1^{\mathcal{M}} : M^3 \rightarrow M$, with $(a_1, a_2, a_3) \mapsto (a_2, a_3)$, and t_2 is assigned to $t_2^{\mathcal{M}} : M^4 \rightarrow M$, with $(a_1, a_2, a_3, a_4) \mapsto a_2 \cdot a_3$.

Fact 2.3. Let \mathcal{M}, \mathcal{N} be L -structures, and let $\alpha : \mathcal{M} \rightarrow \mathcal{N}$ be an [embedding](#). For any L -term $t(x_1, \dots, x_k)$ and $a_1, \dots, a_k \in M$ we have

$$\alpha(t^{\mathcal{M}}(a_1, \dots, a_k)) = t^{\mathcal{N}}(\alpha(a_1), \dots, \alpha(a_k))$$

Proof. By induction on the complexity of t . Let $\bar{a} = (a_1, \dots, a_k)$ and $\bar{x} = (x_1, \dots, x_k)$. Then

- (i) if $t = x_i$, then $t^{\mathcal{M}}(\bar{a}) = a_i$, and $t^{\mathcal{N}}(\alpha(a_1), \dots, \alpha(a_k)) = \alpha(a_i)$, so the conclusion holds.

- (ii) if $t = c$ a constant, then $t^{\mathcal{M}}(\bar{a}) = c^{\mathcal{M}}$, and $t^{\mathcal{N}}(\alpha(\bar{a})) = c^{\mathcal{N}}$, and $\alpha(c^{\mathcal{M}}) = c^{\mathcal{N}}$, as required.
- (iii) if $t = f(t_1(\bar{x}), \dots, t_{m_f}(\bar{x}))$, then

$$\alpha(f^{\mathcal{M}}(t_1^{\mathcal{M}}(\bar{a}), \dots, t_{m_f}^{\mathcal{M}}(\bar{a}))) = f^{\mathcal{N}}(\alpha(t_1^{\mathcal{M}}(\bar{a})), \dots, \alpha(t_{m_f}^{\mathcal{M}}(\bar{a})))$$

since α is an embedding. $t_1(\bar{x}), \dots, t_{m_f}(\bar{x})$ have lower complexity than t , so inductive hypothesis applies.

□

Example 2.4. Exercise: conclude the proof of [Fact 2.3](#).

Definition 2.5 (Atomic formula). The set of **atomic formulas** of L is defined as follows

- (i) if t_1, t_2 are L -terms, then $t_1 = t_2$ is an atomic formula
- (ii) if R is a relation symbol and t_1, \dots, t_{m_R} are terms, then $R(t_1, \dots, t_{m_R})$ is an atomic formula
- (iii) nothing else is an atomic formula.

Definition 2.6 (Formula). The set of L -**formulas** is defined as follows

- (i) any **atomic formula** is an L -formula
- (ii) if ϕ is an L -formula, then so is $\neg\phi$
- (iii) if ϕ and ψ are L -formulas, then so is $\phi \wedge \psi$
- (iv) if ϕ is an L -formula, for any $i \geq 1$, $\exists x_i \phi$ is an L -formula
- (v) nothing else is an L -formula

Example. In L_{gp} , $x_1 \cdot x_1 = x_2$ and $x_1 \cdot x_2 = 1$ are **atomic formulas**, and $\exists x_1(x_1 \cdot x_2) = 1$ is an L_{gp} -formula.

A variable occurs freely in a formula if it does not occur within the scope of a quantifier \exists (the variable is **free**). Otherwise the variable is **bound**. For instance, in $\exists x_1(x_1 \cdot x_2) = 1$, x_1 is bound and x_2 is free.

Important convention: no variable occurs both **freely** and as a bound variable in the same formula.

A **sentence** is a **formula** with no **free** variables. $\exists x_1 \exists x_2(x_1 \cdot x_2 = 1)$ is an L_{gp} -sentence.

Notation: $\phi(x_1, \dots, x_k)$ means that the free variables in ϕ are among x_1, \dots, x_k .

Definition 2.7 (\models). Let $\phi(x_1, \dots, x_k)$ be an L -formula, let \mathcal{M} be an L -structure, and let $\bar{a} = (a_1, \dots, a_k)$ be elements of \mathcal{M} . We define $\mathcal{M} \models \phi(\bar{a})$ as follows.

- (i) if ϕ is $t_1 = t_2$, then $\mathcal{M} \models \phi(\bar{a})$ if and only if $t_1^{\mathcal{M}}(\bar{a}) = t_2^{\mathcal{M}}(\bar{a})$.
- (ii) if ϕ is $R(t_1, \dots, t_{m_k})$ then $\mathcal{M} \models \phi(\bar{a})$ iff

$$(t_1^{\mathcal{M}}(\bar{a}), \dots, t_{m_k}^{\mathcal{M}}(\bar{a})) \in R^{\mathcal{M}}.$$

- (iii) if ϕ is $\psi \wedge \chi$, then $\mathcal{M} \models \phi(\bar{a})$ iff $\mathcal{M} \models \psi(\bar{a})$ and $\mathcal{M} \models \chi(\bar{a})$.
- (iv) if $\phi = \neg\psi$ then $\mathcal{M} \models \phi(\bar{a})$ iff $\mathcal{M} \not\models \psi(\bar{a})$. (this is well-defined since $\psi(\bar{a})$ is shorter than $\phi(\bar{a})$)
- (v) if ϕ is $\exists x_j : \chi(x_1, \dots, x_k, x_j)$ (where $x_j \neq x_i$ for $i = 1, \dots, k$). Then $\mathcal{M} \models \phi(\bar{a})$ iff there is $b \in \mathcal{M}$ such that $\mathcal{M} \models \chi(a_1, \dots, a_k, b)$.

Example. For $\mathcal{R} = \langle \mathbb{R}^*, \cdot, ^{-1}, 1 \rangle$, $\phi(x_1) = \exists x_2 (x_2 \cdot x_2 = x_1)$ then $\mathcal{R} \models \phi(1)$ but $\mathcal{R} \not\models \phi(-1)$.

Notation 2.8 (Useful abbreviations). We write

- $\phi \vee \psi$ for $\neg(\neg\phi \wedge \neg\psi)$
- $\phi \rightarrow \psi$ for $\neg\phi \vee \psi$
- $\phi \leftrightarrow \psi$ for $(\phi \rightarrow \psi) \wedge (\psi \rightarrow \phi)$
- $\forall x_i \phi$ for $\neg\exists x_i (\neg\phi)$

Proposition 2.9. Let \mathcal{M}, \mathcal{N} be *L-structures*, let $\alpha : \mathcal{M} \rightarrow \mathcal{N}$ be an *embedding*. Let $\phi(\bar{x})$ be *atomic* and $\bar{a} \in M^{|\bar{x}|}$, then

$$\mathcal{M} \models \phi(\bar{a}) \iff \mathcal{N} \models \phi(\alpha(\bar{a})).$$

Question: If ϕ is an *L-formula*, not necessarily atomic, does [Proposition 2.9](#) hold?

Lecture 3

Proof of Proposition 2.9. Cases:

- (i) $\phi(\bar{x})$ is of the form $t_1(\bar{x}) = t_2(\bar{x})$ where t_1, t_2 are terms. (Exercise: complete this case, using [Fact 2.3](#))
- (ii) $\phi(\bar{x})$ is of the form $R(t_1(\bar{x}), \dots, t_{m_R}(\bar{x}))$. Then $\mathcal{M} \models R(t_1(\bar{a}), \dots, t_{m_R}(\bar{a}))$ if and only if... (Exercise: complete this case)

□

Exercise 2.10. Show that [Proposition 2.9](#) holds if $\phi(\bar{x})$ is a formula without quantifiers (a quantifier-free formula).

Example 2.11. Do *embeddings* preserve *all formulas*? No. Take $\mathcal{Z} = (\mathbb{Z}, <)$ and $\mathcal{Q} = (\mathbb{Q}, <)$ an *L_{lo}-structure*. Then $\alpha : \mathbb{Z} \rightarrow \mathbb{Q}$ (inclusion) is an embedding, but

$$\phi(x_1, x_2) = \exists x_3 (x_1 < x_3 \wedge x_3 < x_2). \quad (1)$$

$$\mathcal{Q} \models \phi(1, 2) \text{ but } \mathcal{Z} \not\models \phi(1, 2). \quad (2)$$

Fact 2.12. Let $\alpha : \mathcal{M} \rightarrow \mathcal{N}$ be an *isomorphism*. Then if $\phi(\bar{x})$ is an *L-formula* and $\bar{a} \in M^{|\bar{x}|}$, then

$$\mathcal{M} \models \phi(\bar{a}) \iff \mathcal{M} \models \phi(\alpha(\bar{a})).$$

Proof. Exercise.

□

3 Theories and elementarity

Throughout, L is a [language](#), \mathcal{M}, \mathcal{N} are [L-structures](#).

Definition 3.1 (*L-theory*). An *L-theory* T is a set of [L-sentences](#). \mathcal{M} is a **model** of T if $\mathcal{M} \models \sigma$ for all $\sigma \in T$. We write $\mathcal{M} \models T$. The class of all the models of T is written $\text{Mod}(T)$. The theory of \mathcal{M} is the set

$$\text{Th}(\mathcal{M}) = \{ \sigma \mid \sigma \text{ is an } L\text{-structure and } \mathcal{M} \models \sigma \}.$$

Example 3.2. Let T_{gp} be the set of [L_{gp}-sentences](#).

- (i) $\forall x_1 x_2 x_3 (x_1 \cdot (x_2 \cdot x_3) = (x_1 \cdot x_2) \cdot x_3)$
- (ii) $\forall x_1 (x_1 \cdot 1 = 1 \cdot x_1 = x_1)$
- (iii) $\forall x_1 (x_1 \cdot x_1^{-1} = x_1^{-1} \cdot x_1 = 1)$

Clearly for a group G , $G \models T_{gp}$. For a specific G , clearly $\text{Th}(G)$ is larger than T_{gp} !

Definition 3.3. —hyperpage

We'll see $(\mathbb{Q}, <) \equiv (\mathbb{R}, <)$ as L_{lo} -structures.

Definition 3.4.

- (i) an embedding $\beta : \mathcal{M} \rightarrow \mathcal{N}$ is **elementary** if for all formulas $\phi(\bar{x})$ and $\bar{a} \in M^{|\bar{x}|}$,

$$\mathcal{M} \models \phi(\bar{a}) \iff \mathcal{N} \models \phi(\beta(\bar{a}))$$

- (ii) if $M \subseteq N$ and $\text{id} : \mathcal{M} \rightarrow \mathcal{N}$ is an embedding, then \mathcal{M} is said to be a **substructure** of \mathcal{N} .
- (iii) if $M \subseteq N$ and $\text{id} : \mathcal{M} \rightarrow \mathcal{N}$ is an elementary embedding, then \mathcal{M} is said to be an **elementary substructure** of \mathcal{N} , written $\mathcal{M} \preceq \mathcal{N}$.

Example 3.5. Consider $\mathcal{M} = [0, 1] \subseteq \mathbb{R}$, an L_{lo} -structure, where $<$ is the usual order, and $\mathcal{N} = [0, 2] \subseteq \mathbb{R}$ in the same way. Then $\mathcal{M} \simeq \mathcal{N}$ as L_{lo} -structures. Is $\mathcal{M} \equiv \mathcal{N}$? Yes they are isomorphic!

Is $\mathcal{M} \subseteq \mathcal{N}$? Yes (the ordering $<$ coincides on \mathcal{M} and \mathcal{N} .) But $\mathcal{M} \not\preceq \mathcal{N}$, since if $\phi(x) = \exists y (x < y)$, then

$$\mathcal{N} \models \phi(1) \quad \text{and} \quad \mathcal{M} \not\models \phi(1).$$

Definition 3.6. Let \mathcal{M} an L -structure, $A \subseteq M$, then

$$L(A) := L \cup \{ c_a : a \in A \}$$

for c_a each constant symbols. An interpretation of \mathcal{M} as an L -structure extends to an interpretation of \mathcal{M} as an $L(A)$ -structure in the obvious way ($c_a^{\mathcal{M}} = a$). The elements of A are called **parameters**. If \mathcal{M}, \mathcal{N} are L -structures and $A \subseteq M \cap N$, then $\mathcal{M} \equiv_A \mathcal{N}$ when \mathcal{M}, \mathcal{N} satisfy exactly the same $L(A)$ sentences.

Index

\models , [5](#)

bound variable, [5](#)

embedding, [2](#)

formula, [5](#)
 atomic, [5](#)

free variable, [5](#)

L -structure, [2](#)

L -term, [4](#)

L -theory, [7](#)

language, [2](#)

sentence, [5](#)