

Part III – Category Theory (Ongoing course)

Based on lectures by Professor P. T. Johnstone

Notes taken by Bhavik Mehta

Michaelmas 2018

Contents

0	Introduction	2
1	Definitions and Examples	2
2	The Yoneda Lemma	9
	Index	12

0 Introduction

Lecture 1 Category theory is like a language spoken by many different people, with many different dialects. Specifically, different parts of category theory are used in different branches of mathematics. In this course, we aim to speak the language of category theory, without an accent - a broad overview of all aspects of category theory. There will be many examples, some of which may not be understandable. As long as some examples make sense, it is not a point of concern that some examples seem unfamiliar.

1 Definitions and Examples

1.1 Definition (Category). A **category** \mathcal{C} consists of

- (a) a collection $\text{ob } \mathcal{C}$ of **objects** A, B, C, \dots
- (b) a collection $\text{mor } \mathcal{C}$ of **morphisms** f, g, h, \dots
- (c) two operations dom, cod assigning to each $f \in \text{mor } \mathcal{C}$ a pair of objects, its **domain** and **codomain**. We write $A \xrightarrow{f} B$ to mean ‘ f is a morphism and $\text{dom } f = A$ and $\text{cod } f = B$ ’.
- (d) an operation assigning to each $A \in \text{ob } \mathcal{C}$ a morphism $A \xrightarrow{1_A} A$, called its **identity**.
- (e) a partial binary operation **composition** $(f, g) \mapsto fg$ on morphisms, such that fg is defined iff $\text{dom } f = \text{cod } g$ and $\text{dom}(fg) = \text{dom } g$, $\text{cod}(fg) = \text{cod } f$ if fg is defined.

satisfying

- (f) $f1_A = f = 1_B f$ for any $A \xrightarrow{f} B$
- (g) $(fg)h = f(gh)$ whenever fg and gh are defined

1.2 Remark.

- (a) This definition is independent of a model of set theory. If we’re given a particular model of set theory, we call the **category** \mathcal{C} **small** if $\text{ob } \mathcal{C}$ and $\text{mor } \mathcal{C}$ are sets.
- (b) Some texts say fg means ‘ f followed by g ’, i.e. fg defined $\iff \text{cod } f = \text{dom } g$.
- (c) Note that a morphism f is an **identity** iff $fg = g$ and $hf = h$ whenever the compositions are defined. So we could formulate the definition entirely in terms of morphisms.

1.3 Examples.

- (a) The **category Set** has all sets as objects, and all functions between sets as morphisms. (Strictly, morphisms $A \longrightarrow B$ are pairs (f, B) where f is a set-theoretic function.)
- (b) The category **Gp** has all groups as objects, and group homomorphisms as morphisms. Similarly, **Rng** is the category of rings, **Mod** $_R$ the category of R -modules.
- (c) The category **Top** has all topological spaces as objects and continuous functions as morphisms. Similarly **Unif** has uniform spaces and uniformly continuous functions, and **Mf** has manifolds and smooth maps.

- (d) The category **Htpy** has the same objects as **Top**, but morphisms are homotopy classes of continuous functions. More generally, given \mathcal{C} , we call an equivalence relation \simeq on $\text{mor } \mathcal{C}$ a **congruence** if $f \simeq g \implies \text{dom } f = \text{dom } g$ and $\text{cod } f = \text{cod } g$, and $f \simeq g \implies fh \simeq gh$ and $kf \simeq kg$ whenever the composites are defined. Then we have a category \mathcal{C}/\simeq with the same objects as \mathcal{C} , but congruence classes as morphisms.
- (e) Given \mathcal{C} , the **opposite category** \mathcal{C}^{op} has the same objects and morphisms as \mathcal{C} , but dom and cod are interchanged, and fg in \mathcal{C}^{op} is gf in \mathcal{C} . This leads to the **Duality principle** if P is a true statement about categories, so is the statement P^* obtained from P by reversing all arrows.
- (f) A **small** category with one object is a **monoid**, i.e. a semigroup with 1. In particular, a group is a small category with one object, in which every morphism is an isomorphism (f is an **isomorphism** if $\exists g$ such that fg and gf are identities).
- (g) A **groupoid** is a category in which every morphism is an isomorphism. For a topological space X , the fundamental groupoid $\pi(X)$ has all points of X as objects and morphisms $x \longrightarrow y$ are homotopy classes $\text{rel } \{0, 1\}$ of paths $u : [0, 1] \longrightarrow X$ with $u(0) = x$, $u(1) = y$. (If you know how to prove that the fundamental group is a group, you can prove that $\pi(X)$ is a groupoid.)
- (h) A **discrete** category is one whose only morphisms are identities. A **preorder** is a category \mathcal{C} in which, for any pair (A, B) there is at most 1 morphism $A \longrightarrow B$. A small preorder is a set equipped with a binary relation which is reflexive and transitive. In particular, a partially ordered set is a small preorder in which the only isomorphisms are identities.
- (i) The category **Rel** has the same objects as **Set**, but morphisms $A \longrightarrow B$ are arbitrary relations $R \subseteq A \times B$. Given R and $S \subseteq B \times C$, we define

$$S \circ R = \{ (a, c) \in A \times C \mid (\exists b \in B)((a, b) \in R \wedge (b, c) \in S) \}.$$

The identity $1_A : A \longrightarrow A$ is $\{ (a, a) \mid a \in A \}$.

Similarly, the category **Part** of sets and partial functions (i.e. relations such that $(a, b) \in R, (a, b') \in R \implies b = b'$).

- (j) Let K be a field. The category **Mat** $_K$ has natural numbers as objects, and morphisms $n \longrightarrow p$ are $(p \times n)$ matrices with entries from K . Composition is matrix multiplication.

1.4 Definition (Functor). Let \mathcal{C}, \mathcal{D} be **categories**. A **functor** $F : \mathcal{C} \longrightarrow \mathcal{D}$ consists of

- (a) a mapping $A \longmapsto FA$ from $\text{ob } \mathcal{C}$ to $\text{ob } \mathcal{D}$
- (b) a mapping $f \longmapsto Ff$ from $\text{mor } \mathcal{C}$ to $\text{mor } \mathcal{D}$

such that $\text{dom}(Ff) = F(\text{dom } f)$, $\text{cod}(Ff) = F(\text{cod } f)$, $1_{FA} = F(1_A)$ and $(Ff)(Fg) = F(fg)$ whenever fg is defined.

Lecture 2 1.3 Examples (Continued).

- (k) We write **Cat** for the category whose objects are all **small categories**, and whose morphisms are **functors** between them.

1.5 Examples.

- (a) We have **forgetful functors** $\mathbf{Gp} \xrightarrow{U} \mathbf{Set}, \mathbf{Rng} \rightarrow \mathbf{Set}, \mathbf{Top} \rightarrow \mathbf{Set}, \mathbf{Rng} \rightarrow \mathbf{AbGp}$ (forgetting \times), $\mathbf{Rng} \rightarrow \mathbf{Mon}$ (forgetting $+$).
- (b) Given a set A , the free group FA has the property: given any group G and any function $A \xrightarrow{f} UG$, there's a unique homomorphism $FA \xrightarrow{f} G$ extending f . F is a functor $\mathbf{Set} \rightarrow \mathbf{Gp}$: given $A \xrightarrow{f} B$, we define Ff to be the unique homomorphism extending $A \xrightarrow{f} B \hookrightarrow UFB$.
 Functoriality follows from uniqueness: given $B \xrightarrow{g} C$, $F(gf)$ and $(Fg)(Ff)$ are both homoms extending $A \xrightarrow{f} B \xrightarrow{g} C \hookrightarrow UFC$.
- (c) Given a set A , we write PA for the set of all subsets of A . We can make P into a functor $\mathbf{Set} \rightarrow \mathbf{Set}$: given $A \xrightarrow{f} B$, we define $Pf(A') = \{f(a) \mid a \in A'\}$ for $A' \subseteq A$. But we also have a functor $P^* : \mathbf{Set} \rightarrow \mathbf{Set}^{op}$ defined on objects by P , but $P^*f(B') = \{a \in A \mid f(a) \in B'\}$ for $B' \subseteq B$.
 By a **contravariant** functor $\mathcal{C} \rightarrow \mathcal{D}$, we mean a **functor** $\mathcal{C} \rightarrow \mathcal{D}^{op}$ (or $\mathcal{C}^{op} \rightarrow \mathcal{D}$). (A **covariant** functor is one that doesn't reverse arrows).
- (d) Let K be a field. We have a functor $* : \mathbf{Mod}_K \rightarrow \mathbf{Mod}_K^{op}$ defined by $V^* = \{\text{linear maps } V \rightarrow K\}$ and if $V \xrightarrow{f} W$, $f^*(\theta : W \rightarrow K) = \theta f$.
- (e) We have a functor $op : \mathbf{Cat} \rightarrow \mathbf{Cat}$ which is the 'identity' on morphisms. (Note that this is **covariant**).
- (f) A functor between monoids is a monoid homomorphism.
- (g) A functor between posets is an order-preserving map.
- (h) Let G be a group. A functor $F : G \rightarrow \mathbf{Set}$ consists of a set $A = F*$ together with an action of G on A , i.e. a permutation representation of G (where we use $*$ to refer to the unique object of the group). Similarly a functor $G \rightarrow \mathbf{Mod}_K$ is a K -linear representation of G .
- (i) The construction of a fundamental group $\pi_1(X, x)$ of a space X with basepoint x is a functor $\mathbf{Top}_* \rightarrow \mathbf{Gp}$ where \mathbf{Top}_* is the set of spaces with a chosen basepoint. Similarly, the fundamental groupoid is a functor $\mathbf{Top} \rightarrow \mathbf{Gpd}$ where \mathbf{Gpd} is the category of groupoids and functors between them.

1.6 Definition (Natural transformation). Let \mathcal{C}, \mathcal{D} be **categories** and $F, G : \mathcal{C} \rightrightarrows \mathcal{D}$ two **functors**. A **natural transformation** $\alpha : F \rightarrow G$ consists of an assignment $A \mapsto \alpha_A$ from $\text{ob } \mathcal{C}$ to $\text{mor } \mathcal{D}$, such that $\text{dom } \alpha_A = FA$ and $\text{cod } \alpha_A = GA$ for all A , and for all $A \xrightarrow{f} B$ in \mathcal{C} the square

$$\begin{array}{ccc} FA & \xrightarrow{Ff} & FB \\ \downarrow \alpha_A & & \downarrow \alpha_B \\ GA & \xrightarrow{Gf} & GB \end{array}$$

commutes (i.e. $\alpha_B(Ff) = (Gf)\alpha_A$).

1.3 Examples (Continued).

- (l) Given categories \mathcal{C}, \mathcal{D} , we write $[\mathcal{C}, \mathcal{D}]$ for the category whose objects are functors $\mathcal{C} \rightarrow \mathcal{D}$, and whose morphisms are natural transformations.

1.7 Examples.

- (a) Let K be a field, V a vector space over K . There is a linear map $\alpha_V : V \rightarrow V^{**}$ given by

$$\alpha_V(v)(\theta) = \theta(v)$$

for $\theta \in V^*$. This is the V -component of a natural transformation

$$1_{\mathbf{Mod}_K} \rightarrow ** : \mathbf{Mod}_K \rightarrow \mathbf{Mod}_K.$$

Lecture 3

- (b) For any set A , we have a mapping $\sigma_A : A \rightarrow PA$ sending a to $\{a\}$. If $f : A \rightarrow B$, then $Pf\{a\} = \{f(a)\}$, so σ is a natural transformation $1_{\mathbf{Set}} \rightarrow P$.
- (c) Let $F : \mathbf{Set} \rightarrow \mathbf{Gp}$ be the free group functor (Examples 1.5(b)) and $U : \mathbf{Gp} \rightarrow \mathbf{Set}$ the forgetful functor. The inclusions $A \rightarrow UFA$ form a natural transformation $1_{\mathbf{Set}} \rightarrow UF$.
- (d) Let G, H be groups and $f, g : G \rightarrow H$ two homomorphisms. A natural transformation $\alpha : f \rightarrow g$ corresponds to an element $h = \alpha_*$ of H such that $h.f(x) = g(x).h$ for all $x \in G$, or equivalently $f(x) = h^{-1}g(x)h$, i.e. f and g are conjugate group homomorphisms.
- (e) Let A, B be two G -sets regarded as functors $G \rightarrow \mathbf{Set}$. A natural transformation $A \rightarrow B$ is a function f satisfying $f(g.a) = g.f(a)$ for all $a \in A$, i.e. a G -equivariant map.

1.8 Lemma. Let $F, G : \mathcal{C} \rightarrow \mathcal{D}$ be two functors, and $\alpha : F \rightarrow G$ a natural transformation. Then α is an isomorphism in $[\mathcal{C}, \mathcal{D}]$ iff each α_A is an isomorphism in \mathcal{D} .

Proof.

\Rightarrow trivial

\Leftarrow Suppose each α_A has an inverse β_A . Given $f : A \rightarrow B$ in \mathcal{C} , we need to show that

$$\begin{array}{ccc} GA & \xrightarrow{Gf} & GB \\ \downarrow \beta_A & & \downarrow \beta_B \\ FA & \xrightarrow{Ff} & FB \end{array}$$

commutes.

But

$$\begin{aligned} (Ff)\beta_A &= \beta_B \alpha_B (Ff)\beta_A \\ &= \beta_B (Gf) \alpha_A \beta_A \\ &= \beta_B (Gf). \end{aligned}$$

□

1.9 Definition (Equivalent category). Let \mathcal{C}, \mathcal{D} be categories. By an **equivalence** between \mathcal{C} and \mathcal{D} , we mean a pair of **functors** $F : \mathcal{C} \rightarrow \mathcal{D}, G : \mathcal{D} \rightarrow \mathcal{C}$ together with natural isomorphisms $\alpha : 1_{\mathcal{C}} \rightarrow GF$ and $\beta : FG \rightarrow 1_{\mathcal{D}}$. We write $\mathcal{C} \simeq \mathcal{D}$ if \mathcal{C} and \mathcal{D} are equivalent.

We say a property P of **categories** is a **categorical property** if whenever \mathcal{C} has P and $\mathcal{C} \simeq \mathcal{D}$, then \mathcal{D} has P .

For instance, being a groupoid or a preorder are categorical properties, but being a group or a partial order are not.

1.10 Examples.

- (a) The category **Part** is equivalent to the category **Set**_{*} of pointed sets (and basepoint-preserving functions). We define $F : \mathbf{Set}_* \rightarrow \mathbf{Part}$ by $F(A, a) = A \setminus \{a\}$ and if $f : (A, a) \rightarrow (B, b)$

$$Ff(x) = \begin{cases} f(x) & \text{if } f(x) \neq b \\ \text{undefined} & \text{otherwise} \end{cases}$$

and $G : \mathbf{Part} \rightarrow \mathbf{Set}_*$ by $G(A) = A^+ = A \cup \{A\}$ and if $f : A \rightarrow B$ is a partial function, we define $Gf : A^+ \rightarrow B^+$ by

$$Gf = \begin{cases} f(x) & \text{if } x \in A \text{ and } f(x) \text{ defined} \\ B & \text{otherwise} \end{cases}$$

The composite FG is the identity on **Part**, but GF is not the identity, however there's an isomorphism

$$(A, a) \rightarrow ((A \setminus \{a\})^+, A \setminus \{a\})$$

sending a to $A \setminus \{a\}$ and everything else to itself and this is natural.

Note that there can be no isomorphism $\mathbf{Set}_* \rightarrow \mathbf{Part}$ since **Part** has a 1-element isomorphism class $\{\emptyset\}$ and **Set**_{*} doesn't.

- (b) The category **FdMod** _{K} of finite-dimensional vector spaces over K is equivalent to **FdMod** _{K} ^{op}: the functors in both directions are $(-)^*$ and both isomorphisms are the natural transformations of [Examples 1.7\(a\)](#).
- (c) **FdMod** _{K} is also equivalent to **Mat** _{K} : We define $F : \mathbf{Mat}_K \rightarrow \mathbf{FdMod}_K$ by $F(n) = K^n$, and $F(A)$ is the linear map represented by A with respect to the standard bases of K^n and K^p .

To define $G : \mathbf{FdMod}_K \rightarrow \mathbf{Mat}_K$, choose a basis for each finite dimensional vector space, and define $G(V) = \dim V$, $G(V \xrightarrow{f} W)$ as the matrix representing f with respect to the chosen bases. GF is the identity, provided we choose the standard bases for the spaces K^n ; $FG \neq 1$, but the chosen basis gives isomorphisms $FG(V) = K^{\dim V} \rightarrow V$ for each V , which form a natural isomorphism.

Lecture 4 **1.11 Definition** (Faithful, full, essentially surjective). Let $\mathcal{C} \xrightarrow{F} \mathcal{D}$ be a **functor**.

- (a) We say F is **faithful** if, given $f, f' \in \text{mor } \mathcal{C}$ with $\text{dom } f = \text{dom } f'$, $\text{cod } f = \text{cod } f'$ and $Ff = Ff'$ then $f = f'$.
- (b) We say F is **full** if, given $FA \xrightarrow{g} FB$ in \mathcal{D} , there exists $A \xrightarrow{f} B$ in \mathcal{C} with $Ff = g$.

- (c) We say F is **essentially surjective** if, for every $B \in \text{ob } \mathcal{D}$, there exists $A \in \text{ob } \mathcal{C}$ and an isomorphism $FA \rightarrow B$ in \mathcal{D} .

We say a subcategory $\mathcal{C}' \subseteq \mathcal{C}$ is **full** if the inclusion $\mathcal{C}' \rightarrow \mathcal{C}$ is a full functor.

Example. **Gp** is a **full subcategory** of **Mon**, but **Mon** is not a full subcategory of the category **Sgp** of semigroups.

1.12 Lemma. Assuming the axiom of choice, a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is part of an **equivalence** $\mathcal{C} \simeq \mathcal{D}$ iff it is **full**, **faithful** and **essentially surjective**.

Proof.

\Rightarrow Given G, α, β as in [Definition 1.9](#), for each $B \in \text{ob } \mathcal{D}$, β_B is an isomorphism $FGB \rightarrow B$, so F is essentially surjective.

Given $A \xrightarrow{f} B$ in \mathcal{C} , we can recover f from Ff as the composite

$$A \xrightarrow{\alpha_A} GFA \xrightarrow{GFf} GFB \xrightarrow{\alpha_B^{-1}} B.$$

Hence if $A \xrightarrow{f'} B$ satisfies $Ff = Ff'$, then $f = f'$.

Given $FA \xrightarrow{g} FB$, define f to be the composite

$$A \xrightarrow{\alpha_A} GFA \xrightarrow{Gg} GFB \xrightarrow{\alpha_B^{-1}} B$$

Then $GFf = \alpha_B f \alpha_A^{-1} = Gg$, and G is faithful for the same reason as F , so $Ff = g$.

\Leftarrow For each $B \in \text{ob } \mathcal{D}$, choose $GB \in \text{ob } \mathcal{C}$ and an isomorphism $\beta_B : FGB \rightarrow B$ in \mathcal{D} . Given

$$B \xrightarrow{g} B'$$

define $Gg : GB \rightarrow GB'$ to be the unique morphism whose image under F is

$$FGB \xrightarrow{\beta_B} B \xrightarrow{g} B' \xrightarrow{\beta_{B'}^{-1}} FGB'$$

Uniqueness implies functoriality: given

$$B' \xrightarrow{g'} B''$$

then note $(Gg')(Gg)$ and $G(g'g)$ have the same image under F , so they're equal.

By construction, β is a natural transformation $FG \rightarrow 1_{\mathcal{D}}$.

Given $A \in \text{ob } \mathcal{C}$, define $\alpha_A : A \rightarrow GFA$ to be the unique morphism whose image under F is

$$FA \xrightarrow{\beta_{FA}^{-1}} FGFA$$

α_A is an isomorphism, since β_{FA} also has a unique pre-image under F .

Also α is a natural transformation, since any naturality square for α is mapped by F to a commutative square, and F is faithful. \square

1.13 Definition (Skeleton). By a **skeleton** of a **category** \mathcal{C} , we mean a **full subcategory** \mathcal{C}_0 containing one object from each isomorphism class. We say \mathcal{C} is **skeletal** if it's a skeleton of itself.

Example. \mathbf{Mat}_K is **skeletal**, and the image of $F : \mathbf{Mat}_K \rightarrow \mathbf{FdMod}_K$ of [Examples 1.10\(c\)](#) is a **skeleton** of \mathbf{FdMod}_K .

Warning. Almost any assertion about **skeletons** is equivalent to the axiom of choice. See question 2 on example sheet 1.

1.14 Definition (Monomorphism, epimorphism). Let $A \xrightarrow{f} B$ be a **morphism** in \mathcal{C}

- (a) We say f is a **monomorphism** (or f is **monic**) if, given any pair $C \rightrightarrows A$, $fg = fh$ implies $g = h$
- (b) We say f is an **epimorphism** (or **epic**) if it's a monomorphism in \mathcal{C}^{op} i.e. if $gf = hf$ implies $g = h$.

We denote **monomorphisms** by $A \rightarrowtail B$ and **epimorphisms** by $A \twoheadrightarrow B$

Any isomorphism is **monic** and **epic**: more generally if f has a left inverse (i.e. $\exists g$ such that gf is an identity) then it's monic. We call such monomorphisms **split**.

We say \mathcal{C} is a **balanced category** if any morphism which is both **monic** and **epic** is an isomorphism.

1.15 Examples.

- (a) In **Set**, **mono** \iff injective (\implies easy; for \impliedby take $C = 1 = \{*\}$) and **epi** \iff surjective (\implies easy; for \impliedby use two morphisms $B \rightarrow 2 = \{0, 1\}$). So **Set** is **balanced**.
- (b) In **Gp** **mono** \iff injective (for \impliedby use homoms $\mathbb{Z} \rightarrow A$) and **epi** \iff surjective (\impliedby uses free products with amalgamation). So **Gp** is balanced.
- (c) In **Rng**, **mono** \iff injective (proof much as for **Gp**) but the inclusion $\mathbb{Z} \rightarrow \mathbb{Q}$ is an epimorphism, since if $\mathbb{Q} \rightrightarrows R$ agree on all integers, they agree everywhere. So **Rng** isn't balanced.
- (d) In **Top**, **mono** \iff injective and **epi** \iff surjective (proofs as in **Set**). But **Top** isn't balanced since a continuous bijection needn't have a continuous inverse.

2 The Yoneda Lemma

Lecture 5 **2.1 Definition** (Locally small). We say a **category** \mathcal{C} is **locally small** if, for any two objects A, B , the morphisms $A \rightarrow B$ in \mathcal{C} form a set $\mathcal{C}(A, B)$.

If we fix A and let B vary, the assignment $B \mapsto \mathcal{C}(A, B)$ becomes a **functor** $\mathcal{C}(A, -) : \mathcal{C} \rightarrow \mathbf{Set}$: given $B \xrightarrow{f} C$, $\mathcal{C}(A, f)$ is the mapping $g \mapsto fg$. Similarly, $A \mapsto \mathcal{C}(A, B)$ defines a functor $\mathcal{C}(-, B) : \mathcal{C}^{op} \rightarrow \mathbf{Set}$.

2.2 Lemma. Let \mathcal{C} be a **locally small category**, $A \in \text{ob } \mathcal{C}$ and $F : \mathcal{C} \rightarrow \mathbf{Set}$ a **functor**.

- (i) Then natural transformations $\mathcal{C}(A, -) \rightarrow F$ is a bijection with elements of FA .
- (ii) Moreover, this bijection is natural in both A and F .

Proof of Lemma 2.2(i). Given $\alpha : \mathcal{C}(A, -) \rightarrow F$, we define

$$\Phi(\alpha) = \alpha_A(1_A) \in FA.$$

Given $x \in FA$, we define $\Psi(x) : \mathcal{C}(A, -) \rightarrow F$ by

$$\Psi(x)_B(A \xrightarrow{f} B) = Ff(x) \in FB.$$

$\Psi(x)$ is natural: given $g : B \rightarrow C$, we have

$$\begin{aligned} \Psi(x)_C \mathcal{C}(A, g)(f) &= \Psi(x)_C(gf) = F(gf)(x) \\ (Fg)\Psi(x)_B(f) &= (Fg)(Ff)(x) = F(gf)(x). \end{aligned}$$

We also verify Ψ and Φ are inverse:

$$\Phi\Psi(x) = \Psi(x)_A(1_A) = F(1_A)(x) = x.$$

Given α ,

$$\begin{aligned} \Psi\Phi(\alpha)_B(f) &= \Phi(\alpha_A(1_A))_B(f) = Ff(\alpha_A(1_A)) \\ &= \alpha_B \mathcal{C}(A, f)(1_A) = \alpha_B(f) \end{aligned}$$

so $\Psi\Phi(\alpha) = \alpha$. □

2.3 Corollary. The assignment $A \mapsto \mathcal{C}(A, -)$ defines a **full** and **faithful functor** $\mathcal{C}^{op} \rightarrow [\mathcal{C}, \mathbf{Set}]$.

Proof. Put $F = \mathcal{C}(B, -)$ in **Lemma 2.2(i)**: we get a bijection between $\mathcal{C}(B, A)$ and morphisms $\mathcal{C}(A, -) \rightarrow \mathcal{C}(B, -)$ in $[\mathcal{C}, \mathbf{Set}]$. We need to verify this is functorial: but it sends $f : B \rightarrow A$ to the natural transformation $g \mapsto gf$. So functoriality follows from associativity. □

We call this **functor** (or the functor $\mathcal{C} \rightarrow [\mathcal{C}^{op}, \mathbf{Set}]$) sending A to $\mathcal{C}(-, A)$ the **Yoneda embedding** of \mathcal{C} , and denote it by Y .

Proof of Lemma 2.2(ii). Suppose for the moment that \mathcal{C} is **small**, so that $[\mathcal{C}, \mathbf{Set}]$ is **locally small**. Then we have two functors $\mathcal{C} \times [\mathcal{C}, \mathbf{Set}] \rightarrow \mathbf{Set}$: One sends (A, F) to FA , and the other is the composite

$$\mathcal{C} \times [\mathcal{C}, \mathbf{Set}] \xrightarrow{Y \times 1} [\mathcal{C}, \mathbf{Set}]^{op} \times [\mathcal{C}, \mathbf{Set}] \xrightarrow{[\mathcal{C}, \mathbf{Set}](-, -)} \mathbf{Set}$$

Lemma 2.2(ii) says that these are naturally isomorphic.

We can translate this into an elementary statement, making sense even when \mathcal{C} isn't small, given $A \xrightarrow{f} B$ and $F \xrightarrow{\alpha} G$, the two ways of producing an element of GB from a natural transformation $\beta : \mathcal{C}(A, -) \rightarrow F$ give the same result, namely

$$\alpha_B(Ff)\beta_A(1_A) = (GF)\alpha_A\beta_A(1_A)$$

which is equal to $\alpha_B\beta_B(f)$. □

2.4 Definition. We say a functor $F : \mathcal{C} \rightarrow \mathbf{Set}$ is **representable** if it's **isomorphic** to $\mathcal{C}(A, -)$ for some A . By **representation** of F , we mean a pair (A, x) where $x \in FA$ is such that $\Phi(x)$ is an **isomorphism**. We also call x a **universal element** of F .

2.5 Corollary. If (A, x) and (B, y) are both **representations** of F , then there's a unique **isomorphism** $f : A \rightarrow B$ such that $(Ff)(x) = y$.

Proof. Consider the composite

$$\mathcal{C}(B, -) \xrightarrow{\Psi(y)} F \xrightarrow{\Psi(x)^{-1}} \mathcal{C}(A, -)$$

By Corollary 2.3, this is of the form $Y(f)$ for a unique **isomorphism** $f : A \rightarrow B$ and the diagram

$$\begin{array}{ccc} \mathcal{C}(B, -) & \xrightarrow{Y(f)} & \mathcal{C}(A, -) \\ & \searrow \Psi(y) & \swarrow \Psi(x) \\ & F & \end{array}$$

commutes iff $(Ff)x = y$. □

2.6 Examples.

- (a) The **forgetful functor** $\mathbf{Gp} \rightarrow \mathbf{Set}$ is **representable** by $(\mathbb{Z}, 1)$. Similarly, the forgetful functor $\mathbf{Rng} \rightarrow \mathbf{Set}$ is representable by $(\mathbb{Z}[x], x)$ and the forgetful functor $\mathbf{Top} \rightarrow \mathbf{Set}$ is representable by $(\{\ast\}, \ast)$.
- (b) The functor $P^* : \mathbf{Set}^{op} \rightarrow \mathbf{Set}$ (see Examples 1.5(c)) is representable by $(\{0, 1\}, \{1\})$: this is the bijection between subsets and characteristic functions.
- (c) Let G be a group. The unique (up to **isomorphism**) representable functor $G(\ast, -) : G \rightarrow \mathbf{Set}$ is the *Cayley representation* of G , i.e. the set UG with G acting by left multiplication.

- (d) Let A, B be two objects of a **locally small category** \mathcal{C} . We have a functor $\mathcal{C}^{op} \rightarrow \mathbf{Set}$ sending C to $\mathcal{C}(C, A) \times \mathcal{C}(C, B)$. A representation of this, if it exists, is called a (categorical) **product** of A and B , and denoted

$$(A \times B, (A \times B \xrightarrow{\pi_1} A), (A \times B \xrightarrow{\pi_2} B)).$$

This pair has the property that, for any pair $(C \xrightarrow{f} A, C \xrightarrow{g} B)$ there's a unique $C \xrightarrow{h} A \times B$ with $\pi_1 h = f$ and $\pi_2 h = g$.

Products exist in many categories of interest: in **Set**, **Gp**, **Rng**, **Top** they are 'just' Cartesian products, in posets they are binary meets.

Dually we have the notion of **coproduct** $(A + B, A \xrightarrow{\nu_1} A + B, B \xrightarrow{\nu_2} A + B)$.

These also exist in many categories of interest.

Index

- balanced, 8
- categorical property, 6
- category, 2
 - axioms, 2
 - balanced, 8
 - equivalence, 6
 - small, 2
- codomain, 2
- composition, 2
- domain, 2
- duality, 3
- epimorphism, 8
- equivalent, 6
- functor, 3
 - contravariant, 4
 - essentially surjective, 7
 - faithful, 6
 - forgetful, 4
 - full, 6
- identity, 2
- isomorphism, 3
- locally small, 9
- monomorphism, 8
 - split, 8
- morphism, 2
 - epi-, 8
 - mono-, 8
- natural transformations, 4
- object, 2
- representable, 10
- representation, 10
- split monomorphism, 8