

Generalising induction, and coinduction

Bhavik Mehta

Part III Seminar

Friday 30 November

Generalising induction, and coinduction

Generalising induction, and coinduction

- Recursive definitions and principle of induction away from \mathbb{N}

Generalising induction, and coinduction

- Recursive definitions and principle of induction away from \mathbb{N}
- Dualise: what is corecursive data?

Generalising induction, and coinduction

- Recursive definitions and principle of induction away from \mathbb{N}
- Dualise: what is corecursive data?
- Universal algebra, model theory, automata, real analysis, theoretical computer science

Algebras of an endofunctor

Take an endofunctor $F : \mathcal{C} \rightarrow \mathcal{C}$

Definition (F -algebra)

An F -algebra is a pair $(A, \alpha : FA \rightarrow A)$ with $A \in \text{ob } \mathcal{C}$

Algebras of an endofunctor

Take an endofunctor $F : \mathcal{C} \rightarrow \mathcal{C}$

Definition (F -algebra)

An F -algebra is a pair $(A, \alpha : FA \rightarrow A)$ with $A \in \text{ob } \mathcal{C}$

Definition (Algebra homomorphism)

A homomorphism of F -algebras $(A, \alpha) \rightarrow (B, \beta)$ is a morphism $f : A \rightarrow B$ with

$$\begin{array}{ccc} FA & \xrightarrow{Ff} & FB \\ \downarrow \alpha & & \downarrow \beta \\ A & \xrightarrow{f} & B \end{array}$$

Alg F is the category of F -algebras

Coalgebras of an endofunctor

Take an endofunctor $F : \mathcal{C} \rightarrow \mathcal{C}$

Definition (F -coalgebra)

An F -coalgebra is a pair $(A, \alpha : A \rightarrow FA)$ with $A \in \text{ob } \mathcal{C}$

Definition (Coalgebra homomorphism)


A homomorphism of F -coalgebras $(A, \alpha) \rightarrow (B, \beta)$ is a morphism $f : A \rightarrow B$ with

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow \alpha & & \downarrow \beta \\ FA & \xrightarrow{Ff} & FB \end{array}$$

$\mathbf{Coalg} F$ is the category of F -coalgebras


“Monoids”

$$FX := 1 + X \times X$$



“Monoids”

$$FX := 1 + X \times X$$



- An F -algebra gives an *interpretation*, not necessarily a model

“Monoids”

$$FX := 1 + X \times X$$

$$\begin{array}{ccccc}
 1 & \longrightarrow & 1 + A \times A & \longleftarrow & A \times A \\
 & \searrow e & \downarrow e+m & \swarrow m & \\
 & & A & &
 \end{array}$$

- An F -algebra gives an *interpretation*, not necessarily a model

$$f(e_A) = e_B$$

$$f(m_A(x, y)) = m_B(f(x), f(y))$$

$$\begin{array}{ccc}
 1 + A \times A & \longrightarrow & 1 + B \times B \\
 e_A + m_A \downarrow & & \downarrow e_B + m_B \\
 A & \xrightarrow{f} & B
 \end{array}$$

“Monoids”

$$FX := 1 + X \times X$$

$$\begin{array}{ccccc}
 1 & \longrightarrow & 1 + A \times A & \longleftarrow & A \times A \\
 & \searrow e & \downarrow e+m & \swarrow m & \\
 & & A & &
 \end{array}$$

$$\begin{array}{ccc}
 1 + A \times A & \longrightarrow & 1 + B \times B \\
 e_A + m_A \downarrow & & \downarrow e_B + m_B \\
 A & \xrightarrow{f} & B
 \end{array}$$

- An F -algebra gives an *interpretation*, not necessarily a model


$$f(e_A) = e_B$$

$$f(m_A(x, y)) = m_B(f(x), f(y))$$

- Alg** F has a full subcategory isomorphic to **Mon**

Trees


- The set T of finite binary trees gives an F -algebra



$1 \rightarrow T$ gives empty tree, $T \times T \rightarrow T$ combines trees

Trees

- The set T of finite binary trees gives an F -algebra



$1 \rightarrow T$ gives empty tree, $T \times T \rightarrow T$ combines trees

- All binary trees also works

More algebra examples

- $FX := 1 + X + X \times X$
 $FA \rightarrow A$ decomposes into $1 \rightarrow A$, $A \rightarrow A$, $A \times A \rightarrow A$

More algebra examples

- $FX := 1 + X + X \times X$
 $FA \rightarrow A$ decomposes into $1 \rightarrow A, A \rightarrow A, A \times A \rightarrow A$
- $FX := X \times X$ has semigroups, $FX := 2 + X + 2 \times X \times X$ has rings

More algebra examples


- $FX := 1 + X + X \times X$
 $FA \rightarrow A$ decomposes into $1 \rightarrow A$, $A \rightarrow A$, $A \times A \rightarrow A$
- $FX := X \times X$ has semigroups, $FX := 2 + X + 2 \times X \times X$ has rings
- $FX := \mathcal{P}X$, then $\mathcal{P}A \rightarrow A$ could be a point in A , or $\mathcal{P}\mathcal{P}B \rightarrow \mathcal{P}B$, or many things...

More algebra examples

- $FX := 1 + X + X \times X$
 $FA \rightarrow A$ decomposes into $1 \rightarrow A$, $A \rightarrow A$, $A \times A \rightarrow A$
- $FX := X \times X$ has semigroups, $FX := 2 + X + 2 \times X \times X$ has rings
- $FX := \mathcal{P}X$, then $\mathcal{P}A \rightarrow A$ could be a point in A , or $\mathcal{P}\mathcal{P}B \rightarrow \mathcal{P}B$, or many things...
- If $F0 = 0$ then 0 has an F -algebra structure


Coalgebra examples

- $FX = 1 + X$, then $f : A \rightarrow 1 + A$




Coalgebra examples

- $FX = 1 + X$, then $f : A \rightarrow 1 + A$



Coalgebra examples


- $FX = 1 + X$, then $f : A \rightarrow 1 + A$



- $FX := B \times X$, then $A \rightarrow B \times A$ is a deterministic automaton with output in B (but no fixed input state)

Coalgebra examples


- $FX = 1 + X$, then $f : A \rightarrow 1 + A$




- $FX := B \times X$, then $A \rightarrow B \times A$ is a deterministic automaton with output in B (but no fixed input state)
- $FX := \mathcal{P}X$ models non-deterministic automata

Coalgebra examples

- $FX = 1 + X$, then $f : A \rightarrow 1 + A$



- $FX := B \times X$, then $A \rightarrow B \times A$ is a deterministic automaton with output in B (but no fixed input state)
- $FX := \mathcal{P}X$ models non-deterministic automata
- $FX := 1 + X \times X$



and trees

Initial algebras

Definition

An initial F -algebra is an initial object in the category of F -algebras

For any $FA \rightarrow A$, there is a unique morphism $i : I \rightarrow A$, with

$$\begin{array}{ccc} FI & \xrightarrow{Fi} & FA \\ \downarrow & & \downarrow \\ I & \xrightarrow{i} & A \end{array}$$

Initial algebras

Definition

An initial F -algebra is an initial object in the category of F -algebras

For any $FA \rightarrow A$, there is a unique morphism $i : I \rightarrow A$, with

$$\begin{array}{ccc} FI & \xrightarrow{Fi} & FA \\ \downarrow & & \downarrow \\ I & \xrightarrow{i} & A \end{array}$$

Definition

A terminal (or final) F -coalgebra is a terminal object in the category of F -coalgebras

Induction on \mathbb{N}

Take $FX = 1 + X$ on **Set**, then \mathbb{N} forms an F -algebra

$$\begin{array}{ccccc}
 1 & \longrightarrow & 1 + \mathbb{N} & \longleftarrow & \mathbb{N} \\
 & \searrow 0 & \downarrow 0+s & \swarrow s & \\
 & & \mathbb{N} & &
 \end{array}$$

Induction on \mathbb{N}

Take $FX = 1 + X$ on **Set**, then \mathbb{N} forms an F -algebra

$$\begin{array}{ccccc}
 1 & \longrightarrow & 1 + \mathbb{N} & \longleftarrow & \mathbb{N} \\
 & \searrow 0 & \downarrow 0+s & \swarrow s & \\
 & & \mathbb{N} & &
 \end{array}$$

$$\begin{array}{ccc}
 1 + \mathbb{N} & \xrightarrow{1+\varphi} & 1 + A \\
 \downarrow 0+s & & \downarrow x_0+f \\
 \mathbb{N} & \xrightarrow{\varphi} & A
 \end{array}
 \qquad
 \begin{array}{l}
 \varphi(0) = x_0 \\
 \varphi(n+1) = f(\varphi(n))
 \end{array}$$

Recall: Subobjects of an initial object are isomorphic to it

Streams

$FX = B \times X$, fixed set B ; B^∞ is infinite sequences (streams) of B ;
 $B^\infty \rightarrow B \times B^\infty$ is head and tail, $(x_n)_{n=0}^\infty \mapsto (x_0, (x_n)_{n=1}^\infty)$

Streams

$FX = B \times X$, fixed set B ; B^∞ is infinite sequences (streams) of B ;
 $B^\infty \rightarrow B \times B^\infty$ is head and tail, $(x_n)_{n=0}^\infty \mapsto (x_0, (x_n)_{n=1}^\infty)$

$$a_0 \mapsto (b_1, a_2)$$

$$a_1 \mapsto (b_3, a_0)$$

$$a_2 \mapsto (b_4, a_1)$$

$$\begin{array}{ccc} A & \xrightarrow{\varphi} & B^\infty \\ \downarrow f & & \downarrow \\ B \times A & \xrightarrow{1_B \times \varphi} & B \times B^\infty \end{array}$$

Streams

$FX = B \times X$, fixed set B ; B^∞ is infinite sequences (streams) of B ;
 $B^\infty \rightarrow B \times B^\infty$ is head and tail, $(x_n)_{n=0}^\infty \mapsto (x_0, (x_n)_{n=1}^\infty)$

$$\begin{array}{ccc}
 A & \xrightarrow{\varphi} & B^\infty \\
 \downarrow f & & \downarrow \\
 B \times A & \xrightarrow{1_B \times \varphi} & B \times B^\infty
 \end{array}$$

$$a_0 \mapsto (b_1, a_2)$$

$$a_1 \mapsto (b_3, a_0)$$

$$a_2 \mapsto (b_4, a_1)$$

$$\begin{aligned}
 \varphi(a_0) &= \langle b_1, \varphi(a_2) \rangle \\
 &= \langle b_1, \langle b_4, \varphi(a_1) \rangle \rangle \\
 &= \langle b_1, \langle b_4, \langle b_3, \varphi(a_0) \rangle \rangle \rangle \\
 &= (b_1, b_4, b_3, b_1, b_4, \dots)
 \end{aligned}$$

Streams

$FX = B \times X$, fixed set B ; B^∞ is infinite sequences (streams) of B ;
 $B^\infty \rightarrow B \times B^\infty$ is head and tail, $(x_n)_{n=0}^\infty \mapsto (x_0, (x_n)_{n=1}^\infty)$

$$\begin{array}{ccc}
 A & \xrightarrow{\varphi} & B^\infty \\
 \downarrow f & & \downarrow \\
 B \times A & \xrightarrow{1_B \times \varphi} & B \times B^\infty
 \end{array}$$

Initial algebra is boring

$$a_0 \mapsto (b_1, a_2)$$

$$a_1 \mapsto (b_3, a_0)$$

$$a_2 \mapsto (b_4, a_1)$$

$$\begin{aligned}
 \varphi(a_0) &= \langle b_1, \varphi(a_2) \rangle \\
 &= \langle b_1, \langle b_4, \varphi(a_1) \rangle \rangle \\
 &= \langle b_1, \langle b_4, \langle b_3, \varphi(a_0) \rangle \rangle \rangle \\
 &= (b_1, b_4, b_3, b_1, b_4, \dots)
 \end{aligned}$$

Powerset functors

\mathcal{P}_{fin} on **Set** has an initial algebra (V_ω, id) , **hereditarily finite sets**

$$\begin{array}{ccc}
 \mathcal{P}_{\text{fin}} V_\omega & \xrightarrow{\mathcal{P}_{\text{fin}} \varphi} & \mathcal{P}_{\text{fin}} A \\
 \downarrow \text{id} & & \downarrow f \\
 V_\omega & \xrightarrow{\varphi} & A
 \end{array}$$

Powerset functors

\mathcal{P}_{fin} on **Set** has an initial algebra (V_ω, id) , **hereditarily finite sets**

$$\begin{array}{ccc}
 \mathcal{P}_{\text{fin}} V_\omega & \xrightarrow{\mathcal{P}_{\text{fin}} \varphi} & \mathcal{P}_{\text{fin}} A \\
 \downarrow \text{id} & & \downarrow f \\
 V_\omega & \xrightarrow{\varphi} & A
 \end{array}$$

$$\varphi(x) = f(\{\varphi(s) : s \in x\})$$

Powerset functors

\mathcal{P}_{fin} on **Set** has an initial algebra (V_ω, id) , **hereditarily finite sets**

$$\begin{array}{ccc} \mathcal{P}_{\text{fin}} V_\omega & \xrightarrow{\mathcal{P}_{\text{fin}} \varphi} & \mathcal{P}_{\text{fin}} A \\ \downarrow \text{id} & & \downarrow f \\ V_\omega & \xrightarrow{\varphi} & A \end{array}$$

$$\varphi(x) = f(\{\varphi(s) : s \in x\})$$

\mathcal{P}_κ = subsets of cardinality $< \kappa$ has V_κ . What about \mathcal{P} itself?

A necessary condition

Lemma (Lambek)

If (A, α) is an initial F -algebra, then α is an isomorphism

A necessary condition

Lemma (Lambek)

If (A, α) is an initial F -algebra, then α is an isomorphism

Proof.

$$\begin{array}{ccc} FA & & \\ \downarrow \alpha & & \\ A & & FA \end{array}$$



A necessary condition

Lemma (Lambek)

If (A, α) is an initial F -algebra, then α is an isomorphism

Proof.

$$\begin{array}{ccc}
 FA & \xrightarrow{Fi} & FFA \\
 \downarrow \alpha & & \downarrow F\alpha \\
 A & \xrightarrow{i} & FA
 \end{array}$$



A necessary condition

Lemma (Lambek)

If (A, α) is an initial F -algebra, then α is an isomorphism

Proof.

$$\begin{array}{ccccc}
 FA & \xrightarrow{Fi} & FFA & \xrightarrow{F\alpha} & FA \\
 \downarrow \alpha & & \downarrow F\alpha & & \downarrow \alpha \\
 A & \xrightarrow{i} & FA & \xrightarrow{\alpha} & A
 \end{array}$$



A necessary condition

Lemma (Lambek)

If (A, α) is an initial F -algebra, then α is an isomorphism

Proof.

$$\begin{array}{ccccc}
 FA & \xrightarrow{Fi} & FFA & \xrightarrow{F\alpha} & FA \\
 \downarrow \alpha & & \downarrow F\alpha & & \downarrow \alpha \\
 A & \xrightarrow{i} & FA & \xrightarrow{\alpha} & A \\
 & \searrow & & \nearrow & \\
 & 1_A & & &
 \end{array}$$




A necessary condition

Lemma (Lambek)

If (A, α) is an initial F -algebra, then α is an isomorphism

Proof.



Fixed points of an endofunctor

- If (A, α) is initial, then $FA \cong A$
- Dually, if (A, α) is a terminal coalgebra, then $A \cong FA$.

Fixed points of an endofunctor

- If (A, α) is initial, then $FA \cong A$
- Dually, if (A, α) is a terminal coalgebra, then $A \cong FA$.

Definition (Fixed point)

A fixed point of F is an isomorphism $FA \rightarrow A$ (equivalently, $A \rightarrow FA$)

Fixed points of an endofunctor

- If (A, α) is initial, then $FA \cong A$
- Dually, if (A, α) is a terminal coalgebra, then $A \cong FA$.

Definition (Fixed point)

A fixed point of F is an isomorphism $FA \rightarrow A$ (equivalently, $A \rightarrow FA$)

- \mathcal{P} has no fixed points, in particular no initial algebra (Cantor)
- If 0 is a fixed point of F , it is an initial algebra

Least, greatest fixed points

- $\text{Fix}(F)$, a full subcategory of **Alg** F and of **Coalg** F

Least, greatest fixed points

- $\text{Fix}(F)$, a full subcategory of **Alg** F and of **Coalg** F
- Initial algebra is initial in this subcategory

Least, greatest fixed points

- $\text{Fix}(F)$, a full subcategory of **Alg** F and of **Coalg** F
- Initial algebra is initial in this subcategory
- Subobjects of an initial object are isomorphic to it

Least, greatest fixed points

- $\text{Fix}(F)$, a full subcategory of **Alg** F and of **Coalg** F
- Initial algebra is initial in this subcategory
- Subobjects of an initial object are isomorphic to it
- So initial algebra is *least* fixed point

Least, greatest fixed points

- $\text{Fix}(F)$, a full subcategory of **Alg** F and of **Coalg** F
- Initial algebra is initial in this subcategory
- Subobjects of an initial object are isomorphic to it
- So initial algebra is *least* fixed point
- Dually, terminal coalgebra is greatest fixed point

Trees



T is initial for $FX = 1 + X^2$

Trees



T is initial for $FX = 1 + X^2$

$$\begin{array}{ccc}
 1 + T^2 & \longrightarrow & 1 + \mathbb{N}^2 \\
 \downarrow & & \downarrow 1+\text{add} \\
 T & \xrightarrow{\varphi} & \mathbb{N}
 \end{array}$$

Trees



T is initial for $FX = 1 + X^2$

$$\begin{array}{ccc}
 1 + T^2 & \longrightarrow & 1 + \mathbb{N}^2 \\
 \downarrow \scriptstyle \text{dashed} & & \downarrow \scriptstyle 1+\text{add} \\
 T & \xrightarrow{\varphi} & \mathbb{N}
 \end{array}$$

$$\begin{aligned}
 \varphi \left(\begin{array}{c} \wedge \\ \bullet \quad \bullet \\ \bullet \quad \bullet \end{array} \right) &= \varphi(\bullet) + \varphi(\wedge) \\
 &= 1 + (\varphi(\bullet) + \varphi(\bullet)) \\
 &= 3.
 \end{aligned}$$

Trees, continued

$$\begin{array}{ccc}
 1 + T^2 & \longrightarrow & 1 + \mathbb{N}^2 \\
 \downarrow \scriptstyle \text{dashed} & & \downarrow 0+f \\
 T & \xrightarrow{\varphi} & \mathbb{N}
 \end{array}$$

$$f(m, n) = 1 + \max\{m, n\}$$

Trees, continued

$$\begin{array}{ccc}
 1 + T^2 & \longrightarrow & 1 + \mathbb{N}^2 \\
 \downarrow & & \downarrow 0+f \\
 T & \xrightarrow{\varphi} & \mathbb{N}
 \end{array}$$

$$f(m, n) = 1 + \max\{m, n\}$$

$$\begin{aligned}
 \varphi\left(\begin{array}{c} \diagup \diagdown \\ \bullet \quad \bullet \end{array}\right) &= 1 + \max\{\varphi(\bullet), \varphi(\bullet)\} \\
 &= 1 + (1 + \max\{\varphi(\bullet), \varphi(\bullet)\}) \\
 &= 2.
 \end{aligned}$$

Conaturals?

$FX = 1 + X$. Take $\overline{\mathbb{N}} = \mathbb{N} \cup \{\infty\}$, and


$$\alpha : \overline{\mathbb{N}} \longrightarrow 1 + \overline{\mathbb{N}}$$

$$0 \longmapsto *$$

$$n \longmapsto n - 1$$

$$\infty \longmapsto \infty$$

$$\begin{array}{ccc} A & \longrightarrow & \overline{\mathbb{N}} \\ \downarrow & & \downarrow \\ 1 + A & \longrightarrow & 1 + \overline{\mathbb{N}} \end{array}$$



Finite lists

$FX = 1 + B \times X$, B^* are the finite sequences of B (finite lists)

Finite lists

$FX = 1 + B \times X$, B^* are the finite sequences of B (finite lists)

$$\begin{array}{ccc}
 1 + \mathbb{Z} \times \mathbb{Z}^* & \longrightarrow & 1 + \mathbb{Z} \times \mathbb{Q} \\
 \varepsilon + \langle \cdot, \cdot \rangle \downarrow & & \downarrow f \\
 \mathbb{Z}^* & \xrightarrow{\varphi} & \mathbb{Q}
 \end{array}$$

$$f(*) = 1$$

$$f(n, q) = n \times q$$

Finite lists

$FX = 1 + B \times X$, B^* are the finite sequences of B (finite lists)

$$\begin{array}{ccc}
 1 + \mathbb{Z} \times \mathbb{Z}^* & \longrightarrow & 1 + \mathbb{Z} \times \mathbb{Q} \\
 \varepsilon + \langle \cdot, \cdot \rangle \downarrow & & \downarrow f \\
 \mathbb{Z}^* & \xrightarrow{\varphi} & \mathbb{Q}
 \end{array}$$

$$f(*) = 1$$

$$f(n, q) = n \times q$$

$$\varphi([3, -4, 2]) = f(3, \varphi([-4, 2]))$$

$$\varphi([-4, 2]) = f(-4, \varphi([2]))$$

$$\varphi([2]) = f(2, \varphi([]))$$

$$\varphi([]) = f(*) = 1$$

Finite lists

$FX = 1 + B \times X$, B^* are the finite sequences of B (finite lists)

$$\begin{array}{ccc}
 1 + \mathbb{Z} \times \mathbb{Z}^* & \longrightarrow & 1 + \mathbb{Z} \times \mathbb{Q} \\
 \varepsilon + \langle \cdot, \cdot \rangle \downarrow & & \downarrow f \\
 \mathbb{Z}^* & \xrightarrow{\varphi} & \mathbb{Q}
 \end{array}$$

$$f(*) = 1$$

$$f(n, q) = n \times q$$

$$\varphi([3, -4, 2]) = f(3, \varphi([-4, 2]))$$

$$\varphi([-4, 2]) = f(-4, \varphi([2]))$$

$$\varphi([2]) = f(2, \varphi([]))$$

$$\varphi([]) = f(*) = 1$$

$$\begin{aligned}
 \varphi([3, -4, 2]) &= 3 \times (-4 \times (2 \times 1)) \\
 &= -24
 \end{aligned}$$

More coalgebras

Terminal coalgebra of $FX = 1 + B \times X$ is (potentially infinite) lists, B^ω

$$\beta : B^\omega \rightarrow 1 + B \times B^\omega$$

$$\varepsilon \mapsto *$$

$$\langle b_0, \mathbf{b} \rangle \mapsto (b_0, \mathbf{b})$$

More coalgebras

Terminal coalgebra of $FX = 1 + B \times X$ is (potentially infinite) lists, B^ω

$$\beta : B^\omega \rightarrow 1 + B \times B^\omega$$

$$\varepsilon \mapsto *$$

$$\langle b_0, \mathbf{b} \rangle \mapsto (b_0, \mathbf{b})$$

$$f(0) = *$$

$$f(n) = (n, n-1)$$

$$\begin{array}{ccc}
 \mathbb{Z} & \xrightarrow{\varphi} & \mathbb{Z}^\omega \\
 \downarrow f & & \downarrow \beta \\
 1 + \mathbb{Z} \times \mathbb{Z} & \longrightarrow & 1 + \mathbb{Z} \times \mathbb{Z}^\omega
 \end{array}$$

More coalgebras

Terminal coalgebra of $FX = 1 + B \times X$ is (potentially infinite) lists, B^ω

$$\beta : B^\omega \rightarrow 1 + B \times B^\omega$$

$$\varepsilon \mapsto *$$

$$\langle b_0, \mathbf{b} \rangle \mapsto (b_0, \mathbf{b})$$

$$\begin{array}{ccc} \mathbb{Z} & \xrightarrow{\varphi} & \mathbb{Z}^\omega \\ \downarrow f & & \downarrow \beta \\ 1 + \mathbb{Z} \times \mathbb{Z} & \longrightarrow & 1 + \mathbb{Z} \times \mathbb{Z}^\omega \end{array}$$

$$f(0) = *$$

$$f(n) = (n, n-1)$$

$$\begin{aligned} \varphi(2) &= \langle 2, \varphi(1) \rangle \\ &= \langle 2, \langle 1, \varphi(0) \rangle \rangle \\ &= \langle 2, \langle 1, \varepsilon \rangle \rangle \\ &= [2, 1] \end{aligned}$$

$\varphi(-5)$ is infinite: $[-5, -6, -7, \dots]$

Our examples

Functor	Initial algebra	Terminal coalgebra
$1 + X$	naturals	conaturals
$1 + X^2$	finite binary trees	binary trees
$1 + B \times X$	finite lists	lists
$B \times X$	empty	streams
\mathcal{P}_{fin}	V_ω	finitely branching trees

Our examples

Functor	Initial algebra	Terminal coalgebra
$1 + X$	naturals	conaturals
$1 + X^2$	finite binary trees	binary trees
$1 + B \times X$	finite lists	lists
$B \times X$	empty	streams
\mathcal{P}_{fin}	V_ω	finitely branching trees

- Dyadic rationals in $[0, 1]$ as an initial algebra

Our examples

Functor	Initial algebra	Terminal coalgebra
$1 + X$	naturals	conaturals
$1 + X^2$	finite binary trees	binary trees
$1 + B \times X$	finite lists	lists
$B \times X$	empty	streams
\mathcal{P}_{fin}	V_ω	finitely branching trees

- Dyadic rationals in $[0, 1]$ as an initial algebra
- (Freyd) $[0, 1]$ itself as a terminal coalgebra (as a set, poset, totally ordered set or topologically)

Our examples

Functor	Initial algebra	Terminal coalgebra
$1 + X$	naturals	conaturals
$1 + X^2$	finite binary trees	binary trees
$1 + B \times X$	finite lists	lists
$B \times X$	empty	streams
\mathcal{P}_{fin}	V_ω	finitely branching trees

- Dyadic rationals in $[0, 1]$ as an initial algebra
- (Freyd) $[0, 1]$ itself as a terminal coalgebra (as a set, poset, totally ordered set or topologically)
- (Leinster) $L^1[0, 1]$ (with Lebesgue measure) as a terminal coalgebra, and Julia sets

Sufficient conditions?

- Can have a least fixed point and no initial algebra

Sufficient conditions?

- Can have a least fixed point and no initial algebra
- But with some conditions on \mathcal{C} and F , F has a fixed point iff an initial F -algebra exists

Sufficient conditions?

- Can have a least fixed point and no initial algebra
- But with some conditions on \mathcal{C} and F , F has a fixed point iff an initial F -algebra exists
 - for example, if F preserves monomorphisms in **Set**, **Top**

Sufficient conditions?

- Can have a least fixed point and no initial algebra
- But with some conditions on \mathcal{C} and F , F has a fixed point iff an initial F -algebra exists
 - for example, if F preserves monomorphisms in **Set**, **Top**
- But F can preserve monomorphisms and have a fixed point, but no terminal coalgebra

Sufficient conditions?

- Can have a least fixed point and no initial algebra
- But with some conditions on \mathcal{C} and F , F has a fixed point iff an initial F -algebra exists
 - for example, if F preserves monomorphisms in **Set**, **Top**
- But F can preserve monomorphisms and have a fixed point, but no terminal coalgebra

Theorem (Adámek)

Let 0 be initial in \mathcal{C} , and suppose

$$0 \xrightarrow{!} F0 \xrightarrow{F!} F^2 0 \xrightarrow{F^2!} \dots$$

has a colimit, which is preserved by F . Then the colimit carries an initial algebra.


Recursion vs corecursion

- Recursion allows defining a map out of a structure by reducing to easier cases
- Induction defines what to do on constructors
- Corecursion allows defining a map to a complex structure by building up from a seed
- Coinduction defines what destructors do

Further reading


- Initial algebras and terminal coalgebras: a survey (Adámek, Milius, Moss)
- A study of categories of algebras and coalgebras (Hughes)
- A general theory of self-similarity (Leinster)

Sketch proof of Adámek's Theorem



Dyadic rationals

- Category of bipointed sets (X, \top, \perp) with $\top \neq \perp$.
- $FX = X \vee X$



- $\{\perp, \top\}$ is initial object in **BiP**
- Dyadic rationals in $[0, 1]$ are initial algebra

Real interval

- Category of intervals: linearly ordered sets with least and greatest element
- Same functor: glue the greatest element of the first copy to the least element of the second copy
- $F : [0, 1] \mapsto [0, 1] \vee [0, 1] \cong [0, 2]$

$$\begin{array}{ccc}
 A & \longrightarrow & [0, 1] \\
 \downarrow & & \downarrow \\
 A \vee A & \longrightarrow & [0, 2]
 \end{array}$$

$$0.x_1 \cdots x_n 011111 \cdots = 0.x_1 \cdots x_n 100000 \cdots$$