

# Quantum dots

What is a quantum dot?



Controlled tunneling of electrons through the dot:

- Input
- Coulomb blockade
- Spin (Pauli) blockade
- Readout

# Coulomb blockade

Action

$$S[\bar{\Psi}_\alpha, \Psi_\alpha] = \sum_\alpha \bar{\psi}_\alpha (\partial_\tau + \epsilon_\alpha - \mu) \psi_\alpha + E_C \left( \sum_\alpha \bar{\Psi}_\alpha \Psi_\alpha - N_0 \right)^2$$

Hubbard-Stratonovich transformation:

$$\psi^4 \rightarrow \psi^2 V + V^2 + V$$

In this case

$$e^{-E_C(\sum_\alpha \bar{\Psi}_\alpha \Psi_\alpha - N_0)^2} = \int DV e^{-\frac{V^2}{4E_C} + iN_0 V - i \sum_\alpha \bar{\Psi}_\alpha V \Psi_\alpha}$$

# Coulomb blockade

This leads to

$$\begin{aligned}
 G_{\alpha} &= \frac{1}{Z} \int D(\bar{\psi}, \psi) e^{-\underline{S_{free}[\bar{\psi}, \psi]} - \underline{S_{int}[\bar{\psi}, \psi]}} \bar{\psi}_{\alpha} \psi_{\alpha} \\
 &= \frac{1}{Z} \int D(\bar{\psi}, \psi) e^{-\underline{\sum_{\alpha} \bar{\psi}_{\alpha} (\partial_{\tau} + \epsilon_{\alpha} - \mu) \psi_{\alpha}} - \underline{E_C (\sum_{\alpha} \bar{\Psi}_{\alpha} \Psi_{\alpha} - N_0)^2}} \bar{\psi}_{\alpha} \psi_{\alpha} \\
 &= \frac{1}{Z} \int DV e^{-S[V]} \int D(\bar{\psi}, \psi) e^{-\underline{\sum_{\alpha} \bar{\psi}_{\alpha} (\partial_{\tau} + \epsilon_{\alpha} - \mu + iV) \psi_{\alpha}}} \bar{\psi}_{\alpha} \psi_{\alpha} \\
 &= \frac{1}{Z} \int DV e^{-S[V]} \int D(\bar{\psi}, \psi) e^{-S^V[\bar{\psi}, \psi]} \bar{\psi}_{\alpha} \psi_{\alpha}
 \end{aligned}$$

$$S^V[\bar{\psi}, \psi] = S_{free}[\bar{\psi}, \psi]|_{\mu \rightarrow \mu - iV}$$

# Coulomb blockade

Using a gauge transformation of the form

$$\psi(\tau) \rightarrow \psi(\tau) e^{-i \int^\tau d\tau' (V(\tau') - V_0)}$$

The action  $S^V$  can be reduced

$$S^V[\bar{\psi}, \psi] \rightarrow S^{V_0}[\bar{\psi}, \psi]$$

At the cost of an extra gauge factor

$$\frac{1}{Z} \int DV e^{-S[V]} \int D(\bar{\psi}, \psi) e^{-S^{V_0}[\bar{\psi}, \psi]} e^{-i \int^\tau d\tau' (V(\tau') - V_0)} \bar{\psi}_\alpha \psi_\alpha$$

# Coulomb blockade

The integral over the auxiliary field can be split

$$\int DV = \prod_{n=0} \int dV_n = \int dV_0 \prod_{n=1} \int dV_n$$

The integration over the  $V_n$  components gives a Matsubara summation

$$-2E_C T \sum_{n \neq 0} \frac{1}{\omega_n^2} (1 - e^{-i\omega_n \tau}) = -E_C (|\tau| - T \cdot \tau^2)$$

*Note:* this is not a trivial calculation, but the answer is obtained by using

$$\sum_{k=1}^{\infty} \frac{\cos(kx)}{k^2} \approx \frac{\pi^2}{6} - \frac{\pi|x|}{2} + \frac{x^2}{4}$$

# Coulomb blockade

In our action, we can write

$$\begin{aligned} G(\tau) &= \frac{1}{Z} \int DV e^{-S[V]} \int D(\bar{\psi}, \psi) e^{-i \int^\tau d\tau' (V(\tau') - V_0)} e^{-S^{V_0}[\bar{\psi}, \psi]} \bar{\psi}_\alpha \psi_\alpha \\ &= \frac{1}{Z} \int DV e^{-S[V]} \int D(\bar{\psi}, \psi) e^{-i \int^\tau d\tau' (V(\tau') - V_0)} Z^{V_0} G_\alpha^{V_0} \\ &= \frac{F(\tau)}{Z} \int dV_0 e^{-S[V_0]} Z^{V_0} G_\alpha^{V_0} \end{aligned}$$

Where

$$F(\tau) \equiv e^{-E_C(|\tau| - T \cdot \tau^2)}$$

# Coulomb blockade

In

$$G(\tau) = \frac{F(\tau)}{Z} \int dV_0 e^{-S[V_0]} Z^{V_0} G_{\alpha}^{V_0}$$

We can define a free energy  $\mathcal{F}(\mu)$  as

$$Z^{V_0} = e^{-\beta \mathcal{F}^{V_0}(\mu)} = e^{-\beta \mathcal{F}(\mu - iV_0)}$$

So we have, writing out the  $S[V_0]$ :

$$G(\tau) = \frac{F(\tau)}{Z} \int dV_0 e^{-\frac{\beta}{4EC} V_0^2 + i\beta N_0 V_0 - \beta \mathcal{F}(\mu - iV_0)} G_{\alpha}^{V_0}$$

which can be approximated using the stationary phase method

# Stationary phase approximation

When given an integral of the form

$$I = \lim_{\lambda \rightarrow \infty} \int_{-\infty}^{\infty} e^{-\lambda f(x)}$$

The relevant contributions occur around minimum  $x_0$  of  $f$ :

$$\left. \frac{\partial f}{\partial x} \right|_{x=x_0} = 0$$

Which is known as the *saddle point equation*



# Stationary phase approximation

In the case of our action, we have the exponent

$$-\frac{\beta}{4E_C}V_0^2 + i\beta N_0 V_0 - \beta \mathcal{F}(\mu - iV_0)$$

Which has a minimum (after some calculation)

$$0 = \frac{1}{2E_C}V_0 - iN_0 + i\left\langle \hat{N} \right\rangle_{\mu - iV_0}$$

where we used that

$$\frac{\partial}{\partial V_0}\mathcal{F}(\mu - iV_0) = -i\frac{\partial}{\partial \mu}\mathcal{F}(\mu - iV_0) = -i\left\langle \hat{N} \right\rangle_{\mu - iV_0}$$

where  $\hat{N}$  is the number operator

$$\hat{N} \equiv \sum_{\alpha} a_{\alpha}^{\dagger} a_{\alpha}$$

# Density of states

Filling in the solutions, we can calculate the tunneling density of states by

$$\nu(\epsilon) = -\frac{1}{\pi} \text{Im } \text{tr } G$$

After some calculations, we obtain

$$\nu(\epsilon) = \nu_0(\epsilon - E_C \text{sgn}(\epsilon)) \Theta(|\epsilon| - E_C)$$

Where  $\nu_0$  is the density of states (shifted by the energy cost) of the non-interacting systems and  $\Theta$  a step function.

This means that there is an energy gap of size  $2E_C$  around the Fermi energy

# Conclusions

The Coulomb blockade prevents electrons of certain energies to tunnel into the dot.

If it has energy  $\epsilon > E_C$  it is allowed to tunnel