

Assignment 3

1. What is the solution to the difference equation $y[n] = Ky[n-1]$ with initial condition $y[0] = 1$ and $y[-1] = 0$? Next, find the ROC of z-transform of your solution for $y[n]$, $K \leq 1$.

$$y[n] = K^n$$

$$Y(Z) = \frac{1}{1 - KZ^{-1}}$$

The z-transform converges outside of a circle of radius K .

2. Consider a discrete-time signal $x[n]$ passed through a bank of filters $h_k[n]$ where each filter is given by a modulated version of a baseband prototype filter $h[n]$

$$h_k[n] = h[n] \exp \left[j \left(\frac{2n}{N} \right) kn \right]$$

where $h[n]$, a Hamming window, is assumed causal and lies over a duration $0 \leq n < N_w$, and $2n/N$ is the frequency sampling factor. In this problem, you are asked to time-scale expand some simple input signals by time-scale expanding the filter bank outputs.

- (a) State the constraint (with respect to the values N_w and N) such that the input $x[n]$ is recovered when the filter bank outputs are summed.
- (b) If the input to the filter bank is the unit sample $\delta[n]$, then the output of each filter is a complex exponential with "envelope" $a_k[n] = h[n]$ and phase $\theta_k[n] = (2n/N)kn$. Suppose each complex exponential output is time-expanded by two by interpolation of its envelope and phase. Derive a new constraint (with respect to values N_w and N), so that the summed filter bank outputs equal $\delta[n]$.
- (c) Suppose now that the filter bank input is

$$x[n] = \delta[n] + \delta[n-n_0]$$

and that the filter bank outputs are time-expanded as in part (b). Derive a sufficient condition on N_w , N , and n_0 so that the summed filter bank output is given by

$$y[n] = \delta[n] + \delta[n-2n_0]$$

that is, the unit samples are separated by $2n_0$ samples rather than n_0 samples.

3. Recall the "voiced" excitation to the digital model of speech production,

$$e[n] = \sum_{q=-\infty}^{\infty} \delta[n - qP]$$

Find the expression for the long term temporal auto-correlation, $r_e(\eta)$.

4. (MATLAB) In this MATLAB exercise,

- (a) Perform the following operations with the vowel utterance of your choice:
 - i. Compute the short term auto-correlation function for a hamming window of $N = 512$ for $\eta = 0, 1, 2, \dots, 256$.

- ii. Compute the $N = 512$ point magnitude spectrum of the waveform based on hamming window and stDFT.
(Note: The stDFT and conventional DFT are equivalent here because only the magnitude spectrum is required.)
 - iii. Repeat step (i) and (ii) after centre clipping the waveform.
- (b) Comment the changes in both the auto-correlation and the spectrum. What do these changes indicate about the effects of the clipping operations on the waveform?
- (c) Estimate the pitch using the two auto-correlation results. Which result would provide better performance in an auto-correlation procedure?