

Emergence of Classical Constants from a Minimal Recursive Triadic Framework

J. W. Milton

Clarity Coalition

10 June 2026

v1.2

Abstract

Five classical mathematical constants (the Leibniz limit $\pi/4$, the golden ratio φ , Euler's number e , $\sqrt{2}$, and $\ln 2$) emerge as limits of a single family of three-variable recurrences on a triple (N, D, C) with functionally distinct roles: a *negotiation* state that is iteratively refined, a *definition/limitation* parameter that bounds, and a *contribution/integration* parameter that accumulates. These roles are irreducible to fewer than three even when two positions share the same numerical seed value.

Each branch is specified by an initial triple (N_0, D_0, C_0) and a *traversal rule* governing the evolution of (D_k, C_k) . The five branches fall into three traversal classes: Advancing (Class A, external parameter injection), Self-redefined (Class B, internal transformation), and Fixed (Class C, constant parameters). The limits are classical: each branch reduces to a known series, fixed point, or iterative method, so the contribution of this framework is not in computing these numbers anew, but in *structural unification*: a single three-role grammar from which all five mechanisms descend, combined with proved theorems (diagonal invariance, swap symmetry, perfect-square denominators, offset consistency, parametric families).

The $\pi/4$ branch is structurally unique: it alone requires three numerically distinct seeds $\{1, 3, 5\}$ and advances its parameters externally via a fixed step $\Delta = 4$ rooted in a geometric axis structure. The remaining four branches operate on seeds from $\{0, 1, 2\}$ with purely

internal or fixed parameter evolution. Four open conjectures address whether these assignments are forced by the framework or merely convenient choices.

MSC 2020: 40A05 (primary); 11B39, 65B10 (secondary).

Keywords: classical constants; recursive sequences; triadic recurrence; Fibonacci ratio; alternating series; structural unification.

Contents

1	Introduction	3
2	The tholonic triad	4
2.1	Geometric origin of the axis labels	4
3	The irreducible three-variable structure	6
4	The ladder family	7
4.1	Classification of traversal types	7
5	Branch specification and convergence proofs	8
5.1	The $\pi/4$ branch	9
5.2	The φ branch	10
5.3	The $\sqrt{2}$ branch	11
5.4	The e branch	11
5.5	The $\ln 2$ branch	11
6	Structural properties	12
7	Convergence	15
8	Open problems	15
9	Related work	17
10	Discussion	17
11	Conclusion	18
A	Reference implementation	18

1 Introduction

The Leibniz series for $\pi/4$,

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \cdots = \sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1},$$

has been known since the late seventeenth century and remains a landmark example of slowly convergent alternating series [6, 9]. Leibniz himself saw philosophical significance in binary arithmetic and its connection to the I Ching [7], viewing the interplay of 0 and 1 as a model of structured emergence from minimal premises.

Five real numbers recur persistently across mathematics: π , the golden ratio $\varphi = (1 + \sqrt{5})/2$, Euler’s number e , $\sqrt{2}$, and $\ln 2$. Each has multiple independent characterizations and has been analyzed in isolation over centuries. The question of whether a *single algebraic apparatus* organizes all five as co-equal outputs of one recursive scheme is less a question of computing them anew than one of *structural unification*: does a common grammar exist?

This paper answers that question affirmatively for the *tholonic ladder*: a family of three-variable recurrences whose branches are distinguished by different initial triples and traversal rules, but whose variable roles remain consistent across all five instances.

What this paper provides. Minimal definitions; a complete branch-specification table; full proofs of convergence to each classical limit; structural theorems with proofs; quantitative convergence data; and four open conjectures.

What this paper does not provide. New proofs of the underlying series identities: those are classical and referenced. The contribution is the shared grammar and the structural contrasts.

Organization. §2 defines the tholonic triad and motivates its geometry. §3 argues the irreducibility of three variables. §4 introduces the ladder family and its traversal-class taxonomy. §5 provides the complete branch specification and proves the five convergence propositions. §6 establishes the structural properties. §7 records convergence data. §8 states four open

conjectures. § 9 covers related work. § 10 discusses scope and implications. § 11 concludes.

2 The tholonic triad

Definition 2.1 (Tholonic triad). A *tholonic triad* is an ordered triple (N, D, C) of non-negative reals together with three directed relational roles:

- N (*negotiation*): the running state; the quantity being iteratively refined, emerging from the interaction of the other two.
- D (*definition/limitation*): the constraining, bounding force; what *limits* the state.
- C (*contribution/integration*): the accumulating, synthesizing force; what *grows* the state.

The triad is not merely a vector: the three positions have distinct semantic roles that persist across all five branches, regardless of whether two positions share a numerical value.

2.1 Geometric origin of the axis labels

The tholonic geometry [8] assigns successive powers of two, $2^0, 2^1, \dots, 2^5$, to six vertices of a triangular configuration: outer role vertices $N = 2^0$, $D = 2^1$, $C = 2^2$, and three inner midpoint vertices carrying $2^3, 2^4, 2^5$. Each directed axis runs from one outer role vertex through an inner midpoint to another outer role vertex; its three vertex values sum to 7 times a characteristic multiplier:

- Definition axis ($N \rightarrow D$, through 2^5): $1 + 32 + 2 = 35 = 7 \times 5$
- Contribution axis ($C \rightarrow N$, through 2^4): $4 + 16 + 1 = 21 = 7 \times 3$
- Instantiation axis ($D \rightarrow C$, through 2^3): $2 + 8 + 4 = 14 = 7 \times 2$

The Instantiation axis multiplier, denoted $\text{istep} = 2$, enters the $\pi/4$ branch:

$$d_{\text{step}} = \text{istep}^2 = 4, \quad c_{\text{step}} = 2 \times \text{istep} = 4.$$

Both equal 4, fixing step size $\Delta = 4$ without free parameters. The seed triple $(N_0, D_0, C_0) = (1, 3, 5)$ receives a geometric interpretation. Each axis connects two of the three roles and excludes the third, and its multiplier supplies the seed (or step) of precisely the excluded role: the Definition axis ($N \rightarrow D$) excludes C and carries $5 = C_0$; the Contribution axis ($C \rightarrow N$) excludes D and carries $3 = D_0$; the Instantiation axis ($D \rightarrow C$) excludes N and carries $\text{istep} = 2$, which fixes $\Delta = 4$ above. The seeds 3 and 5 are simultaneously the first members of the $(4n-1, 4n+1)$ parity classes traversed by the recursion, and $N_0 = 1$ is the multiplicative identity. Each role draws its seed from the complementary axis, so the axis geometry supplies both the seeds and the step that drives them. This motivates the seed-partition theorem of Proposition 6.1.




Figure 1. Left: the tholonic triad. Outer vertices carry roles N, D, C with values $2^0, 2^1, 2^2$; inner midpoint vertices carry $2^3, 2^4, 2^5$. Each directed axis sums to a multiple of 7 (multipliers 5, 3, 2). Right: the Instantiation axis multiplier $\text{istep} = 2$ uniquely fixes the $\pi/4$ branch step $\Delta = 4$ and the seed triple $(N_0, D_0, C_0) = (1, 3, 5)$.

Figure 1: Left: the tholonic triad. Outer vertices carry roles N, D, C with values $2^0, 2^1, 2^2$; inner midpoint vertices carry $2^3, 2^4, 2^5$. Each directed axis sums to a multiple of 7 (multipliers 5, 3, 2). Right: the Instantiation axis multiplier $\text{istep} = 2$ uniquely fixes $\Delta = 4$ and the seed triple $(N_0, D_0, C_0) = (1, 3, 5)$.

3 The irreducible three-variable structure

Lemma 3.1 (Triadic irreducibility). *Let $\mathcal{R}$ be a recurrence on m real variables $x^{(1)}, \dots, x^{(m)}$ in which the update to one distinguished state variable $x^{(1)}$ is of the form*

$$x_{k+1}^{(1)} = g\left(x_k^{(1)}; \alpha_k, \beta_k\right),$$

where α_k limits the correction magnitude and β_k controls the additive or multiplicative integration of each step. Suppose that (i) α_k and β_k are functionally independent in the sense that g is not expressible as a function of a single combined argument $h(\alpha_k, \beta_k)$, and (ii) the limit $L = \lim_{k \rightarrow \infty} x_k^{(1)}$ is not a trivial fixed point of a one- or two-variable recurrence. Then $\mathcal{R}$ requires $m \geq 3$.

Proof. The distinguished variable $x^{(1)}$ must persist across iterations to record the refined state, so it cannot be eliminated. Suppose $m = 2$. Then both control arguments are determined by the single auxiliary variable y_k : $\alpha_k = a(y_k)$ and $\beta_k = b(y_k)$, and the admissible pairs (α_k, β_k) are confined to the one-parameter image of $y \mapsto (a(y), b(y))$. On that set, g factors through the single argument y itself: $g(x; \alpha_k, \beta_k) = \tilde{g}(x; y_k)$, exhibiting exactly the combined-argument factorization that hypothesis (i) excludes on the admissible set. Hypothesis (ii) rules out the degenerate alternative in which the controls are constant or absent and L is reachable with fewer variables. Hence two functionally independent auxiliary variables are required, and with the state variable, $m \geq 3$. $\square$

A critical structural property is that **every branch requires exactly three variables**, regardless of whether some share the same numerical value.

In the e branch, $D_0 = C_0 = 1$; in the $\sqrt{2}$ branch, $D_0 = C_0 = 2$. However, the triad is not about three *numbers*; it is about three *functional roles*:

1. N : the state being negotiated; what emerges.
2. D : the defining boundary; what constrains.
3. C : the integrating accumulator; what synthesizes.

In the e branch, $D = 1$ is the fixed numerator that never changes; the unchanging boundary. C starts at 1 but grows as a factorial. They begin at the same value but diverge immediately in behavior: one constrains, the other integrates.

In the $\sqrt{2}$ branch, $D = 2$ is the target being bounded toward; the defining constraint. $C = 2$ is the averaging divisor; the integrator that synthesizes overshoot and undershoot. Without D/N , there is no correction; without C -division, no synthesis.

The triad is irreducible: a state, a limiter, and a contributor constitute the minimum structure for a recursion that converges to a non-trivial constant.

4 The ladder family

Definition 4.1 (Tholonic ladder recurrence). Given $(N_0, D_0, C_0) \in \mathbb{R}^3$, a *tholonic ladder branch* is a discrete dynamical system

$$N_{k+1} = f(N_k; D_k, C_k), \quad k \in \mathbb{N}_0,$$

together with a *traversal rule* specifying how (D_k, C_k) is updated at each step.

Definition 4.2 (Branch). A *branch* is the tuple $((N_0, D_0, C_0), f, \text{traversal rule})$.

4.1 Classification of traversal types

Class A: Advancing. (D_k, C_k) each receive a fixed external increment at every step. New information is injected per iteration. Only the $\pi/4$ branch belongs to this class.

Class B: Self-redefined. (D_k, C_k) evolve by a transformation of their own current values (Fibonacci swap for φ , factorial scaling for e) with no external increment. The factorial rule consumes the iteration counter k as a multiplier, but injects no new constant beyond it.

Class C: Fixed. (D_k, C_k) are held constant at their seed values for all k . Convergence is driven by the update map f alone: as a pure fixed-point iteration in the $\sqrt{2}$ branch (where k never enters f), or with the iteration index entering f directly in the $\ln 2$ branch.

This trichotomy partitions the five branches cleanly: one Class A ($\pi/4$), two Class B (φ, e), two Class C ($\sqrt{2}, \ln 2$). The classification can equivalently be stated on two independent axes: whether (D_k, C_k) evolve at all, and whether the branch consumes the iteration counter k . Class A is evolution by external constant; Class B is evolution by internal or counter-indexed transformation; Class C is no evolution.

Table 1: Two-axis refinement of the traversal classification.

Branch	(D, C) evolve?	Counter k consumed?	Class
$\pi/4$	yes (external increment Δ)	no	A
φ	yes (internal swap)	no	B
e	yes (scaling by k)	yes, in traversal	B
$\sqrt{2}$	no	no	C
$\ln 2$	no	yes, in f	C

5 Branch specification and convergence proofs

Table 2: Complete branch specification.

Limit	N_0	D_0	C_0	N_{k+1}	Traversal rule	Class
$\pi/4$	1	3	5	$N - \frac{1}{D} + \frac{1}{C}$	$D \leftarrow D + 4, \quad C \leftarrow C + 4$	A
φ	1	1	2	$N_0 + \frac{D}{C}$	$D \leftarrow C, \quad C \leftarrow C + D$	B
e	0	1	1	$N + \frac{D}{C}$	$C \leftarrow C \cdot \max(k, 1)$	B
$\sqrt{2}$	1	2	2	$\frac{N + D/N}{C}$	D, C fixed	C
$\ln 2$	0	1	1	$N + (-1)^k \frac{D}{k + C}$	D, C fixed	C

In every branch D plays the bounding/constraining role and C the integrating/synthesizing role, despite the operations differing. This functional consistency is not imposed; it emerges from the mathematics.




Figure 2: Classification of the five tholonic ladder branches by traversal type. Class A (red) injects new information each step via an external increment. Class B (blue) evolves parameters by an internal rule. Class C (green) holds all parameters fixed; convergence is driven by the update map alone, with the iteration index entering f directly in the $\ln 2$ branch.

5.1 The $\pi/4$ branch

Proposition 5.1. *With $(N_0, D_0, C_0) = (1, 3, 5)$ and step $\Delta = 4$,*

$$N_\infty = 1 + \sum_{n=1}^{\infty} \left(\frac{1}{4n+1} - \frac{1}{4n-1} \right) = 1 + \sum_{n=1}^{\infty} \frac{-2}{16n^2 - 1} = \frac{\pi}{4}.$$

Proof. At iteration $n \geq 1$, $D_n = 4n - 1$ and $C_n = 4n + 1$. The update adds $-1/(4n - 1) + 1/(4n + 1) = -2/(16n^2 - 1)$. Starting from $N_0 = 1$, the accumulated sum is a grouping of consecutive Leibniz pairs. By the alternating series theorem applied to $\sum_{k=0}^{\infty} (-1)^k/(2k + 1)$, the grouped sum converges to $\pi/4$ [4]. $\square$

Remark 5.2 (Primitive invariant). The recurrence targets $\pi/4$ directly; $\pi = 4(\pi/4)$ is a geometric normalization. We treat $\pi/4$ as the *primitive invariant* of this branch.

Corollary 5.3 (Perfect-square form). *Multiplying both sides of Proposi-*

tion 5.1 by 8:

$$\pi = \sum_{n=1}^{\infty} \frac{8}{16n^2 - 1},$$

expressing π in terms of perfect squares offset by unity. See also Proposition 6.5.

Functional roles. D_k subtracts from N_k (constraining); C_k adds (contributing). Both advance in lockstep, driven by external step $\Delta = 4$.




Figure 3: Partial sums N_k of the $\pi/4$ branch converging to $\pi/4$ (dashed). The damped alternating behavior is characteristic of conditionally convergent series; amplitude decays as $O(1/k)$. Red diamonds mark $k \in \{5, 25, 125\}$; the shaded band shows the $\pm 1/(2k+1)$ envelope.

Figure 3: Partial sums N_k of the $\pi/4$ branch converging to $\pi/4$ (dashed). Amplitude decays as $O(1/k)$. Red diamonds mark $k \in \{5, 25, 125\}$; the shaded band shows the $\pm 1/(2k+1)$ envelope.

5.2 The φ branch

Proposition 5.4. *With $(N_0, D_0, C_0) = (1, 1, 2)$, update $N_{k+1} = N_0 + D_k/C_k$, and the Fibonacci traversal $D \leftarrow C$, $C \leftarrow C + D$, the limit is $N_\infty = \varphi = (1 + \sqrt{5})/2$.*

Proof. With seeds $(D_0, C_0) = (1, 2) = (F_2, F_3)$, the traversal maintains $(D_k, C_k) = (F_{k+2}, F_{k+3})$, consecutive Fibonacci numbers. Ratios of consecutive Fibonacci numbers converge: $F_{k+2}/F_{k+3} \rightarrow 1/\varphi$ [2, 5], with error contracting geometrically by the factor $\varphi^{-2} \approx 0.382$ per step (Binet's formula). Hence $N_{k+1} = 1 + F_{k+2}/F_{k+3} \rightarrow 1 + 1/\varphi = \varphi$, by the defining identity $\varphi = 1 + 1/\varphi$. $\square$

Functional roles. D is the smaller (prior) Fibonacci term; the boundary. C is the larger accumulated term; the integration of the two most recent

generations. The update consumes the ratio D/C directly, so the Fibonacci traversal is the engine of convergence, not an accessory to it.

5.3 The $\sqrt{2}$ branch

Proposition 5.5. *With $(N_0, D_0, C_0) = (1, 2, 2)$ and fixed parameters, $N_\infty = \sqrt{2}$, with quadratic convergence.*

Proof. The update $N_{k+1} = (N_k + 2/N_k)/2$ is the Newton–Babylonian method for $x^2 = 2$ [1, 3]. Let $g(x) = (x + 2/x)/2$. Then $g(\sqrt{2}) = \sqrt{2}$ and $g'(\sqrt{2}) = (1 - 2/x^2)/2|_{x=\sqrt{2}} = 0$, confirming quadratic convergence. $\square$

Functional roles. $D = 2$ is the target; the defining constraint. $C = 2$ is the averaging divisor. Though $D = C$ numerically, their roles are distinct.

5.4 The e branch

Proposition 5.6. *With $(N_0, D_0, C_0) = (0, 1, 1)$ and $C_k = k!$, $N_\infty = e$.*

Proof. The update $N_{k+1} = N_k + 1/k!$ accumulates $\sum_{k=0}^K 1/k!$, converging to e [10]. $\square$

Functional roles. $D = 1$ is the fixed numerator; the unchanging boundary. C starts at 1 but grows as a factorial: the expanding denominator.

Remark 5.7 (Offset variant). When initialized with $N_0 = 2$, the branch converges to $e + 2$, confirming the offset $= N_0$ (Proposition 6.6).

5.5 The $\ln 2$ branch

Proposition 5.8. *With $(N_0, D_0, C_0) = (0, 1, 1)$ and fixed parameters, $N_\infty = \ln 2$.*

Proof. The update accumulates $(-1)^k/(k+1)$ starting at $k = 0$, giving $\sum_{k=0}^{\infty} (-1)^k/(k+1) = \ln 2$ [4]. $\square$

Functional roles. $D = 1$ is the fixed numerator; $C = 1$ is the integration base. They are numerically identical but structurally distinct: one limits, one integrates.

Remark 5.9 (Offset variant). When $N_0 = 1/2$, the branch converges to $\ln 2 + 1/2$. The offset equals N_0 exactly, as it must: the update is pure accumulation, so the seed enters the limit additively (Proposition 6.6).

6 Structural properties

Proposition 6.1 (Seed partition). *Among the five branches, the $\pi/4$ branch is the unique branch with three numerically distinct seeds; its seed set is $\{1, 3, 5\}$. The remaining four branches have seeds drawn from $\{0, 1, 2\}$.*

Table 3: Seed partition across the five branches.

Branch	Seeds	Distinct values	Seed set
$\pi/4$	(1, 3, 5)	3	$\{1, 3, 5\}$
φ	(1, 1, 2)	2	$\{1, 2\}$
e	(0, 1, 1)	2	$\{0, 1\}$
$\sqrt{2}$	(1, 2, 2)	2	$\{1, 2\}$
$\ln 2$	(0, 1, 1)	2	$\{0, 1\}$

The value 1 appears 10 out of 15 times. Only $\pi/4$ demands the richer seed set $\{1, 3, 5\}$ and external parameter injection.

Remark 6.2 (Seeds and primality). The seeds 1, 3, 5 are the first three odd integers ≥ 1 . Whether 1 is treated as “prime” does not affect the convergence proof; it is the generative unit from which the triadic recursion begins.

Proposition 6.3 (Diagonal invariance). *For the Class A recurrence with $D_k = C_k$ for all k , the partial sums telescope to zero and $N_\infty = N_0$. Non-triviality requires $D \neq C$.*

Proof. At each step, $-1/D_k + 1/C_k = 0$ when $D_k = C_k$. The running total is unchanged. $\square$

Proposition 6.4 (Swap symmetry). *For the Class A recurrence with fixed N_0 , let $N_\infty(D_0, C_0)$ denote the limit. Then $N_\infty(D_0, C_0) + N_\infty(C_0, D_0) = 2N_0$.*

Proof. Swapping $D \leftrightarrow C$ negates the signed contribution $-1/D_k + 1/C_k$ at every step, so the two accumulated partial sums are antisymmetric around N_0 . $\square$

Proposition 6.5 (Perfect-square denominators). *In the $\pi/4$ branch, $D_n \cdot C_n = (4n - 1)(4n + 1) = 16n^2 - 1$ for $n \geq 1$. This yields*

$$\pi = \sum_{n=1}^{\infty} \frac{8}{16n^2 - 1}.$$




Figure 4: Swap symmetry (Proposition 6.3): the two runs $(D_k, C_k) = (3, 5)$ and $(5, 3)$ mirror around the midline $N_k = 1$ at every iteration k . Run A decreases monotonically to $\pi/4$. Run B increases monotonically to $2 - \pi/4$. Equal brackets δ_k confirm $N_A(k) + N_B(k) = 2N_0 = 2$ at each step. Inset: numerical residual $|N_k + N_k - 2|$ is at floating-point zero throughout.

Figure 4: Swap symmetry (Proposition 6.4): runs $(3, 5)$ and $(5, 3)$ mirror around $N_0 = 1$. Run A decreases to $\pi/4$; Run B increases to $2 - \pi/4$. Equal brackets confirm $N_A(k) + N_B(k) = 2$ at each step.

Proof. With $D_n = 4n - 1$ and $C_n = 4n + 1$, the product is $16n^2 - 1$. The identity for π follows from Proposition 5.1 by algebraic rearrangement. $\square$

Table 4: Denominator products.

n	D_n	C_n	$D_n \cdot C_n$	Perfect-square form
1	3	5	15	$16(1)^2 - 1$
2	7	9	63	$16(2)^2 - 1$
3	11	13	143	$16(3)^2 - 1$

Proposition 6.6 (Offset consistency, translation equivariance). *Let a branch have a pure-accumulation update $N_{k+1} = N_k + t_k$, where the term t_k depends only on (D_k, C_k, k) and not on N_k (the e and $\ln 2$ branches). Then the limit is translation-equivariant in the seed:*

$$N_\infty(N_0) = N_0 + N_\infty(0),$$

so the affine offset of N_∞ from the target constant equals N_0 exactly.

Proof. By induction, $N_K = N_0 + \sum_{k=0}^{K-1} t_k$, and the sum is independent of N_0 . Letting $K \rightarrow \infty$ gives the identity. $\square$

In both cases the offset is the initial baseline. In the canonical forms (Table 2) with $N_0 = 0$, both converge directly to their targets.




Figure 5: The denominator products $D_n \cdot C_n = 16n^2 - 1$ of the left branch at scales $K = 10, 100, 1000$ (left to right). Each scale reproduces the same parabolic shape, reflecting the self-similar structure of the recursion. The dotted red curve shows the reference parabola $16n^2$; the products are uniformly offset by -1 (perfect squares minus unity).

Figure 5: The products $D_n \cdot C_n = 16n^2 - 1$ at scales $K = 10, 100, 1000$. Each scale reproduces the same parabolic shape. Dotted red: reference parabola $16n^2$.

Table 5: Offset behavior.

Branch	Variant N_0	N_∞	Offset
e	2	$e + 2$	$+2 = N_0$
$\ln 2$	$1/2$	$\ln 2 + 1/2$	$+1/2 = N_0$

Proposition 6.7 (Parametric families). *Two branches embed in one-parameter families with the same traversal class and role structure: (i) replacing $D = 2$ by $D = a > 0$ in the $\sqrt{2}$ branch yields $N_\infty = \sqrt{a}$, with quadratic convergence, for every a ; (ii) replacing the additive base $N_0 = 1$ by $m \in \mathbb{N}$ in the φ branch, with the generalized traversal $D \leftarrow C$, $C \leftarrow mC + D$, yields the metallic means $N_\infty = \sigma_m = (m + \sqrt{m^2 + 4})/2$; the case $m = 1$ recovers φ .*

Proof. (i) The update $N_{k+1} = (N_k + a/N_k)/C$ with $C = 2$ is Newton's method for $x^2 = a$ [1]; the argument of Proposition 5.5 applies verbatim, with $g'(\sqrt{a}) = 0$ giving quadratic convergence. (ii) The traversal maintains consecutive terms of the generalized Fibonacci recurrence $G_{k+1} = mG_k + G_{k-1}$, whose consecutive ratios converge (for any positive seeds, since σ_m is the dominant root) to the positive root of $x^2 = mx + 1$: thus $D_k/C_k \rightarrow 1/\sigma_m$, and $N_{k+1} = m + D_k/C_k \rightarrow m + 1/\sigma_m = \sigma_m$ by the defining identity $\sigma_m = m + 1/\sigma_m$. $\square$

These deformations show that the documented branches are distinguished points inside parametric families rather than isolated curiosities, and they make precise why the local-isolation reformulation of Conjecture 8.3 is the natural one: within a family, varying the parameter moves the limit contin-

uously through other algebraic targets, so uniqueness claims must be made relative to fixed parameters.

7 Convergence

Table 6: Numerical convergence at $K = 10^5$ iterations.

Branch	N_K (computed)	Target	Residual	Rate
$\pi/4$	0.785 399 4 ...	0.785 398 2 ...	1.2×10^{-6}	$O(1/K)$
φ	1.618 033 9 ...	1.618 033 9 ...	$< 10^{-15}$	linear, rate φ^{-2}
e	2.718 281 8 ...	2.718 281 8 ...	$< 10^{-15}$	$O(1/K!)$
$\sqrt{2}$	1.414 213 6 ...	1.414 213 6 ...	$< 10^{-15}$	quadratic
$\ln 2$	0.693 142 2 ...	0.693 147 2 ...	5.0×10^{-6}	$O(1/K)$

The slow convergence of $\pi/4$ and $\ln 2$ is intrinsic to their alternating-series structure. The three remaining branches converge to machine precision well within 10^5 iterations. Series-acceleration methods (Euler, van Wijngaarden, Levin) would reduce residuals for the two slow branches but are not required by the framework.

Explicit error bounds. For the two alternating branches the alternating series theorem supplies a priori envelopes: after k iterations, $|N_k - \pi/4| \leq 1/(4k + 3)$ (the first omitted Leibniz term) and $|N_k - \ln 2| \leq 1/(k + 1)$. At $K = 10^5$ these give 2.5×10^{-6} and 10^{-5} , consistent with the observed residuals 1.2×10^{-6} and 5.0×10^{-6} (roughly half the envelope, as expected for an alternating tail). For $\sqrt{2}$, Newton’s method gives the exact one-step relation $|N_{k+1} - \sqrt{2}| = |N_k - \sqrt{2}|^2/(2N_k)$; for φ , the error contracts geometrically by the factor $\varphi^{-2} \approx 0.382$ per step.

Convergence and traversal class. The two slowest branches are Class A ($\pi/4$) and one Class C branch ($\ln 2$). The fastest are Class B (e , super-exponential; φ , linear with geometric ratio $\varphi^{-2} \approx 0.382$) and the other Class C branch ($\sqrt{2}$, quadratic).

8 Open problems

The structural results of § 6 are proved. The following questions remain open; positive resolutions would move the framework from taxonomy to theorem.




Figure 6: Convergence of all five branches: $\log_{10} |N_k - N_\infty|$ vs k at $K = 10^5$ iterations. Class B branches (e, g) reach machine precision within tens of steps; Class C Newton–Babylonian ($\sqrt{2}$) converges quadratically and hits the floor by $k \approx 6$. Class A ($\pi/4$) and the Class C alternating branch ($\ln 2$) decay as $O(1/k)$, tracking the dotted slope -1 reference line throughout.

Figure 6: Convergence of all five branches: $\log_{10} |N_k - N_\infty|$ vs. k at $K = 10^5$ iterations. Class B branches reach machine precision within tens of steps; Class C Newton–Babylonian ($\sqrt{2}$) converges quadratically by $k \approx 6$. Class A ($\pi/4$) and Class C ($\ln 2$) decay as $O(1/k)$.

Conjecture 8.1 (Uniqueness of the $\pi/4$ seed). *The seed triple $(1, 3, 5)$ is the unique Class A admissible seed with $D_0 < C_0$, $C_0 - D_0 = 2$, D_0 and C_0 consecutive in the $\{4n - 1\}$ and $\{4n + 1\}$ sequences, that yields $N_\infty = \pi/4$ under step size $\Delta = 4$.*

Conjecture 8.2 (Exclusion from the $\{0, 1, 2\}$ lattice). *No triple $(N_0, D_0, C_0) \in \{0, 1, 2\}^3$ with any Class B or Class C traversal rule yields $N_\infty = \pi/4$. That is, $\pi/4$ is genuinely inaccessible from the primitive seed lattice.*

Conjecture 8.3 (Finite classification). *There exists a finite rule set $\mathcal{R}$ such that the only convergent branches with limits in $\{\pi/4, \varphi, e, \sqrt{2}, \ln 2\}$ are exactly the five documented here.*

If Conjecture 8.3 fails, the appropriate reformulation is: show the five branches are *locally isolated* within the admissible space.

Conjecture 8.4 (Non-circularity of the $\pi/4$ grouping). *The pairing $N_{k+1} = N_k - 1/D_k + 1/C_k$ with Class A advancement can be derived from constraints (a) f is rational of degree ≤ 1 in D^{-1} and C^{-1} , (b) D and C carry opposite signs, and (c) step Δ is derived from the Instantiation axis, without assuming π as input.*

Conjecture 8.4 is the key independence question. Establishing it would show that the $\pi/4$ grouping is structurally forced, not chosen post hoc.

9 Related work

Series rearrangements. The Leibniz series is conditionally convergent, so rearrangements can alter its sum [4]. The $\pi/4$ branch uses *grouped* (paired) rearrangement, not a sign-pattern reordering, so its limit is standard. **Continued fractions and metallic means.** The golden ratio is the simplest nontrivial continued fraction, $\varphi = [1; 1, 1, 1, \dots]$, and the worst-approximable irrational [2]. The φ branch recovers this via Fibonacci recursion without competing with continued-fraction theory [5]; the metallic means extend the same template parametrically (Proposition 6.7). **Newton–Babylonian method.** The $\sqrt{2}$ branch is the classical Heron iteration for $x^2 = 2$ [1, 3]. The tholonic framing adds role interpretation without altering the proof or convergence order. **Classical series identities.** Standard analysis establishes $\sum_{k=0}^{\infty} 1/k! = e$ [10] and $\sum_{k=1}^{\infty} (-1)^{k+1}/k = \ln 2$ [4]. The ladder presents these as Class B and C branches in a unified format. **Structural unification.** We are unaware of a framework that identifies all five of $\pi/4$, φ , e , $\sqrt{2}$, $\ln 2$ as branches of a common three-variable recurrence with consistent role assignments. **Leibniz and binary structure.** Leibniz developed binary arithmetic simultaneously with the $\pi/4$ series and drew connections to the I Ching [7, 11].

10 Discussion

Organizational vs. predictive claims. The ladder framework as proved is *organizational*: a unified format from which five known limits descend. The path from organizational to predictive runs through the Conjectures of §8. **Post hoc flexibility risk.** The standard referee objection is: “you chose seeds and rules after knowing the targets.” The structural theorems partially defuse this. Diagonal invariance and swap symmetry are consequences of the Class A form independent of any targeted limit. The seed partition (Proposition 6.1) is a sharp combinatorial observation: a framework tuned post hoc would not automatically produce a clean $\{1, 3, 5\}$ vs. $\{0, 1, 2\}$ partition.

The remaining vulnerability is Conjecture 8.4. Until non-circularity is established for the $\pi/4$ grouping, one cannot rule out that it is one among infinitely many valid Leibniz rearrangements that happen to fit the Class A format.

Convergence rates as a structural signal. The two $O(1/k)$ branches

are Class A ($\pi/4$) and a Class C branch ($\ln 2$); the other three reach machine precision within tens of iterations (e super-exponentially, $\sqrt{2}$ quadratically, φ linearly at geometric ratio φ^{-2}). The per-step ordering places Class C $\sqrt{2}$ (quadratic) above Class B φ (linear), so the correlation between traversal class and convergence order is imperfect; characterizing exactly when class predicts rate is an open regularity.

The infinite landscape. The five branches are recognizable loci in an infinite space of admissible seed–traversal combinations. Proposition 6.7 exhibits two parametric families explicitly: the square roots $\sqrt{a}$ and the metallic means σ_m . The scientific question is which of the infinite admissible paths yield limits of independent mathematical significance, and whether that set is finite and well-characterized.

11 Conclusion

We have presented a three-variable recurrence family (N, D, C) , the tholonic ladder, with five branches converging to $\pi/4$, φ , e , $\sqrt{2}$, and $\ln 2$. The limits are classical; the contribution lies in the structural theorems: a seed partition cleanly separating the $\pi/4$ branch, diagonal invariance and swap symmetry for the Class A template, a perfect-square denominator identity linking π to the step structure, an offset-consistency law for shifted initializations, and two parametric families (square roots and metallic means) embedding the $\sqrt{2}$ and φ branches.

The central claim (**same recursive triadic geometry, different traversal dynamics, different limits**) is supported by these proved contrasts. The open work, stated as four numbered conjectures, concerns whether the seed assignments and traversal rules are forced by a small set of axioms (predictive framework) or represent a human-chosen sample (organizational taxonomy). The structural evidence points toward the former; the proofs are not yet in hand.

A Reference implementation

The following Python code implements all five branches. No bare numeric literals appear inside any recursion loop.

```
1 |#!/usr/bin/env python3
```

```

2
3 def _compute_pi_over_4(max_iter, istep=2):
4     N, D, C = 1.0, 3.0, 5.0
5     d_step = istep ** 2; c_step = istep * 2
6     for _ in range(max_iter):
7         N = N - (1.0 / D) + (1.0 / C)
8         D += d_step; C += c_step
9     return N
10
11 def _compute_phi(max_iter):
12     N = 1.0
13     D, C = 1, 2 # exact integer Fibonacci terms (floats
14                 # overflow near k = 1476)
15     base = N
16     for _ in range(max_iter):
17         N = base + (D / C); D, C = C, (C + D)
18     return N
19
20 def _compute_e(max_iter):
21     N, D, C = 0.0, 1.0, 1.0
22     for count in range(max_iter):
23         N = N + (D / C); C *= count + 1 if count > 0 else 1
24     return N
25
26 def _compute_sqrt2(max_iter):
27     N, D, C = 1.0, 2.0, 2.0
28     for _ in range(max_iter):
29         N = (N + (D / N)) / C
30     return N
31
32 def _compute_ln2(max_iter):
33     N, D, C = 0.0, 1.0, 1.0
34     for count in range(max_iter):
35         term = D / (count + C)
36         N = N + term if count % 2 == 0 else N - term
37     return N

```

Listing 1: Reference implementation.

Output at 10^5 iterations:

```

pi/4   = 0.7853994133911905
phi    = 1.618033988749895
e      = 2.7182818284590455
sqrt2  = 1.414213562373095

```

$$\ln 2 = 0.6931421805849816$$

B Summary table with convergence rates

Table 7: Summary of all five branches.

Target	(N_0, D_0, C_0)	Seed set	Class	Classical mechanism	Rate
$\pi/4$	(1, 3, 5)	$\{1, 3, 5\}$	A	Leibniz paired grouping	$O(1/K)$
φ	(1, 1, 2)	$\{1, 2\}$	B	Fibonacci ratio convergence	linear, rate φ^{-2}
e	(0, 1, 1)	$\{0, 1\}$	B	$\sum_{k=0}^{\infty} 1/k!$	$O(1/K!)$
$\sqrt{2}$	(1, 2, 2)	$\{1, 2\}$	C	Newton–Babylonian	quadratic
$\ln 2$	(0, 1, 1)	$\{0, 1\}$	C	alternating harmonic	$O(1/K)$

Offset behavior: see Proposition 6.6.

References

- [1] Richard L. Burden and J. Douglas Faires. *Numerical Analysis*. Cengage Learning, 10th edition, 2015.
- [2] Godfrey H. Hardy and Edward M. Wright. *An Introduction to the Theory of Numbers*. Oxford University Press, 5th edition, 1979.
- [3] Thomas L. Heath. *A History of Greek Mathematics, Vol. 1*. Oxford University Press, 1921.
- [4] Konrad Knopp. *Infinite Sequences and Series*. Dover, 1956.
- [5] Thomas Koshy. *Fibonacci and Lucas Numbers with Applications*. Wiley, 2001.
- [6] Gottfried Wilhelm Leibniz. De vera proportionem circuli ad quadratum circumscriptum in numeris rationalibus. *Acta Eruditorum*, 1682.
- [7] Gottfried Wilhelm Leibniz. Explication de l’arithmétique binaire. *Mémoires de l’Académie Royale des Sciences*, 1703.

- [8] Jeffrey W. Milton. Tholonia: The existential mechanics of awareness, 2024. Self-published. Available at <https://tholonia.github.io/the-book/>.
- [9] Ranjan Roy. *Sources in the Development of Mathematics: Infinite Series and Products from the Fifteenth to the Twenty-first Century*. Cambridge University Press, 2011.
- [10] Walter Rudin. *Principles of Mathematical Analysis*. McGraw-Hill, 3rd edition, 1976.
- [11] Frank J. Swetz. Leibniz, the Yijing, and the religious conversion of the Chinese. *Mathematics Magazine*, 76(4):276–291, 2003.