STATS 275: Statistical Consulting Preliminary and Formal Analysis ¹

Babak Shahbaba

Department of Statistics, UCI

Shahbaba (UCI)

Data exploration

- Data screening: assessment of data quality
 - Examining measurements
 - Examining summary statistics such as mean, standard deviation, min, max, ...
 - Identifying data outliers (we don't remove outliers unless they arise from errors that cannot be fixed)
 - Detecting patterns of missing data
 - Assessing pairwise relationships to detect redundancy
- Preliminary graphical analysis
- Preliminary tabular analysis

Model formulation

- Next, we typically specify probability models representing the underlying mechanism of the observed data
- Model parameters usually capture our scientific questions
- Types of models
 - parametric, nonparametric, or semiparametric
 - substantive (specific to the underlying subject matter) or purely empirical (typical statistical models)
- Models specifications usually decompose systematic and random variations

Formal inference

- After specifying the model, we perform formal statistical inference
- Inference typically involves making decisions: hypothesis testing, estimation, or prediction
- Two general frameworks for statistical inference are frequentist (likelihood-based) and Bayesian
- Bayesian inference involves specifying priors and obtaining posterior distributions, which are used to make inference
- Frequentist inference usually involve estimation (point or interval) and significance tests (cautions should be taken when a large number of significance tests are performed simultaneously)

Post analysis

- Model checking and diagnosis
- Tabular and graphical presentation of results
- Interpretation of results in clear, plain language, and in the subject-matter context
- Publication!