

# STATS 275: Statistical Consulting Design of Studies <sup>1</sup>

Babak Shahbaba

Department of Statistics, UCI

---

<sup>1</sup>Based on “Principles of Applied Statistics” by Cox and Donnelly <img alt="Beamer navigation icons" data-bbox="700 935 988 964"/>

# Why we design studies?

- To avoid systematic error arising from irrelevant sources that do not cancel out in the long term
- To reduce non-systematic error to a reasonable level by replication and other appropriate techniques
- To estimate realistically the extent of uncertainty in the final conclusion
- To ensure the scale of effort is appropriate; Goldilocks rule: not too limited, not too wasteful, just right!

# What is involved in a study design?

- Units of analysis (e.g., patients, patient-month)
- Deciding the type of study: observational studies vs. experiments
- Avoiding systematic error using balanced design or adjustment
- Controlling random error by replication and inclusion of background variables
- Controlling the scale of effort to achieve a certain power of significance tests or a desired value for the standard error of the estimates.

# Sampling a specific population

- Learning about a specific target population by studying a subset (sample)
  - ▶ Simple random sampling
  - ▶ Stratification
  - ▶ Use of control variables
  - ▶ Multi-stage and temporal sampling

# Experiment

- Investigating *treatment* effect by assigning *experimental units* to different treatments and observing one or more *responses*
  - ▶ Randomization
  - ▶ Forming blocks of units
  - ▶ Replication

# Observational study

- Investigators are typically passive observers (either cannot control the system or choose not to) investigating the relationship between various features of units of analysis
  - ▶ Cross-sectional: Individuals are observed at one point in time, although some background information might also be used
  - ▶ Prospective observational studies: following a group or cohort of individuals forward in time
  - ▶ Retrospective observational studies: for each identified case, one or more controls are included in the sample, and explanatory features of all units of study are determined retrospectively