


PIZZA

SQL PROJECT


HOUSE



WELCOME TO PIZZA DAY!

NEED TO UNDERSTAND LOT MORE
ABOUT THE PIZZA BUSINESS

CEO HAS SOME
QUERY LETS REOLVE
IT.



Q1. RETRIEVE THE TOTAL NUMBER OF ORDERS PLACED.

```
select count(*) from orders;
```

	count(*)
▶	21350



Q2. CALCULATE THE TOTAL REVENUE GENERATED FROM PIZZA SALES.

```
SELECT  
    ROUND(SUM(p.price * od.quantity), 2) AS Total_sales  
FROM  
    pizzas p  
    JOIN  
    orders_details od ON p.pizza_id = od.pizza_id;
```

	Total_sales
▶	817860.05



Q3. IDENTIFY THE HIGHEST-PRICED PIZZA.

```
SELECT
    pt.name, p.price
FROM
    pizzas p
    JOIN
    pizza_types pt ON p.pizza_type_id = pt.pizza_type_id
ORDER BY price DESC
LIMIT 1;
```

	name	price
▶	The Greek Pizza	35.95



Q4. IDENTIFY THE MOST COMMON PIZZA SIZE ORDERED.

```
SELECT
    size, COUNT(*) AS most_ordered_size
FROM
    pizzas p
    RIGHT JOIN
    orders_details od ON p.pizza_id = od.pizza_id
GROUP BY size
ORDER BY most_ordered_size DESC limit 1;
```

	size	most_ordered_size
▶	L	18526



Q5. LIST THE TOP 5 MOST ORDERED PIZZA TYPES ALONG WITH THEIR QUANTITIES.

```
SELECT
    pt.name, SUM(od.quantity) as quantity
FROM
    orders_details od
    JOIN
    pizzas p ON p.pizza_id = od.pizza_id
    JOIN
    pizza_types pt ON pt.pizza_type_id = p.pizza_type_id
GROUP BY pt.name
ORDER BY 2 DESC
LIMIT 5;
```

	name	quantity
►	The Classic Deluxe Pizza	2453
	The Barbecue Chicken Pizza	2432
	The Hawaiian Pizza	2422
	The Pepperoni Pizza	2418
	The Thai Chicken Pizza	2371



Q6. JOIN THE NECESSARY TABLES TO FIND THE TOTAL QUANTITY OF EACH PIZZA CATEGORY ORDERED.

```
SELECT
    pt.category, SUM(od.quantity) AS quantity
FROM
    orders_details od
    JOIN
    pizzas p ON p.pizza_id = od.pizza_id
    JOIN
    pizza_types pt ON pt.pizza_type_id = p.pizza_type_id
GROUP BY pt.category
ORDER BY quantity DESC;
```

	category	quantity
►	Classic	14888
	Supreme	11987
	Veggie	11649
	Chicken	11050



Q7. DETERMINE THE DISTRIBUTION OF ORDERS BY HOUR OF THE DAY.

```
SELECT
    HOUR(order_time) AS Hours, COUNT(*) AS order_count
FROM
    orders
GROUP BY hours
ORDER BY order_count DESC;
```

	Hours	order_count
▶	12	2520
	13	2455
	18	2399
	17	2336
	19	2009
	16	1920
	20	1642
	14	1472
	15	1468
	11	1231
	21	1198
	22	663
	23	28
	10	8
	9	1



Q8. JOIN RELEVANT TABLES TO FIND THE CATEGORY-WISE DISTRIBUTION OF PIZZAS.

```
SELECT  
    category, COUNT(*) AS count  
FROM  
    pizza_types  
GROUP BY category;
```

	category	count
▶	Chicken	6
	Classic	8
	Supreme	9
	Veggie	9



Q9. GROUP THE ORDERS BY DATE AND CALCULATE THE AVERAGE NUMBER OF PIZZAS ORDERED PER DAY.

```
SELECT
    ROUND(AVG(qty)) Daily_Average_Order
FROM
    (SELECT
        order_date, SUM(quantity) AS qty
    FROM
        orders o
    JOIN orders_details od ON od.order_id = o.order_id
    GROUP BY order_date) AS daily_orders;
```

	Daily_Average_Order
▶	138



Q10. DETERMINE THE TOP 3 MOST ORDERED PIZZA TYPES BASED ON REVENUE.

```
SELECT
    pt.name,
    ROUND(SUM(od.quantity * p.price), 2) AS sales
FROM
    orders_details od
    JOIN
    pizzas p ON p.pizza_id = od.pizza_id
    JOIN
    pizza_types pt ON pt.pizza_type_id = p.pizza_type_id
GROUP BY pt.name
ORDER BY sales DESC
LIMIT 3;
```

	name	sales
▶	The Thai Chicken Pizza	43434.25
	The Barbecue Chicken Pizza	42768
	The California Chicken Pizza	41409.5



Q11. CALCULATE THE PERCENTAGE CONTRIBUTION OF EACH PIZZA TYPE TO TOTAL REVENUE.

```
with helper as
(
select pt.category, sum(od.quantity * p.price) as revenue from orders_details od
join pizzas p on p.pizza_id = od.pizza_id
join pizza_types pt on pt.pizza_type_id = p.pizza_type_id
group by pt.category)

select category, round(revenue *100 / (select sum(revenue) from helper),2) as Revenue_Share
from helper order by revenue_share desc;
```

	category	Revenue_Share
▶	Classic	26.91
	Supreme	25.46
	Chicken	23.96
	Veggie	23.68



Q12. ANALYZE THE CUMULATIVE REVENUE GENERATED OVER TIME.

```
select * , sum(revenue) over(order by order_date) as cum_revenue from
(select o.order_date, round(sum(od.quantity * p.price),2) as revenue
  from orders_details od
 join orders o on o.order_id = od.order_id
 join pizzas p on p.pizza_id = od.pizza_id
 group by order_date
 order by order_date) as sales;
```

	order_date	revenue	cum_revenue
▶	2015-01-01	2713.85	2713.85
	2015-01-02	2731.9	5445.75
	2015-01-03	2662.4	8108.15
	2015-01-04	1755.45	9863.6
	2015-01-05	2065.95	11929.55
	2015-01-06	2428.95	14358.5
	2015-01-07	2202.2	16560.7
	2015-01-08	2838.35	19399.05
	2015-01-09	2127.35	21526.399999999998
	2015-01-10	2463.95	23990.35
	2015-01-11	1872.3	25862.649999999998
	2015-01-12	1919.05	27781.699999999997
	2015-01-13	2049.6	29831.299999999996
	2015-01-14	2527.4	32358.699999999997
	2015-01-15	1984.8	34343.5



Q13. DETERMINE THE TOP 3 MOST ORDERED PIZZA TYPES BASED ON REVENUE FOR EACH PIZZA CATEGORY.

```
with helper as (  
  select pt.category, pt.name, sum(od.quantity * p.price) as revenue  
  from orders_details od join pizzas p on p.pizza_id = od.pizza_id  
  join pizza_types pt on pt.pizza_type_id = p.pizza_type_id  
  group by pt.category, pt.name  
  order by revenue desc)  
  
select * from  
(select *,  
 rank() over (partition by category order by revenue desc) as rn from helper )  
as rn_table  
where rn <= 3;
```

category	name	revenue	rn
Chicken	The Thai Chicken Pizza	43434.25	1
Chicken	The Barbecue Chicken Pizza	42768	2
Chicken	The California Chicken Pizza	41404.25	3
Classic	The Classic Deluxe Pizza	38180.5	1
Classic	The Hawaiian Pizza	32273.25	2
Classic	The Pepperoni Pizza	30161.75	3
Supreme	The Spicy Italian Pizza	34831.25	1
Supreme	The Italian Supreme Pizza	33476.75	2
Supreme	The Sicilian Pizza	30940.5	3
Veggie	The Four Cheese Pizza	32265.700000000065	1
Veggie	The Mexicana Pizza	26780.75	2
Veggie	The Five Cheese Pizza	26066.5	3





THANK YOU
AND ENJOY
A SLICE OF
PIZZA!