

# Introduction to Abstract Algebra I: Homework 2

Due on October 15, 2025 at 23:59

*Victor Ostrik*

**Hashem A. Damrah**

UO ID: 952102243



**Exercise 2.10.** Let  $n$  be a positive integer and let  $n\mathbb{Z} = \{nm \mid m \in \mathbb{Z}\}$ .

- (i) Show that  $\langle n\mathbb{Z}, + \rangle$  is a group.
- (ii) Show that  $\langle n\mathbb{Z}, + \rangle \cong \langle \mathbb{Z}, + \rangle$ .

*Solution to (i).* We verify the group axioms for  $\langle n\mathbb{Z}, + \rangle$ .

Let  $a, b \in n\mathbb{Z}$ . Then  $a = nm$  and  $b = nk$  for some  $m, k \in \mathbb{Z}$ . Hence,

$$a + b = nm + nk = n(m + k).$$

Since  $m + k \in \mathbb{Z}$ , it follows that  $a + b \in n\mathbb{Z}$ .

Addition in  $\mathbb{Z}$  is associative, and  $n\mathbb{Z}$  inherits this property. Thus, for all  $a, b, c \in n\mathbb{Z}$ ,

$$(a + b) + c = a + (b + c).$$

The additive identity is 0, since for any  $a \in n\mathbb{Z}$ ,

$$a + 0 = a.$$

For any  $a = nm \in n\mathbb{Z}$ , the additive inverse is  $-a = n(-m)$ , because

$$a + (-a) = nm + n(-m) = n(m - m) = 0.$$

Therefore, all group axioms are satisfied, and  $\langle n\mathbb{Z}, + \rangle$  is a group. □

*Solution to (ii).* Define a map  $\varphi : \mathbb{Z} \rightarrow n\mathbb{Z}$  by  $\varphi(m) = nm$ , for all  $m \in \mathbb{Z}$ . We show that  $\varphi$  is an isomorphism.

For any  $m_1, m_2 \in \mathbb{Z}$ ,

$$\varphi(m_1 + m_2) = n(m_1 + m_2) = nm_1 + nm_2 = \varphi(m_1) + \varphi(m_2).$$

Hence,  $\varphi$  preserves addition.

Suppose  $\varphi(m_1) = \varphi(m_2)$ . Then  $nm_1 = nm_2$ , and since  $n \neq 0$ , we may divide by  $n$  to obtain  $m_1 = m_2$ . Thus,  $\varphi$  is injective.

For any  $a \in n\mathbb{Z}$ , there exists  $k \in \mathbb{Z}$  such that  $a = nk$ . Then  $\varphi(k) = nk = a$ , so  $\varphi$  is surjective.

Since  $\varphi$  is an isomorphism, it follows that  $\langle n\mathbb{Z}, + \rangle \cong \langle \mathbb{Z}, + \rangle$ . □

**Exercise 2.19.** Let  $S$  be the set of all real numbers except  $-1$ . Define  $*$  on  $S$  by

$$a * b = a + b + ab.$$

- (i) Show that  $*$  gives a binary operation on  $S$ .
- (ii) Show that  $\langle S, * \rangle$  is a group.
- (iii) Find the solution of the equation  $2 * x * 3 = 7$  in  $S$ .

*Solution to (i).* We need to show that for any  $a, b \in S$ , the result of the operation  $a * b$  is also in  $S$ .

Let  $a, b \in S$ . Then,  $a * b = a + b + ab$ . We need to check that  $a * b \neq -1$ . Suppose for contradiction that  $a * b = -1$ . Then,  $a + b + ab = -1$ . Rearranging gives  $ab + a + b + 1 = 0$ , which can be factored as  $(a + 1)(b + 1) = 0$ .

Since  $a, b \in S$ , we have  $a \neq -1$  and  $b \neq -1$ , so neither factor can be zero. This is a contradiction. Therefore,  $a * b \neq -1$ , and thus  $a * b \in S$ .

Hence,  $*$  is a binary operation on  $S$ . □

*Solution to (ii).* We verify the group axioms for  $\langle S, * \rangle$ .

As shown in part (i), for any  $a, b \in S$ ,  $a * b \in S$ .

For any  $a, b, c \in S$ ,

$$(a * b) * c = (a + b + ab) * c = (a + b + ab) + c + (a + b + ab)c.$$

Expanding this gives

$$a + b + ab + c + ac + bc + abc.$$

Similarly,

$$a * (b * c) = a * (b + c + bc) = a + (b + c + bc) + a(b + c + bc).$$

Expanding this also gives

$$a + b + c + bc + ab + ac + abc.$$

Since both expressions are equal,  $*$  is associative.

We need to find an element  $e \in S$  such that for all  $a \in S$ ,  $a * e = e * a = a$ . Let  $e = 0$ . Then,

$$a * 0 = a + 0 + a \cdot 0 = a,$$

and

$$0 * a = 0 + a + 0 \cdot a = a.$$

Thus, 0 is the identity element.

Fix  $a \in S$ . We want to find a  $b \in S$  with  $a * b = 0$ . Solving the equation  $a + b + ab = 0$ , we get  $(a + 1)(b + 1) = 1$ . Thus  $b + 1 = 1/(a + 1)$ , so

$$b = \frac{1}{a + 1} - 1 = -\frac{a}{a + 1}.$$

Since  $a \neq -1$  the denominator  $a + 1 \neq 0$ , so this  $b$  is well-defined and satisfies  $b \neq -1$ . Notice that  $b \neq -1$ , since that would imply that  $-a/(a + 1) = -1$ , which gives us  $0 = 1$ , which is a contradiction. Hence every element has a two-sided inverse in  $S$ .

Since all group axioms are satisfied,  $\langle S, * \rangle$  is a group.  $\square$

*Solution to (iii).* We need to solve the equation  $2 * x * 3 = 7$  in  $S$ . First, we compute  $2 * x$ :

$$2 * x = 2 + x + 2x = x + 2 + 2x = 3x + 2.$$

Next, we compute  $(2 * x) * 3$ :

$$(3x + 2) * 3 = (3x + 2) + 3 + (3x + 2) \cdot 3 = 3x + 2 + 3 + 9x + 6 = 12x + 11.$$

We set this equal to 7:

$$12x + 11 = 7.$$

Solving for  $x$ , we get

$$12x = -4 \Rightarrow x = -\frac{1}{3}.$$

Since  $-1/3 \in S$ , the solution is  $x = -1/3$ .  $\square$

**Exercise 2.28.** An element  $a \neq e$  in a group is said to have order 2 if  $a * a = e$ . Prove that if  $G$  is a group and  $a \in G$  has order 2, then for any  $b \in G$ ,  $b' * a * b$  also has order 2.

*Solution.* Assume  $a \neq e$  is an element of order 2 in the group  $G$ , i.e.,  $a * a = e$ . Then, for any  $b \in G$ , we compute the square of the element  $b' * a * b$ :

$$(b' * a * b) * (b' * a * b) = b' * a * (b * b') * a * b = b' * a * a * b = b' * b = e.$$

Thus,  $(b' * a * b)$  has order 2.  $\square$

**Exercise 2.29.** Show that if  $G$  is a finite group with identity  $e$  and with an even number of elements, then there is  $a \neq e$  in  $G$  such that  $a * a = e$ .

*Solution.* Let  $G$  be a finite group with identity  $e$  and an even number of elements. Consider the set  $G \setminus \{e\}$ , which has an odd number of elements.

For each element  $a \in G \setminus \{e\}$ , consider its inverse  $a'$ . If  $a \neq a'$ , then we can pair  $a$  with  $a'$ . Each such pair contributes two distinct elements to the set. Since the total number of elements in  $G \setminus \{e\}$  is odd, there must be at least one element that is its own inverse, i.e., an element  $a$  such that  $a = a'$ .

For this element, we have  $a * a = e$ . Since  $a \neq e$ , we have found an element in  $G$  such that  $a * a = e$ .  $\square$

**Exercise 2.30.** Let  $\mathbb{R}^*$  be the set of all real numbers except 0. Define  $*$  on  $\mathbb{R}^*$  by letting  $a * b = |a|b$ .

- (i) Show that  $*$  gives an associative binary operation on  $\mathbb{R}^*$ .
- (ii) Show that there is a left identity for  $*$  and a right inverse for each element in  $\mathbb{R}^*$ .
- (iii) Is  $\mathbb{R}^*$  with this binary operation a group?
- (iv) Explain the significance of this exercise.

*Solution to (i).* We need to show that for any  $a, b \in \mathbb{R}^*$ , the result of the operation  $a * b$  is also in  $\mathbb{R}^*$ .

Let  $a, b \in \mathbb{R}^*$ . Then,  $a * b = |a|b$ . Since  $|a| > 0$  and  $b \neq 0$ , it follows that  $|a|b \neq 0$ . Thus,  $a * b \in \mathbb{R}^*$ .

Next, we verify associativity. For any  $a, b, c \in \mathbb{R}^*$ ,

$$(a * b) * c = (|a|b) * c = ||a|b|c = |a||b|c.$$

Similarly,

$$a * (b * c) = a * (|b|c) = |a|(|b|c) = |a||b|c.$$

Since both expressions are equal,  $*$  is associative.

Therefore,  $*$  is an associative binary operation on  $\mathbb{R}^*$ .  $\square$

*Solution to (ii).* We need to find a left identity element  $e \in \mathbb{R}^*$  such that for all  $a \in \mathbb{R}^*$ ,  $e * a = a$ . Let  $e = 1$ . Then,

$$e * a = 1 * a = |1|a = a.$$

Thus, 1 is a left identity.

Next, we need to find a right inverse for each element  $a \in \mathbb{R}^*$ . We want to find  $b \in \mathbb{R}^*$  such that  $a * b = e$ . Using the left identity found above, we have:

$$a * b = |a|b = 1.$$

Solving for  $b$ , we get:

$$b = \frac{1}{|a|}.$$

Since  $|a| > 0$ , it follows that  $b \neq 0$ , and thus  $b \in \mathbb{R}^*$ . Therefore, every element in  $\mathbb{R}^*$  has a right inverse.

Hence, there is a left identity and a right inverse for each element in  $\mathbb{R}^*$ .  $\square$

*Solution to (iii).* To determine if  $\mathbb{R}^*$  with the operation  $*$  is a group, we need to check if it satisfies all group axioms.

We have already shown that  $*$  is an associative binary operation on  $\mathbb{R}^*$ , and that there is a left identity element (1) and a right inverse for each element.

However, we need to check if the left identity is also a right identity. For any  $a \in \mathbb{R}^*$ ,

$$a * e = a * 1 = |a| \cdot 1 = |a|.$$

Since  $|a|$  is not necessarily equal to  $a$  (for example, if  $a = -2$ , then  $|a| = 2$ ), the left identity is not a right identity.

Therefore,  $\mathbb{R}^*$  with this binary operation does not satisfy all the group axioms, and hence it is not a group.  $\square$

*Solution to (iv).* The significance of this exercise is to illustrate that having a left identity and right inverses for each element does not guarantee that a set with a binary operation forms a group. The failure of the left identity to also be a right identity shows that the group axioms are not fully satisfied. This shows the importance of verifying all group properties when determining if a structure is indeed a group.  $\square$

**Exercise 2.31.** If  $*$  is a binary operation on a set  $S$ , an element  $x$  of  $S$  is an *idempotent* for  $*$  if  $x * x = x$ . Prove that a group has exactly one idempotent element. (You may use any theorems proved so far in the text.)

*Solution.* Let  $G$  be a group with identity element  $e$ . We first show that  $e$  is an idempotent element. Obviously,  $e * e = e$ .

Now, suppose there is another idempotent element  $x \in G$  such that  $x * x = x$ . We will show that  $x = e$ . Since  $G$  is a group, every element has an inverse. Let  $x'$  be the inverse of  $x$ . Then,

$$x' * (x * x) = x' * x.$$

Using associativity, we have  $(x' * x) * x = e * x = x$ . But since  $x' * x = e$ , we have  $e * x = x$ . Therefore,  $x = e$ .

Thus, the only idempotent element in the group  $G$  is the identity element  $e$ . Hence, a group has exactly one idempotent element.  $\square$

**Exercise 2.32.** Show that every group  $G$  with identity  $e$  and such that  $x * x = e$  for all  $x \in G$  is abelian. [Hint: Consider  $(a * b) * (a * b)$ .]

*Solution.* Let  $a, b \in G$ . Since  $x * x = e$  for all  $x \in G$ , each element is its own inverse; i.e.  $x' = x$  for every  $x \in G$ .

Consider  $(a * b) * (a * b)$ . By hypothesis  $(a * b) * (a * b) = e$ , so

$$(a * b)' = (a * b).$$

On the other hand, the general inverse formula in any group gives  $(a * b)' = b' * a'$ . Using  $x' = x$  for  $a$  and  $b$  we obtain

$$(a * b)' = b' * a' = b * a.$$

Combining the two expressions for  $(a * b)'$  yields  $a * b = b * a$ . Thus,  $G$  is abelian.  $\square$

**Exercise 2.33.** Let  $G$  be an abelian group and let  $c^n = c * c * \cdots * c$  for  $n$  factors  $c$ , where  $c \in G$  and  $n \in \mathbb{Z}^+$ . Give a mathematical induction proof that  $(a * b)^n = (a^n) * (b^n)$  for all  $a, b \in G$ .

*Solution.* We will prove by induction on  $n$  that for all  $a, b \in G$ ,  $(a * b)^n = (a^n) * (b^n)$ .

For the base case, we have  $n = 1$ , which gives us

$$(a * b)^1 = a * b.$$

Also,

$$a^1 * b^1 = a * b.$$

Thus, the base case holds.

Assume that the statement holds for some  $k \in \mathbb{Z}^+$ , i.e., assume that

$$(a * b)^k = (a^k) * (b^k).$$

We need to show that it holds for  $k + 1$ . Consider:

$$(a * b)^{k+1} = (a * b)^k * (a * b).$$

By the inductive hypothesis, we have:

$$(a * b)^{k+1} = ((a^k) * (b^k)) * (a * b).$$

Since  $G$  is abelian, we can rearrange the terms:

$$(a * b)^{k+1} = (a^k * a) * (b^k * b) = a^{k+1} * b^{k+1}.$$

Thus, by the principle of mathematical induction, the statement holds for all  $n \in \mathbb{Z}^+$ .  $\square$

**Exercise 2.34.** Suppose that  $G$  is a group and  $a, b \in G$  satisfy  $a * b = b * a'$  where as usual,  $a'$  is the inverse for  $a$ . Prove that  $b * a = a' * b$ .

*Solution.* Given that  $a * b = b * a'$ , we want to show that  $b * a = a' * b$ .

Starting from the given equation, we can multiply both sides on the left by  $a'$ :

$$a' * (a * b) = a' * (b * a').$$

Using associativity, we have:

$$(a' * a) * b = a' * b * a'.$$

Simplifying, we get:

$$b = a' * b * a'.$$

Now, we can multiply both sides on the right by  $a$ :

$$b * a = (a' * b * a') * a.$$

Again, using associativity, we have:

$$b * a = a' * b * (a' * a).$$

Since  $a' * a = e$ , this simplifies to:

$$b * a = a' * b.$$

Thus, we have shown that  $b * a = a' * b$ .  $\square$

**Exercise 2.36.** Let  $G$  be a group with a finite number of elements. Show that for any  $a \in G$ , there exists  $n \in \mathbb{Z}^+$  such that  $a^n = e$ . See Exercise 33 for the meaning of  $a^n$ . [Hint: Consider  $e, a, a^2, a^3, \dots, a^m$ , where  $m$  is the number of elements in  $G$ , and use the cancellation laws.]

*Solution.* Let  $G$  be a finite group with  $m$  elements. Consider the sequence of elements:

$$e, a, a^2, a^3, \dots, a^m.$$

Since  $G$  has only  $m$  elements, by the pigeonhole principle, at least two of these elements must be equal. Thus, there exist integers  $i$  and  $j$  with  $0 \leq i < j \leq m$  such that:

$$a^i = a^j.$$

Using the cancellation law (which holds in groups), we can multiply both sides on the left by  $(a^i)'$ , the inverse of  $a^i$ , to obtain:

$$e = a^{j-i}.$$

Letting  $n = j - i$ , we have found a positive integer  $n$  such that  $a^n = e$ . Thus, for any element  $a \in G$ , there exists  $n \in \mathbb{Z}^+$  such that  $a^n = e$ .  $\square$