

Exercise 8: Numerical Integration

MAD

bacdavid@student.ethz.ch

Outline

1. Information
2. Goals
3. Theory / Recap (55')
4. Exercises (5')

Information

General

- Slides by LAB are on the website (look at them before proceeding!)
- These slides: <https://polybox.ethz.ch/index.php/s/9NFCvtriBbRBDnS>
- **No exercises next week (1. Mai), but lecture!**

Goals

Goals of Today

- Understand why we use numerical integration
- Understand the basic approach for numerical integration
- Know how to integrate numerically with Rectangle, Trapezoidal, & Simpson's rule
- Understand derivation of Newton-Cotes formula & how to use it

Theory / Recap

Integral


- Exact Integral:

$$I = \int_a^b f(x) dx$$

- ...can be solved analytically or not

- Used for differential equations:

$$\dot{x} = f(x)$$



https://de.wikipedia.org/wiki/Integralrechnung#/media/File:Integral_as_region_under_curve.svg

Exercise 1

- Solve the ODE:

$$\dot{x} = t \cdot e^t$$

- Tipp:

$$\int_a^b uv' dx = [uv]_a^b - \int_a^b u' v dx$$

Numerical Integration

- Approximate:

$$I = \int_a^b f(x) dx = \sum_{i=0}^{N-1} I_i$$


- Approximate I_i : (table)

- Rectangular Rule
- Trapezoidal Rule
- Simpson's Rule

Approximations for I_i


Rectangular Rule	$I_{Ri} = f(x_{i+1/2}) \cdot \Delta_i$
Trapezoidal Rule	$I_{Ti} = \frac{f(x_i) + f(x_{i+1})}{2} \cdot \Delta_i$
Simpson's Rule	$I_{Si} = \frac{f(x_i) + 4f(x_{i+1/2}) + f(x_{i+1})}{6} \cdot \Delta_i$

Rectangular Rule




<http://tutorial.math.lamar.edu/Classes/CalcI/AreaProblem.aspx>

Trapezoidal Rule



<http://tutorial.math.lamar.edu/Classes/CalcI/AreaProblem.aspx>

Simpson's Rule



https://de.wikipedia.org/wiki/Datei:Simpsons_method_illustration.svg

Exercise 2

- Solve the integral

$$\int_{-1}^1 x^2 dx$$

1. Exact
2. Rectangle Rule
3. Trapezoidal Rule
4. Simpson's Rule

Approximations for I_i

Rectangular Rule	$I_{Ri} = f(x_{i+1/2}) \cdot \Delta_i$
Trapezoidal Rule	$I_{Ti} = \frac{f(x_i) + f(x_{i+1})}{2} \cdot \Delta_i$
Simpson's Rule	$I_{Si} = \frac{f(x_i) + 4f(x_{i+1/2}) + f(x_{i+1})}{6} \cdot \Delta_i$

Newton-Cotes Formula

- Approximate:

$$I \approx (b - a) \cdot \sum_{k=0}^n C_k^n f(x_k), \quad \text{with } C_k^n = \frac{1}{b - a} \int_a^b l_k^n(x) dx \quad \text{„weights“}$$

- Properties of C_k^n :

- $\sum_{k=0}^n C_k^n = 1$
- $C_k^n = C_{n-k}^n$

- Note:

- Can be used for entire integral I or for smaller intervals I_i
- The “weights” C_k^n are used supporting points f_k

Exercise 3

- use $n = 2$ and equally spaced x_0, x_1, x_2 to approximate I

$$I \approx (b - a) \cdot \sum_{k=0}^n C_k^n f(x_k),$$

with $C_k^n = \frac{1}{b - a} \int_a^b l_k^n(x) dx$ „weights“

- Tipp: $C_1^2 = \frac{2}{3}$

Exercises

Exercises

1. Use Newton-Cotes to derive Simpson's Rule (no short-cuts)
2. Implement Simpson's & Trapezoidal Integration (coding)

Questions?

