

Title

- Text

Exercise 10: Numerical Integration III

MAD

bacdavid@student.ethz.ch

Outline

1. Information
2. Goals
3. Theory/ Recap
4. Exercises

Information

General

- Lecture material & problem sets available [here](#)
- Tutorial material available [here](#)

Goals


Goals of Today

- Recap Richardson's Extrapolation
- Understand how Richardson's Extrapolation can be used to estimate an error
- Understand recursion in programming
- Understand Adaptive Quadrature
- Be able to derive Gauss Quadrature
- Be able to use Gauss Quadrature with tables

Theory / Recap

Adaptive Quadrature

- Use adaptive stepsize
- Decrease stepsize in specific region until desired accuracy is achieved



<https://www.cs.uic.edu/~hogand/cs107/notes-numerical.html>

Reminder: Error Estimate with Richardson's Extrapolation

- We can write a Taylor Series expansion of a numerical method:

$$G(h) = G(0) + c_1 h + c_2 h^2 + \dots$$

- Hence the error generated by the stepsize $h/2$:

$$\epsilon(h/2) = |G(0) - G(h/2)| = \left| -\frac{c_1}{2} h - \frac{c_2}{4} h^2 + \dots \right|$$

- On the other hand:

$$G(h/2) - G(h) = -\frac{c_1}{2} h - \frac{3c_2}{4} h^2 + \dots$$

- Hence:

$$\epsilon(h/2) \approx |G(h/2) - G(h)|$$

Example 1: Error estimate

- Approximate the error $\epsilon\left(\frac{h}{2}\right)$ of the Euler Forward method and compare it to the true error:


$$f'(x) \approx \frac{f(x+h) - f(x)}{h}$$

- Given:
 - $x = 0$
 - $h = 0.2$
 - $f(x) = x^2$
- Remember:
 - $\epsilon(h/2) \approx |G(h/2) - G(h)|$

Reminder: Recursion

- The function calls itself

```
function f(x)
  if base case then
    Return base value
  else
    Return recursive value
  end if
end function
```



<https://stackoverflow.com/questions/8183426/factorial-using-recursion-in-java>

Example 2: Recursion for gcd

- Write pseudo-code for a function that finds the greatest common divisor, using Euclid's algorithm:

$$\text{gcd}(x, y) = \begin{cases} x & \text{if } y = 0 \\ \text{gcd}(y, x \bmod y) & \text{if } y > 0 \end{cases}$$

- Calculate the gcd of 8 and 12 with the algorithms, draw a diagram emphasizing the calls to the functions and the return values

Solution 2

```
function gcd(x, y)
    if y is 0 then
        Return x
    else
        Return gcd(y, x mod y)
    end if
end function
```

Adaptive Quadrature Algorithm

```
function ADAPTIVESIMPSON(a, b)
    Apply Simpson's rule in [a, b]
    Split interval into [a, m] & [m, b]; with  $m = (a + b)/2$ 
    Apply Simpson's rule on [a, m] & [m, b]
    Estimate Error with Richardson's extrapolation

    if error  $> \epsilon$  then
        Return ADAPTIVESIMPSON(a, m) + ADAPTIVESIMPSON(m, b)
    else
        Return value of Simpson's rule (accurate one)
    end if
end function
```

Gauss Quadrature

- n weights + n abscissas = $2n$ parameters
- Integrate $P_{2n-1}(x)$ exactly (integral increases order by one):

$$\int_a^b f(x) dx \approx \int_a^b P_{2n-1}(x) dx = \sum_{i=1}^n \alpha_i \cdot f(x_i)$$

Note

- Transform integral to be in the interval $[-1, 1]$ in order to use lookup-tables:

$$z = \frac{2x - (a + b)}{b - a}$$

- We then have:

$$\int_{-1}^1 f(z) dz \approx \sum_{i=1}^n \alpha_i \cdot f(z_i)$$

- You can now look up z_i and α_i in tables
- Method is exact for polynomials of degree $2n - 1$

Example 3: Gauss Quadrature with lookup tables

- Find – by Gauss Quadrature - the exact integral value:

$$I = \int_0^2 x^3 dx$$

- Tipps:
 - Exact for order $2n - 1$
 - $z = \frac{2x-(a+b)}{b-a}$

n=1	x_i	α_i
1	0	2
n=2	x_i	α_i
1	$-\sqrt{\frac{1}{3}} \approx -0,57735026919$	1
2	$\sqrt{\frac{1}{3}} \approx 0,57735026919$	1
n=3	x_i	α_i
1	$-\sqrt{\frac{3}{5}} \approx -0,774596669241$	$\frac{5}{9} \approx 0,555555555556$
2	0	$\frac{8}{9} \approx 0,888888888889$
3	$\sqrt{\frac{3}{5}} \approx 0,774596669241$	$\frac{5}{9} \approx 0,555555555556$
n=4	x_i	α_i
1	$-\sqrt{\frac{3}{7} + \frac{2}{7}\sqrt{\frac{6}{5}}} \approx -0,861136311594053$	$\frac{18-\sqrt{30}}{36} \approx 0,347854845137454$
2	$-\sqrt{\frac{3}{7} - \frac{2}{7}\sqrt{\frac{6}{5}}} \approx -0,339981043584856$	$\frac{18+\sqrt{30}}{36} \approx 0,652145154862546$
3	$\sqrt{\frac{3}{7} - \frac{2}{7}\sqrt{\frac{6}{5}}} \approx 0,339981043584856$	$\frac{18+\sqrt{30}}{36} \approx 0,652145154862546$
4	$\sqrt{\frac{3}{7} + \frac{2}{7}\sqrt{\frac{6}{5}}} \approx 0,861136311594053$	$\frac{18-\sqrt{30}}{36} \approx 0,347854845137454$

Exercises

Exercise 1

- Use AQ to integrate „batman function“

Exercise 2

- Use various integration schemes – think why some work better than others!

Questions?

