

THE LIFE AND MATHEMATICS OF JAMES H. BRAMBLE

STUDENTS OF BRAMBLE

1. NEWS AND FAMILY

James Henry Bramble, one of the most distinguished computational mathematicians of all times and our beloved mentor and friend, died July 20, 2021 at his home in Austin, Texas. He was 90.

Bramble was born on December 1, 1930 in Annapolis, Maryland to Edith and Clinton Bramble. He had two sisters, Mary Aller and Barbara Lawrence, both deceased. In 1978, Bramble married Margaret (Peggy) Hays, (she passed away March 28, 2019). He is survived by his four children, Margot Dermody of Pittsburgh, Pennsylvania, Tamara Lamenzo of Needham, Massachusetts, Mitzi Bramble of Harwich, Massachusetts, and James Bramble of Austin, his step-son, Alan Hays of Ithaca, and eight grand-children. He also is survived by his first wife, Mary Eppie Boze.

He earned his bachelor degree from Brown University in 1953, a Master's degree and Ph.D. degree from the University of Maryland in 1955 and 1958 respectively. After working for General Electric and the Naval Ordnance Laboratory, he became a professor at the University of Maryland. In 1968, he was recruited to Cornell.

After retiring from Cornell in 1994 and then retiring from Texas A&M in 2007, Bramble and his wife, Peggy, remained in Austin, but returned to their lakeside cottage in Ithaca every summer. They enjoyed entertaining friends and family, playing golf and tennis, and boating.

He will be remembered fondly by family, friends, and colleagues worldwide.

1.1. Cornell days. In his 26 years at Cornell, as professor of Mathematics in the College of Arts and Sciences, Bramble helped move his department to the forefront of applied mathematics and became a leading figure in his field, bridging rigorous theory and practical numerical computation.

"Cornell hired Professor Bramble to help establish a world-class group in numerical analysis," said Tara Holm, professor of mathematics and department chair. "That group began with his adviser, Larry Payne, and later included professors Lars Wahlbin and Al Schatz. Bramble was one of the top three in his field in the country and of worldwide renown. He put applied mathematics at Cornell on the world map, as it were. Bramble's work remains central in applied mathematics and continues to be highly cited today."

Bramble also made a significant impact with Cornell's Center for Applied Mathematics. He served as its director from 1975-81, helping to secure university funding for graduate students. In addition, Bramble served as associate chair and director of undergraduate studies in the Department of Mathematics for three years, and as graduate faculty representative in mathematics for three years, Holm said.

1.2. Texas A&M. In 1994, Bramble retired from Cornell as professor emeritus. He continued his career at Texas A&M University, where he retired in 2007 as Distinguished Professor Emeritus.

He was an editor of the journal *Mathematics of Computation* for 25 years, and chief editor from 1975-83. He received an honorary doctorate from Chalmers University of Technology in Sweden in 1985 in recognition of his important contributions to mathematics.

In 1970, along with Ivo Babuška and Bruce Kellog, Bramble co-founded the Finite Element Circus (<https://sites.google.com/view/fecircus>), a regular meeting devoted to the theory and applications of the finite element method and related areas of numerical analysis and partial differential equations. In the early years, meetings were held as often as four times a year, but soon a format was established of a one-and-one-half day meeting held every spring and fall at varying locations. The Finite Element Circus has played a critical role in making the finite element method one of the major numerical methods for solving partial differential equations, and in training generations of researchers in the finite element method.

1.3. Ph.D. students of Bramble. Bramble taught numerous undergraduate and graduate students, mentored 22 Ph.D. students, and had research collaborators throughout the world. Here is a list:

- (1) [Maxine Rockoff](#)(University of Pennsylvania, 1964), currently an adjunct associate research scientist at Columbia University.
- (2) [Karl Gustafson](#)(University of Pennsylvania, 1965), currently a professor of mathematics at University of Colorado Boulder.
- (3) [Richard Bruce Simpson](#)(University of Maryland, 1966), used to be a professor of computer science at university of Waterloo, passed away in 2020.
- (4) [James Kuttler](#)(University of Maryland, 1967), currently an instructor in JHU Whiting School of Engineering.
- (5) [Stephan Hilbert](#)(University of Maryland, 1969), used to be a professor of mathematics at Ithaca College.
- (6) [Richard Falk](#)(Cornell University, 1971), currently a professor of mathematics at Rutgers.
- (7) Thomas King(Cornell University, 1971).
- (8) [Steven Serbin](#)(Cornell University, 1971), used to be a professor of mathematics at University of Tennessee, passed away in 2008.

- (9) [Garth Baker](#)(Cornell University, 1973), used to be a lecturer of mathematics at Bermuda College.
- (10) [John Trangenstein](#)(Cornell University, 1975), currently a professor of mathematics at Duke University.
- (11) [Joseph E. Pasciak](#)(Cornell University, 1977), currently a professor of mathematics at Texas A&M University.
- (12) [Ragnar Winther](#)(Cornell University, 1977), currently a professor of mathematics at University of OSLO.
- (13) Peter Sammon(Cornell University, 1978).
- (14) [Mark J. Pekker](#)(Cornell University, 1982), currently a professor of mathematics at The University of Alabama in Huntsville.
- (15) [Jinchao Xu](#)(Cornell University, 1989), currently a professor of mathematics at Pennsylvania State University.
- (16) Ping Lee(Cornell University, 1990).
- (17) Mark Hanisch(Cornell University, 1991).
- (18) [Tong Sun](#)(Texas A&M, 1997), currently a professor of mathematics at Bowling Green State University.
- (19) [Jay Gopalakrishnan](#)(Texas A&M, 1999), currently a professor of mathematics at Portland State University.
- (20) [Constantin Bacuta](#)(Texas A&M, 2000), currently a professor of mathematics at University of Delaware.
- (21) [Tzanio Kolev](#)(Texas A&M, 2004), currently a computational mathematician at Lawrence Livermore National Laboratory.
- (22) [Dimitar Trenev](#) (Texas A&M, 2009), currently working at ExxonMobil Corporate Strategic Research.

2. SCIENTIFIC CONTRIBUTIONS

Bramble began his research career developing analytical methods for partial differential equations – that is, equations that relate functions and their derivatives. In most of his career, he worked on numerical solutions of partial differential equations and multigrid method. His early work involved the mathematical analysis of finite difference methods for elliptic problems. His later work was more focused on the numerical analysis of finite element methods. He also worked on questions concerning rapid solution of large-scale systems that result from such approximations. A typical question is: Among all the theoretically good approximations to a general class of problems, are there some that can be solved efficiently by taking advantage of modern computer architectures such as parallelism? Answers to questions like this one can bring many problems into the realm of practical feasibility. He designed approximations to solutions to problems in partial differential equations that adequately describe the problem and that can be efficiently solved using modern computing power. In particular, he was interested in Maxwell's Equations and acoustics, including scattering problems.

A partial list of his major research interests include:

- (1) Finite difference methods
- (2) Finite element method
 - (a) Bramble-Hilbert Lemma and error analysis
 - (b) Super-convergence
- (3) Domain decomposition methods
 - (a) Non-overlapping domain decomposition method
 - (b) Overlapping domain decomposition method
- (4) Multigrid methods
- (5) PML methods

2.1. Finite difference method. Dr. Bramble worked on the mathematical analysis of numerical methods for partial differential equations. In the 1960s, he helped pioneer the mathematical analysis of finite difference methods for elliptical problems.

In a sequence of papers from 1962 to 1966, [22, 8, 12, 24, 23, 9, 21], Bramble and collaborators had systematical studies on error analysis for finite difference methods for elliptic boundary value problems of both 2nd and 4th order with various different (Dirichlet, Neumann and mixed) boundary conditions. For example, by using the discrete maximum principle, he proved that the finite difference schemes for domains with curved boundaries still have $O(h^2)$ accuracy, despite the fact the truncation error is only of order (h) near the boundary. As another example, in paper [12], he provided a special example in which the matrix is neither diagonally dominant nor of non-negative type, but for which a maximum principle still holds. This is a deep observation about the maximum principle for finite difference methods.

Finite difference method was extended and analyzed by Dr. Bramble for Dirichlet problem of Poisson's equation on a domain with curve boundary [22], and then for Neumann problem of Poisson's equation on a domain with curve boundary [24], and approximation of solutions of mixed boundary value problems for Poisson's equation by finite difference method was also considered by Dr. Bramble in [23]. Even further, Dr. Bramble considered and analyzed higher order finite difference method for bounded domain in higher dimension, such as fourth-order finite difference analogues of the Dirichlet problem for Poisson's equation in three and four dimensions were consider in [8].

A second order finite difference analog of the first biharmonic boundary value problem was also done by Dr. Bramble in [9]. Error estimates for finite difference methods for indefinite problem such as in forced vibration problems was also given by Dr. Bramble in [21]. In paper [12], Dr. Bramble showed a special example in which case the matrix is neither diagonally dominant nor of non-negative type, but for which a maximum principle still holds. This is a deep observation of the maximum principle of finite difference methods.

2.2. Finite element method. In the ensuing decades, one of the major focuses of Bramble's research was the development and analysis of the finite element method, and especially its theoretical foundations, applied to the approximation of solutions of elliptic and parabolic partial differential equations.

2.2.1. Bramble-Hilbert lemma. One of his most famous results, the Bramble-Hilbert Lemma, a theoretical tool for proving error estimates for the finite element method, was a collaborative effort with his Ph.D. student, Stephen Hilbert [11].

The Bramble-Hilbert Lemma states that for any bounded functional $F : W^{m,p}(\Omega) \mapsto \mathbb{R}$ that annihilates polynomials of degrees $m - 1$, then $F(v) \leq C|v|_{m,p,\Omega}$ for any $v \in W^{m,p}(\Omega)$. What is equally important, as demonstrated in the original paper [11], is how this Lemma is used together with a scaling argument to derive error estimates in Sobolev norms for approximation of smooth functions by finite elements. This gives a very general and very elegant way to derive error estimates for piecewise polynomial approximation, for numerical quadrature and for many other applications. It is a basic tool for finite element error analysis and used by every researcher in this field.

2.2.2. The eigenvalue problems. Together with John Osborn, Jim developed in [13] error estimates for the approximating the eigenvalues of nonself-adjoint operators. Various methods for defining the approximate eigenvalues are treated, including methods in which the functions in the subspaces of trial functions are not required to satisfy any boundary conditions. The estimates are optimal in the sense that they give the best estimate that could be expected on the basis of the approximability properties of the underlying subspaces. The work extends on previous results on obtaining lower bounds for the inverse Poincare constant and estimating first nonzero Steklov eigenvalue for second order elliptic PDEs, [19].

2.2.3. Time-stepping methods. For second order hyperbolic equations, a new class of single step fully discrete schemes is developed. The schemes are high order accurate in time, [6]. Together with Baker and Thomée in [7] convergence estimates for semidiscrete approximations for Parabolic Equations are investigated. The results include L_2 and maximum norm estimates, and superconvergence estimates. Several of the estimates are derived under weak assumptions on the initial data.

2.2.4. Post-processing and super-convergence. In [20] together with Schatz, *superconvergence* results for a class of Ritz-Galerkin methods for approximating elliptic boundary problems are developed. Convolution operators with locally supported functions are designed that are easy to compute from the Ritz-Galerkin approximation and lead to higher order of approximation of the solution in both L^2 and L^∞ norms for translation invariant grids. This convolutional operator does not depend on the specific elliptic

operator and is easily constructed. This technique of using convolution operator for translation invariant grids is reminiscent of the convolutional neural network [25] for classification of image, which was also motivated by the fact that the main feature of an image is translational invariant.

2.3. Domain decomposition methods. Domain decomposition methods can be, roughly speaking, divided into two different classes: (1) non-overlapping domain decomposition methods, and (2) overlapping domain decomposition methods. In both classes of methods, Bramble has made fundamental contributions.

2.3.1. Non-overlapping domain decomposition methods. In the sequence of 4 papers [14, 15, 16, 17], Bramble, Pasciak and Schatz gave a systematic development of basic non-overlapping domain decomposition methods, including an estimate of the condition number of the resulting preconditioned systems.

2.3.2. Overlapping domain decomposition methods. Overlapping domain decomposition methods can be traced back to Schwarz [] for the two subdomain case. In his original paper, Schwarz also provided a convergence analysis using the maximum principle (???). In ???, Lions used a variational approach to give a convergence analysis in the context of Sobolev space H^1 for the two-subdomain case.

In [18], the first optimal convergence analysis is given for the overlapping domain decomposition methods with coarse space.

research collaborators

2.4. Multigrid and multilevel methods.

2.4.1. Multigrid methods. Multigrid methods are among the most efficient iterative methods for the solution of linear systems which arise in many large scale scientific calculations. Every researcher working with the numerical solution of partial differential equations should at least be familiar with these powerful techniques. Multigrid methods can be traced back to the early seventies. An early and major advocate of multigrid method was A. Brandt. A uniform convergence analysis for a multigrid method for 2nd order elliptic boundary value problems was provided by Bank and Dupont []. A mathematical foundation of the multigrid method was set up by Hackbusch and his collaborators []. An invaluable contribution to this field is Bramble's book *Multigrid Methods* [10], which presents results concerning the rates of convergence of multigrid iterations.

2.4.2. Multilevel representation of norms and applications. Based on the general multilevel foundation used for multigrid methods and using real method of interpolation theory, together with Pasciak, Xu, and Bacuta, in a series of five papers, [3, 5, 1, 4, 2], a new approach was developed for obtaining sharp stability estimates (using fractional norms) for certain elliptic boundary value problems. In [3, 5] new interpolation results that generalizes

Kellogg's subspace interpolation theory are developed for proving regularity estimates for the Laplace and Biharmonic operators. Regularity for the Poisson equation on non-convex polygonal domains at the critical case are investigated in [1, 4]. For the plane elasticity problem on a convex polygonal domains regularity is proved with constants independent of the Lamé coefficients, [2].

REFERENCES

- [1] Constantin Bacuta, James Bramble, and Jinchao Xu. Regularity estimates for elliptic boundary value problems in besov spaces. *Mathematics of Computation*, 72(244):1577–1595, 2003.
- [2] Constantin Bacuta and James H Bramble. Regularity estimates for solutions of the equations of linear elasticity in convex plane polygonal domains. *Zeitschrift für angewandte Mathematik und Physik ZAMP*, 54(5):874–878, 2003.
- [3] Constantin Bacuta, James H Bramble, and Joseph E Pasciak. Shift theorems for the biharmonic dirichlet problem. In *Recent Progress in Computational and Applied PDEs*, pages 1–26. Springer, 2002.
- [4] Constantin Bacuta, James H Bramble, and Jinchao Xu. Regularity estimates for elliptic boundary value problems with smooth data on polygonal domains. 2003.
- [5] Constantin Bacuta, James H. Bramble, and Joseph E. Pasciak. Using finite element tools in proving shift theorems for elliptic boundary value problems. *Numerical linear algebra with applications*, 10(1-2):33–64, 2003.
- [6] Garth A Baker and James H Bramble. Semidiscrete and single step fully discrete approximations for second order hyperbolic equations. *RAIRO. Analyse numérique*, 13(2):75–100, 1979.
- [7] Garth A Baker, James H Bramble, and Vidar Thomée. Single step galerkin approximations for parabolic problems. *Mathematics of Computation*, 31(140):818–847, 1977.
- [8] James H Bramble. Fourth-order finite difference analogues of the dirichlet problem for poisson's equation in three and four dimensions. *Mathematics of Computation*, 17(83):217–222, 1963.
- [9] James H Bramble. A second order finite difference analog of the first biharmonic boundary value problem. *Numerische Mathematik*, 9(3):236–249, 1966.
- [10] James H Bramble. *Multigrid methods*. Chapman and Hall/CRC, 2019.
- [11] James H Bramble and SR Hilbert. Estimation of linear functionals on sobolev spaces with application to fourier transforms and spline interpolation. *SIAM Journal on Numerical Analysis*, 7(1):112–124, 1970.
- [12] James H Bramble and Bert E Hubbard. On a finite difference analogue of an elliptic boundary problem which is neither diagonally dominant nor of non-negative type. *Journal of Mathematics and Physics*, 43(1-4):117–132, 1964.
- [13] James H Bramble and JE Osborn. Rate of convergence estimates for nonselfadjoint eigenvalue approximations. *Mathematics of computation*, 27(123):525–549, 1973.
- [14] James H Bramble, Joseph E Pasciak, and Alfred H Schatz. The construction of preconditioners for elliptic problems by substructuring. i. *Mathematics of Computation*, 47(175):103–134, 1986.
- [15] James H Bramble, Joseph E Pasciak, and Alfred H Schatz. The construction of preconditioners for elliptic problems by substructuring. ii. *Mathematics of Computation*, 49(179):1–16, 1987.
- [16] James H Bramble, Joseph E Pasciak, and Alfred H Schatz. The construction of preconditioners for elliptic problems by substructuring. iii. *Mathematics of Computation*, 51(184):415–430, 1988.

- [17] James H Bramble, Joseph E Pasciak, and Alfred H Schatz. The construction of preconditioners for elliptic problems by substructuring. iv. *Mathematics of Computation*, 53(187):1–24, 1989.
- [18] James H Bramble, Joseph E Pasciak, Jun Ping Wang, and Jinchao Xu. Convergence estimates for product iterative methods with applications to domain decomposition. *Mathematics of Computation*, 57(195):1–21, 1991.
- [19] James H Bramble and LE Payne. Bounds in the neumann problem for second order uniformly elliptic operators. *Pacific Journal of Mathematics*, 12(3):823–833, 1962.
- [20] James H Bramble and Alfred H Schatz. Higher order local accuracy by averaging in the finite element method. *Mathematics of Computation*, 31(137):94–111, 1977.
- [21] JH Bramble. Error estimates for difference methods in forced vibration problems. *SIAM Journal on Numerical Analysis*, 3(1):1–12, 1966.
- [22] JH Bramble and BE Hubbard. On the formulation of finite difference analogues of the dirichlet problem for poisson’s equation. *Numerische Mathematik*, 4(1):313–327, 1962.
- [23] JH Bramble and BE Hubbard. Approximation of solutions of mixed boundary value problems for poisson’s equation by finite differences. *Journal of the ACM (JACM)*, 12(1):114–123, 1965.
- [24] JH Bramble and BE Hubbard. A finite difference analog of the neumann problem for Poisson’s equation. *Journal of the Society for Industrial and Applied Mathematics, Series B: Numerical Analysis*, 2(1):1–14, 1965.
- [25] Yann LeCun, Léon Bottou, Yoshua Bengio, and Patrick Haffner. Gradient-based learning applied to document recognition. *Proceedings of the IEEE*, 86(11):2278–2324, 1998.