

Monodromy conjecture for Newton non-degenerate hypersurfaces

Author: Oriol Baeza Guasch (FME, UPC)
Supervisor: Guillem Blanco Fernández (KU Leuven)
Co-supervisor: Josep Àlvarez Montaner (FME, UPC)

1 Introduction

2 Preliminaries

- Resolution of singularities
- Zeta function
- Bernstein-Sato polynomial
- Strong Monodromy Conjecture
- Periods of integrals

3 Plane curves

4 Newton non-degenerate

5 Examples

- Example 1
- Example 2
- Example 3
- Example 4

Contents

1 Introduction

2 Preliminaries

- Resolution of singularities
- Zeta function
- Bernstein-Sato polynomial
- Strong Monodromy Conjecture
- Periods of integrals

3 Plane curves

4 Newton non-degenerate

5 Examples

- Example 1
- Example 2
- Example 3
- Example 4

Introduction

Open problem in the theory of singularities, formulated by Igusa in the 70s.

Monodromy conjecture, topological version

Let $f \in \mathbb{C}[x_1, \dots, x_n]$ be non-constant. If s_0 is a pole of $Z_{\text{top}}(f, \varphi; s)$, then

- (*standard*) $e^{2\pi i \Re(s_0)}$ is a monodromy eigenvalue of $f: \mathbb{C}^n \rightarrow \mathbb{C}$ at some point of $\{f = 0\}$.
- (*strong*) s_0 is a root of the Bernstein-Sato polynomial b_f .

Introduction

Open problem in the theory of singularities, formulated by Igusa in the 70s.

Monodromy conjecture, topological version

Let $f \in \mathbb{C}[x_1, \dots, x_n]$ be non-constant. If s_0 is a pole of $Z_{\text{top}}(f, \varphi; s)$, then

- (*standard*) $e^{2\pi i \Re(s_0)}$ is a monodromy eigenvalue of $f: \mathbb{C}^n \rightarrow \mathbb{C}$ at some point of $\{f = 0\}$.
- (*strong*) s_0 is a root of the Bernstein-Sato polynomial b_f .

Known cases:

- Plane curves (Loeser '88)
- Newton non-degenerate polynomials* (Loeser '90)
- Some types of hyperplanes arrangements (Budur-Saito-Yuzvinsky '10, Walther '17, Bapat-Walters '15)
- Semi-quasihomogeneous singularities (Budur-Blanco-van der Veer '21)

Introduction

*The case of Newton non-degenerate polynomials requires some additional hypothesis on the *residue numbers*.

Theorem 5.5.1 [Loe90]

Let f be a comfortable polynomial verifying $f(0) = 0$, with Newton diagram $\Gamma(f)$, and Newton non-degenerate. Suppose that all compact faces τ_0 verify

- ① $\frac{k(\tau_0)}{N(\tau_0)} < 1$,
- ② For every face τ of codimension 1 of $\Gamma(f)$, distinct of τ_0 and having non-empty intersection with τ_0 , we have $\varepsilon(\tau_0, \tau) \notin \mathbb{Z}$.

Then, the real parts of the poles of the zeta function of f are roots of the Bernstein-Sato polynomial of f .

Introduction

*The case of Newton non-degenerate polynomials requires some additional hypothesis on the *residue numbers*.

Theorem 5.5.1 [Loe90]

Let f be a comfortable polynomial verifying $f(0) = 0$, with Newton diagram $\Gamma(f)$, and Newton non-degenerate. Suppose that all compact faces τ_0 verify

- ① $\frac{k(\tau_0)}{N(\tau_0)} < 1$,
- ② For every face τ of codimension 1 of $\Gamma(f)$, distinct of τ_0 and having non-empty intersection with τ_0 , we have $\varepsilon(\tau_0, \tau) \notin \mathbb{Z}$.

Then, the real parts of the poles of the zeta function of f are roots of the Bernstein-Sato polynomial of f .

Question

Can this condition be removed? What is the situation for the cases where it does not hold?

Contents

1 Introduction

2 Preliminaries

- Resolution of singularities
- Zeta function
- Bernstein-Sato polynomial
- Strong Monodromy Conjecture
- Periods of integrals

3 Plane curves

4 Newton non-degenerate

5 Examples

- Example 1
- Example 2
- Example 3
- Example 4

Preliminaries — Resolution of singularities

- $f: U \rightarrow \mathbb{C}$ a holomorphic function defined on an open set $U \subset \mathbb{C}^n$
- Hypersurface $X = f^{-1}(0)$

Definition (Singularity)

We define the set of singular points of X by the set

$$\text{Sing}(X) := \left\{ x \in U \mid f(x) = \frac{\partial f}{\partial x_1} = \cdots = \frac{\partial f}{\partial x_n} = 0 \right\}$$

Additionally, if $x \in \text{Sing}(X)$ is the only singularity in a small enough neighborhood $V \ni x$ we will say it is *isolated*.

Definition (Resolution)

A *resolution* of X is a proper morphism $\pi: Y \rightarrow X$ where

- 1 Y is a smooth variety.
- 2 The restriction outside the singular locus $\pi|_{Y \setminus \pi^{-1}(\text{Sing}(X))}: Y \setminus \pi^{-1}(\text{Sing}(X)) \rightarrow X \setminus \text{Sing}(X)$ is a birational isomorphism.

Additionally, we will say that the resolution is *good* if also

- 3 For every singular point $p \in \pi^{-1}(\text{Sing}(X))$, there exists an open neighborhood $U_p \subset Y$, and open $V \subset \mathbb{K}^n$ with a chart

$$\begin{aligned} y: U_p &\xrightarrow{\cong} V \\ p &\mapsto 0 \end{aligned}$$

such that $U \cap \pi^{-1}(\text{Sing}(X)) = \{y_{i_1} = \cdots = y_{i_r} = 0\}$ for certain indices $0 < i_1 < \cdots < i_r \leq n$.

Preliminaries — Resolution of singularities

Definition (Embedded resolution)

Let X be a smooth algebraic variety, $f: X \rightarrow \mathbb{K}$ a polynomial and abbreviate $S = \text{Sing}(f^{-1}(0))$ be the set of singular points on the zero set of f . An *embedded resolution* of f is a proper morphism $\pi: Y \rightarrow X$ where

- 1 Y is a smooth variety.
- 2 The restriction outside the singular locus $\pi|_{Y \setminus \pi^{-1}(S)}: Y \setminus \pi^{-1}(S) \rightarrow X \setminus S$ is a birational isomorphism.
- 3 For every singular point $p \in \pi^{-1}(S)$, there exists an open neighborhood $U_p \subset Y$, and an open $V \subset \mathbb{K}^n$ with a chart

$$\begin{aligned} y: U_p &\xrightarrow{\cong} V \\ p &\mapsto 0 \end{aligned}$$

over which $\pi^*f = u(y) y_{i_1}^{N_1} \cdots y_{i_r}^{N_r}$, with $u(0) \neq 0$ a unit, and $N_i \geq 0$ integers.

Guaranteed in characteristic zero, thanks to a result by Hironaka [Hir64].

Preliminaries — Resolution of singularities

Let $f \in \mathbb{C}[x_1, \dots, x_n]$ be a non-constant polynomial, and $\pi: X \rightarrow \mathbb{C}$ an embedded resolution of f .

- $(E_i)_{i \in J}$ the irreducible components of $\pi^{-1}(f^{-1}(0))$. Locally $E_i: \{x_i = 0\}$.
- By the local expression of π^*f , we know it vanishes with order N_j on a generic point of E_j , and we may write globally

$$\operatorname{div}(\pi^*f) = \sum_{j \in J} N_j E_j$$

Preliminaries — Resolution of singularities

Let $f \in \mathbb{C}[x_1, \dots, x_n]$ be a non-constant polynomial, and $\pi: X \rightarrow \mathbb{C}$ an embedded resolution of f .

- $(E_i)_{i \in J}$ the irreducible components of $\pi^{-1}(f^{-1}(0))$. Locally $E_i: \{x_i = 0\}$.
- By the local expression of π^*f , we know it vanishes with order N_j on a generic point of E_j , and we may write globally

$$\operatorname{div}(\pi^*f) = \sum_{j \in J} N_j E_j$$

Geometric setup

In the resolved space, the intersection of exceptional divisors is, at worst, like intersection of coordinate hyperplanes.

- Normal crossings: divisors are smooth and intersect transversely.
- Simple normal crossings: additionally, no three intersect at the same point.

Preliminaries — Resolution of singularities

- To each divisor $E_j, j \in J$ we have already associated a numerical quantity N_j representing the order of vanishing of a generic point of π^*f in E_j .
- Similarly, we define the integers k_j such that the order of vanishing of a generic point of the pullback of the standard volume form in E_j is $k_j - 1$. In particular, we can write the divisor

$$\operatorname{div}(\pi^*(dx_1 \wedge \cdots \wedge dx_n)) = \sum_{j \in J} (k_j - 1)E_j$$

Preliminaries — Resolution of singularities

- To each divisor $E_j, j \in J$ we have already associated a numerical quantity N_j representing the order of vanishing of a generic point of π^*f in E_j .
- Similarly, we define the integers k_j such that the order of vanishing of a generic point of the pullback of the standard volume form in E_j is $k_j - 1$. In particular, we can write the divisor

$$\operatorname{div}(\pi^*(dx_1 \wedge \cdots \wedge dx_n)) = \sum_{j \in J} (k_j - 1)E_j$$

The quantities (N_j, k_j) are the *numerical data* associated to each divisor E_j . From these, we may define

- *Candidate value* associated to the divisor E_j

$$\sigma_j = \frac{k_j}{N_j}$$

- *Residue number* associated to divisors E_i and E_j

$$\varepsilon(i, j) = -N_j \sigma_i + k_j = -N_j \frac{k_i}{N_i} + k_j$$

Preliminaries — Zeta function

Consider $f \in \mathbb{R}[x_1, \dots, x_n]$ a non-constant polynomial and a test function $\varphi: \mathbb{R}^n \rightarrow \mathbb{C}$. We construct the associated archimedean *zeta function* as

$$Z(s) = Z(f, \varphi; s) := \int_{\mathbb{R}^n} |f(x)|^s \varphi(x) \, dx$$

- Converges and is holomorphic in $\Re(s) > 0$.
- Question by I. Gel'fand in ICM 1954: possible meromorphic continuation and distribution of the poles?

Preliminaries — Zeta function

Consider $\pi: Y \rightarrow X = \mathbb{R}^n$ an embedded resolution

$$Z(s) = \int_Y |\pi^* f(y)|^s |\text{Jac}_\pi(y)| (\pi^* \varphi) dy$$

We only need a finite number of charts U_p , and with a partition of unity $\{\rho_p\}_p$ subordinate to it, we can write $Z(s)$ as a finite sum

$$Z(s) = \sum_p \int_{U_p} \rho_p(y) |u(y)|^s |v(y)| (\pi^* \varphi)(y) \prod_{i \in J_p} |y_i|^{N_i s + k_i - 1} dy_i$$

Theorem 2.1 [Vey24]

Let $f \in \mathbb{R}[x_1, \dots, x_n]$ be a non-constant polynomial and $\varphi: \mathbb{R}^n \rightarrow \mathbb{C}$ a \mathcal{C}^∞ function with compact support. Then $Z(f, \varphi; s)$ has a meromorphic continuation to \mathbb{C} , and its poles of the form $-\frac{k_j + \nu}{N_j}$, with $j \in J$ and ν a non-negative integer.

Preliminaries — Zeta function

Definition (Topological zeta function)

Let $f \in \mathbb{C}[x_1, \dots, x_n]$ be a non-constant polynomial and choose an embedded resolution $\pi: Y \rightarrow \mathbb{A}_{\mathbb{C}}^n$ of $\{f = 0\}$. The (*global*) *topological zeta function* of f is

$$Z_{\text{top}}(f; s) := \sum_{I \subset J} \chi(E_I^\circ) \prod_{i \in I} \frac{1}{k_i + N_i s}$$

and the *local topological zeta function* of f at $a \in \{f = 0\}$ is

$$Z_{\text{top}, a}(f; s) := \sum_{I \subset J} \chi(E_I^\circ \cap \pi^{-1}\{a\}) \prod_{i \in I} \frac{1}{k_i + N_i s}$$

where in both cases I runs through all possible subsets of J .

- Introduced heuristically through the 'limit $p \rightarrow 1$ ' of an expression of the p -adic zeta function, later formalized by Denef and Loeser.
- Independent of the chosen embedded resolution [DL92, Thm. 3.2].

Preliminaries — Bernstein-Sato polynomial

- Bernstein [Ber72] in the case of polynomials
- Kashiwara [Kas76] in the case of holomorphic functions
- Björk [Bjö73] in the case of formal power series

Theorem

Let $f \in R$ be a polynomial. Then, there exists a polynomial $P(s) \in \mathcal{D}[s]$ and a polynomial $e_{f,P}(s) \in \mathbb{C}[s]$ such that the relation

$$P(s)f^{s+1} = e_{f,P}(s)f^s$$

holds formally in the \mathcal{D} -module $R_f[s] \cdot f^s$.

Preliminaries — Bernstein-Sato polynomial

- Bernstein [Ber72] in the case of polynomials
- Kashiwara [Kas76] in the case of holomorphic functions
- Björk [Bjö73] in the case of formal power series

Theorem

Let $f \in R$ be a polynomial. Then, there exists a polynomial $P(s) \in \mathcal{D}[s]$ and a polynomial $e_{f,P}(s) \in \mathbb{C}[s]$ such that the relation

$$P(s)f^{s+1} = e_{f,P}(s)f^s$$

holds formally in the \mathcal{D} -module $R_f[s] \cdot f^s$.

Example

Let $f = x_1^2 + \cdots + x_n^2$ in $\mathbb{C}[x_1, \dots, x_n]$. Then, we have

$$\left(\frac{\partial^2}{\partial x_1^2} + \cdots + \frac{\partial^2}{\partial x_n^2} \right) f^{s+1} = 4(s+1) \left(s + \frac{n}{2} \right) f^s$$

Preliminaries — Bernstein-Sato polynomial

- Bernstein [Ber72] in the case of polynomials
- Kashiwara [Kas76] in the case of holomorphic functions
- Björk [Bjö73] in the case of formal power series

Theorem

Let $f \in R$ be a polynomial. Then, there exists a polynomial $P(s) \in \mathcal{D}[s]$ and a polynomial $b_{f,P}(s) \in \mathbb{C}[s]$ such that the relation

$$P(s)f^{s+1} = e_{f,P}(s)f^s$$

holds formally in the \mathcal{D} -module $R_f[s] \cdot f^s$.

Definition (Bernstein-Sato polynomial)

The Bernstein-Sato polynomial $b_f(s)$ is the monic generator of the ideal in $\mathbb{C}[s]$ consisting of polynomials $e_{f,P}(s)$ satisfying such a functional equation.

First algorithm to compute it introduced by Oaku in [Oak97], using non-commutative Gröbner basis in the Weyl algebra.

Preliminaries — Bernstein-Sato polynomial

Applying the functional equation to integrate by parts.

$$b_f(s)Z(s) = \int_{\mathbb{R}^n} b_f(s)f(x)^s \varphi(x) \, dx = \int_{\mathbb{R}^n} P(s) \cdot f(x)^{s+1} \varphi(x) \, dx = \int_{\mathbb{R}^n} f(x)^{s+1} \underbrace{P^*(s) \cdot \varphi(x)}_{\varphi_1(x)} \, dx$$

where P^* is the adjoint operator of P . Thus, we obtain a meromorphic continuation to the half-plane $\Re(s) > -1$ as

$$Z(s) = \frac{1}{b_f(s)} \int_{\mathbb{R}^n} f(x)^{s+1} \varphi_1(x) \, dx$$

Preliminaries — Bernstein-Sato polynomial

Applying the functional equation to integrate by parts.

$$b_f(s)Z(s) = \int_{\mathbb{R}^n} b_f(s)f(x)^s \varphi(x) dx = \int_{\mathbb{R}^n} P(s) \cdot f(x)^{s+1} \varphi(x) dx = \int_{\mathbb{R}^n} f(x)^{s+1} \underbrace{P^*(s) \cdot \varphi(x)}_{\varphi_1(x)} dx$$

where P^* is the adjoint operator of P . Thus, we obtain a meromorphic continuation to the half-plane $\Re(s) > -1$ as

$$Z(s) = \frac{1}{b_f(s)} \int_{\mathbb{R}^n} f(x)^{s+1} \varphi_1(x) dx$$

Iterating this procedure for $r > 1$

$$Z(s) = \frac{1}{b_f(s+r-1) \cdots b_f(s+1)b_f(s)} \int_{\mathbb{R}^n} f(x)^{s+r} \varphi_r(x) dx$$

Theorem 2.3 [Vey24]

Let $f \in \mathbb{R}[x_1, \dots, x_n]$ be a non-constant polynomial and $\varphi: \mathbb{R}^n \rightarrow \mathbb{C}$ a \mathcal{C}^∞ function with compact support. Then $Z(f, \varphi; s)$ has a meromorphic continuation to \mathbb{C} , and its poles are of the form $\lambda - \nu$ for λ a root of b_f and ν a non-negative integer.

Preliminaries — Bernstein-Sato polynomial

Roots of the Bernstein-Sato polynomial

- -1 is always a root. We write $\tilde{b}_f(s) = b_f(s)/(s+1)$ for the *reduced* BS polynomial.
- All roots are negative rational numbers [Mal75; Kas76].
- A set of candidate roots can be obtained from a resolution of singularities [Kol97; Lic89].

Preliminaries — Bernstein-Sato polynomial

Roots of the Bernstein-Sato polynomial

- -1 is always a root. We write $\tilde{b}_f(s) = b_f(s)/(s+1)$ for the *reduced* BS polynomial.
- All roots are negative rational numbers [Mal75; Kas76].
- A set of candidate roots can be obtained from a resolution of singularities [Kol97; Lic89].

Theorem

With the notations introduced for the resolution of singularities, we have that every root of the Bernstein-Sato polynomial b_f is of the form

$$-\frac{k_j + \nu}{N_j}, \quad j \in J, \nu \in \mathbb{Z}_{\geq 0}$$

Preliminaries — Bernstein-Sato polynomial

Roots of the Bernstein-Sato polynomial

- -1 is always a root. We write $\tilde{b}_f(s) = b_f(s)/(s+1)$ for the *reduced* BS polynomial.
- All roots are negative rational numbers [Mal75; Kas76].
- A set of candidate roots can be obtained from a resolution of singularities [Kol97; Lic89].

Theorem

With the notations introduced for the resolution of singularities, we have that every root of the Bernstein-Sato polynomial b_f is of the form

$$-\frac{k_j + \nu}{N_j}, \quad j \in J, \nu \in \mathbb{Z}_{\geq 0}$$

Question

Can we proceed similarly with the Bernstein-Sato polynomial to give a continuation for the topological zeta function?

Preliminaries — Strong Monodromy Conjecture

Let $f: (\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}, 0)$ be the germ of a holomorphic function on the origin. Fix $0 < \delta < \varepsilon$ and denote T disk of radius δ , $T' = T \setminus \{0\}$ and

$$X = B_\varepsilon \cap f^{-1}(T), \quad X' = X \setminus f^{-1}(0), \quad X_t = f^{-1}(t) \cap X, \quad t \in T$$

Theorem [Mil16, §4]

For small enough δ, ε , the restriction $f': X' \rightarrow T'$ is a locally trivial smooth fiber bundle, and the diffeomorphism type of any fiber X_t is independent of δ, ε and t .




Figure: Adapted from [Viu21, Fig. 4].


Preliminaries — Strong Monodromy Conjecture

Let $f: (\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}, 0)$ be the germ of a holomorphic function on the origin. Fix $0 < \delta < \varepsilon$ and denote T disk of radius δ , $T' = T \setminus \{0\}$ and

$$X = B_\varepsilon \cap f^{-1}(T), \quad X' = X \setminus f^{-1}(0), \quad X_t = f^{-1}(t) \cap X, \quad t \in T$$

Theorem [Mil16, §4]

For small enough δ, ε , the restriction $f': X' \rightarrow T'$ is a locally trivial smooth fiber bundle, and the diffeomorphism type of any fiber X_t is independent of δ, ε and t .



For an isolated singularity

$$H_i(X_t, \mathbb{C}) = \begin{cases} \mathbb{Z}, & i = 0 \\ \mathbb{Z}^\mu, & i = n \\ 0, & \text{else} \end{cases}$$

Figure: Adapted from [Viu21, Fig. 4].

Preliminaries — Strong Monodromy Conjecture

The action of $\pi_1(T', t) \cong \mathbb{Z}$ of the base of the fibration induces a diffeomorphism f on each fiber X_t , usually referred to as the *geometric monodromy*.

Also induces an automorphism on the homology and integer singular cohomology

$$h_* : H_*(X_t, \mathbb{C}) \longrightarrow H_*(X_t, \mathbb{C}), \quad h^* : H^\bullet(X_t, \mathbb{C}) \longrightarrow H^\bullet(X_t, \mathbb{C})$$

These are usually referred as the *algebraic (complex) monodromy*

Preliminaries — Strong Monodromy Conjecture

The action of $\pi_1(T', t) \cong \mathbb{Z}$ of the base of the fibration induces a diffeomorphism f on each fiber X_t , usually referred to as the *geometric monodromy*.

Also induces an automorphism on the homology and integer singular cohomology

$$h_* : H_*(X_t, \mathbb{C}) \longrightarrow H_*(X_t, \mathbb{C}), \quad h^* : H^\bullet(X_t, \mathbb{C}) \longrightarrow H^\bullet(X_t, \mathbb{C})$$

These are usually referred as the *algebraic (complex) monodromy*

Monodromy Theorem [RR06]

The operator h_* is quasi-unipotent, that is, there are p and q such that $(h_*^p - \text{id})^q = 0$. In other words, the eigenvalues of the monodromy are roots of unity. Moreover, one can take $q = n + 1$.

Preliminaries — Strong Monodromy Conjecture

- Formulated by the Igusa in the late seventies, after some examples computed.
- Originally stated in terms of the p -adic zeta function, but has analogous statements in topological and motivic settings.

Monodromy conjecture, topological version

Let $f \in \mathbb{C}[x_1, \dots, x_n]$ be non-constant. If s_0 is a pole of $Z_{\text{top}}(f, \varphi; s)$, then

- (*standard*) $e^{2\pi i \Re(s_0)}$ is a monodromy eigenvalue of $f: \mathbb{C}^n \rightarrow \mathbb{C}$ at some point of $\{f = 0\}$.
- (*strong*) s_0 is a root of the Bernstein-Sato polynomial b_f .

Preliminaries — Strong Monodromy Conjecture

- Formulated by the Igusa in the late seventies, after some examples computed.
- Originally stated in terms of the p -adic zeta function, but has analogous statements in topological and motivic settings.

Monodromy conjecture, topological version

Let $f \in \mathbb{C}[x_1, \dots, x_n]$ be non-constant. If s_0 is a pole of $Z_{\text{top}}(f, \varphi; s)$, then

- (*standard*) $e^{2\pi i \Re(s_0)}$ is a monodromy eigenvalue of $f: \mathbb{C}^n \rightarrow \mathbb{C}$ at some point of $\{f = 0\}$.
- (*strong*) s_0 is a root of the Bernstein-Sato polynomial b_f .

Strong version implies the standard one

Proposition 7.1 [Mal83]

If $-\alpha$ is a root of $\tilde{b}_{f,0}(s)$, then $e^{2\pi i \alpha}$ is an eigenvalue of the monodromy of f at the origin.

Preliminaries — Strong Monodromy Conjecture

- Formulated by the Igusa in the late seventies, after some examples computed.
- Originally stated in terms of the p -adic zeta function, but has analogous statements in topological and motivic settings.

Monodromy conjecture, topological version

Let $f \in \mathbb{C}[x_1, \dots, x_n]$ be non-constant. If s_0 is a pole of $Z_{\text{top}}(f, \varphi; s)$, then

- (*standard*) $e^{2\pi i \Re(s_0)}$ is a monodromy eigenvalue of $f: \mathbb{C}^n \rightarrow \mathbb{C}$ at some point of $\{f = 0\}$.
- (*strong*) s_0 is a root of the Bernstein-Sato polynomial b_f .

Remark

The strong version may even be formulated more precisely as stating that $b_f(s) \cdot Z_{\text{top}}(f, \varphi; s)$ is a polynomial.

Preliminaries — Periods of integrals

Fix $t = 1$ and choose a class of homology $\gamma(1) \in H_n(X_1, \mathbb{C})$, and from it we deduce a class $\gamma(t) \in H_n(X_t, \mathbb{C})$ in a neighborhood.

Consider $\pi \in \Omega^n$, with $\pi|_{X_t}$ holomorphic of maximal degree, thus closed. Define

$$I(t) := \int_{\gamma(t)} \pi$$

Preliminaries — Periods of integrals

Fix $t = 1$ and choose a class of homology $\gamma(1) \in H_n(X_1, \mathbb{C})$, and from it we deduce a class $\gamma(t) \in H_n(X_t, \mathbb{C})$ in a neighborhood.

Consider $\pi \in \Omega^n$, with $\pi|_{X_t}$ holomorphic of maximal degree, thus closed. Define

$$I(t) := \int_{\gamma(t)} \pi \qquad I'(t) = \frac{d}{dt} \int_{\gamma(t)} \pi = \int_{\gamma(t)} \frac{d\pi}{df}$$

- It is a holomorphic multivalued function on T' .
- If γ is a vanishing cycle (bounded and $\gamma(t) \xrightarrow{t \rightarrow 0} 0$), we have $\lim_{t \rightarrow 0} I(t) = 0$.
- Malgrange shows in [Mal74, p. 413] that we have a system of meromorphic differential equations satisfied by $I(t)$.

Preliminaries — Periods of integrals

Fix $t = 1$ and choose a class of homology $\gamma(1) \in H_n(X_1, \mathbb{C})$, and from it we deduce a class $\gamma(t) \in H_n(X_t, \mathbb{C})$ in a neighborhood.

Consider $\pi \in \Omega^n$, with $\pi|_{X_t}$ holomorphic of maximal degree, thus closed. Define

$$I(t) := \int_{\gamma(t)} \pi \qquad I'(t) = \frac{d}{dt} \int_{\gamma(t)} \pi = \int_{\gamma(t)} \frac{d\pi}{df}$$

- It is a holomorphic multivalued function on T' .
- If γ is a vanishing cycle (bounded and $\gamma(t) \xrightarrow[t \rightarrow 0]{} 0$), we have $\lim_{t \rightarrow 0} I(t) = 0$.
- Malgrange shows in [Mal74, p. 413] that we have a system of meromorphic differential equations satisfied by $I(t)$.

Consider a vanishing cycle γ satisfying $(h - \lambda)^p \gamma(1) = 0$, with p minimal, then

$$I(t) = \sum_{\substack{\mu \in L(\lambda) \\ 0 \leq q \leq p-1}} c_{\mu,q}(\pi) t^\mu (\log t)^q$$

where $L(\lambda) = \{\mu > 0 \mid e^{2\pi i \mu} = \lambda\}$.

Preliminaries — Periods of integrals

Next, consider $\omega \in \Omega^{n+1}$, and notice that there exists $\pi \in \Omega^n$ with $d\pi = \omega$, on the account of Poincaré's lemma. Then, we have

$$I'(t) = \int_{\gamma(t)} \frac{d\pi}{df} = \int_{\gamma(t)} \frac{\omega}{df} = \sum_{\substack{\mu \in L(\lambda)-1 \\ 0 \leq q \leq p-1}} d_{\mu,q}(\omega) t^\mu (\log t)^q$$

Preliminaries — Periods of integrals

Next, consider $\omega \in \Omega^{n+1}$, and notice that there exists $\pi \in \Omega^n$ with $d\pi = \omega$, on the account of Poincaré's lemma. Then, we have

$$I'(t) = \int_{\gamma(t)} \frac{d\pi}{df} = \int_{\gamma(t)} \frac{\omega}{df} = \sum_{\substack{\mu \in L(\lambda) - 1 \\ 0 \leq q \leq p-1}} d_{\mu,q}(\omega) t^\mu (\log t)^q$$

Defining $\mu_k := \inf\{\mu \in L(\lambda) - 1 \mid \exists q \geq k - 1, \omega \in \Omega^{n+1} \text{ with } d_{\mu,q}(\omega) \neq 0\}$.

Proposition 3.3 [Mal73]

The polynomial $(s + \mu_1) \dots (s + \mu_p)$ divides \tilde{b} .

Preliminaries — Periods of integrals

Next, consider $\omega \in \Omega^{n+1}$, and notice that there exists $\pi \in \Omega^n$ with $d\pi = \omega$, on the account of Poincaré's lemma. Then, we have

$$I'(t) = \int_{\gamma(t)} \frac{d\pi}{df} = \int_{\gamma(t)} \frac{\omega}{df} = \sum_{\substack{\mu \in L(\lambda) - 1 \\ 0 \leq q \leq p-1}} d_{\mu,q}(\omega) t^\mu (\log t)^q$$

Defining $\mu_k := \inf\{\mu \in L(\lambda) - 1 \mid \exists q \geq k - 1, \omega \in \Omega^{n+1} \text{ with } d_{\mu,q}(\omega) \neq 0\}$.

Proposition 3.3 [Mal73]

The polynomial $(s + \mu_1) \dots (s + \mu_p)$ divides \tilde{b} .

Conclusion

To apply this argument it only remains to show that there actually exists some non-zero homology class on $H_n(X_t, \mathbb{C})$. Or, equivalently, that a certain multivalued differential form defines a non-zero cohomology class.

Given by Loeser [Loe90, Thm. 3.7] for NND (toroidal version of [EV92]), and by Deligne and Mostow for plane curves [DM86, Prop. 2.14].

Contents

- 1 Introduction
- 2 Preliminaries
 - Resolution of singularities
 - Zeta function
 - Bernstein-Sato polynomial
 - Strong Monodromy Conjecture
 - Periods of integrals
- 3 Plane curves
- 4 Newton non-degenerate
- 5 Examples
 - Example 1
 - Example 2
 - Example 3
 - Example 4

Plane curves — Definition

We think of a plane curve as a set of points with coordinates (x, y) in the complex plane, described as the zero locus of some equation $f(x, y) = 0$, with f a holomorphic function.

Remark

The need to work on a bigger space than that of polynomials arises naturally.

$$f(x, y) = x^2 - y^3 = 0 \quad \implies \quad y = x^{2/3}$$

$$f(x, y) = x + y - xy \quad \implies \quad y = - \sum_{r \geq 1} x^r$$

Nonetheless, thanks to Weierstrass preparation theorem, we may (and will) consider f to be a polynomial.

Each curve $C: f = 0$ can be decomposed in a unique way as a finite union of branches $B_j: g_j = 0$, arising from the factorization

$$f = \prod g_j^{a_j}, \quad C = \sum a_j B_j$$

Plane curves — Resolution of singularities

Resolution of singularities for plane curves can be described as a composition of blowups.

- Only need a finite number of blowups to obtain smooth strict transform.
- Can be extended so that $\pi^{-1}(C)$ has simple normal crossings (all curves are smooth, with transverse intersection and no three meeting at a point).


Plane curves — Resolution of singularities

Resolution of singularities for plane curves can be described as a composition of blowups.

- Only need a finite number of blowups to obtain smooth strict transform.
- Can be extended so that $\pi^{-1}(C)$ has simple normal crossings (all curves are smooth, with transverse intersection and no three meeting at a point).

For example, a resolution of the polynomial $f = x^4 - y^5 + x^2y^2$ is given by

$$\begin{aligned} (x, y) &\mapsto (x_1y_1, y_1), & (x_1, y_1) &\mapsto (x_2, x_2y_2), & (x_2, y_2) &\mapsto (x_3, x_3y_3) \\ \implies \pi^*f &= x_3^{10}y_3^4(x_3^2 - y_3 + 1) \end{aligned}$$



Plane curves — Resolution of singularities

Definition (Dual graph)

Denote the *dual graph* of the curve C associated to the resolution π as $\Delta_\pi(C)$.

- Vertices V_i will correspond to the exceptional divisors E_i .
- Edges $V_i V_k$ whenever the curves E_i and E_k intersect.

Additionally, we will add arrowhead vertices W_j for each strict transform $B_j^{(N)}$, and call it *augmented dual graph* $\Delta_\pi^+(C)$.

Definition (Dual graph)

Denote the *dual graph* of the curve C associated to the resolution π as $\Delta_\pi(C)$.

- Vertices V_i will correspond to the exceptional divisors E_i .
- Edges $V_i V_k$ whenever the curves E_i and E_k intersect.

Additionally, we will add arrowhead vertices W_j for each strict transform $B_j^{(N)}$, and call it *augmented dual graph* $\Delta_\pi^+(C)$.

Constructing one blowup at a time, we can see that it is a tree.

- 1 If O_i is proximate only to O_{i-1} , add a new vertex V_i and a new edge $V_{i-1} V_i$.
- 2 If O_i is proximate to both O_{i-1} and some O_j with $j < i - 1$, subdivide the existing edge $V_{i-1} V_j$ by adding a new vertex V_i .

Plane curves — Resolution of singularities

Definition (Dual graph)

Denote the *dual graph* of the curve C associated to the resolution π as $\Delta_\pi(C)$.

- Vertices V_i will correspond to the exceptional divisors E_i .
- Edges $V_i V_k$ whenever the curves E_i and E_k intersect.

Additionally, we will add arrowhead vertices W_j for each strict transform $B_j^{(N)}$, and call it *augmented dual graph* $\Delta_\pi^+(C)$.

Constructing one blowup at a time, we can see that it is a tree.

- 1 If O_i is proximate only to O_{i-1} , add a new vertex V_i and a new edge $V_{i-1} V_i$.
- 2 If O_i is proximate to both O_{i-1} and some O_j with $j < i - 1$, subdivide the existing edge $V_{i-1} V_j$ by adding a new vertex V_i .

Remark


A sequence of proximity relations (or any equivalent information) about the curve determines the dual graph. The reverse is possible if we additionally label vertices.

Plane curves — Resolution of singularities

Observation

If a branch has Puiseux characteristic exponents $(n; \beta_1, \dots, \beta_g)$, then Δ_π^+ consists of a single core chain of edges from the initial vertex to the arrowhead vertex, with g side edge branches.

For example, consider the polynomial $f = y^4 - 2y^2x^3 - 4yx^5 + x^6 - x^7$, with Puiseux characteristic exponents $(4; 6, 7)$.



$$P = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 & 0 \\ -1 & -1 & 1 & 0 & 0 \\ 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & -1 & 1 \end{pmatrix}$$

$$P^t \cdot P = \begin{pmatrix} 3 & 0 & -1 & 0 & 0 \\ 0 & 2 & -1 & 0 & 0 \\ -1 & -1 & 3 & 0 & -1 \\ 0 & 0 & 0 & 2 & -1 \\ 0 & 0 & -1 & -1 & 1 \end{pmatrix}$$

Proposition

The topological zeta function is independent of a choice of resolution.

For plane curves, its expression may be simplified to

$$Z_{\text{top},0}(f; s) = \sum_{j \in J} \frac{\chi(E_j^\circ)}{k_j + N_j s} + \sum_{i \neq j \in J} \frac{\chi(E_i \cap E_j)}{(k_i + N_i s)(k_j + N_j s)}$$

Proposition

The topological zeta function is independent of a choice of resolution.

For plane curves, its expression may be simplified to

$$Z_{\text{top},0}(f; s) = \sum_{j \in J} \frac{\chi(E_j^\circ)}{k_j + N_j s} + \sum_{i \neq j \in J} \frac{\chi(E_i \cap E_j)}{(k_i + N_i s)(k_j + N_j s)}$$

- E_i° is a \mathbb{P}_1 with a point removed for each other divisor intersecting it, that is, the valence v_i in the dual graph.

Proposition

The topological zeta function is independent of a choice of resolution.

For plane curves, its expression may be simplified to

$$Z_{\text{top},0}(f; s) = \sum_{j \in J} \frac{\chi(E_j^\circ)}{k_j + N_j s} + \sum_{i \neq j \in J} \frac{\chi(E_i \cap E_j)}{(k_i + N_i s)(k_j + N_j s)}$$

- E_i° is a \mathbb{P}_1 with a point removed for each other divisor intersecting it, that is, the valence v_i in the dual graph.
- The intersection of any two divisors is either empty or a single point, in which case $\chi(E_i \cap E_j) = 1$.

Proposition

The topological zeta function is independent of a choice of resolution.

For plane curves, its expression may be simplified to

$$Z_{\text{top},0}(f; s) = \sum_{j \in V(\Delta_{\pi}^+)} \frac{2 - v_j}{k_j + N_j s} + \sum_{(i,j) \in E(\Delta_{\pi}^+)} \frac{1}{(k_i + N_i s)(k_j + N_j s)}$$

- E_i° is a \mathbb{P}_1 with a point removed for each other divisor intersecting it, that is, the valence v_i in the dual graph.
- The intersection of any two divisors is either empty or a single point, in which case $\chi(E_i \cap E_j) = 1$.

Plane curves — Topological zeta function

Observation

To study the final poles of the zeta function, we must study the quotient k/N over the divisors and the residue numbers $\varepsilon(i, j) = k_j - k_i \frac{N_j}{N_i}$

Plane curves — Topological zeta function

Observation

To study the final poles of the zeta function, we must study the quotient k/N over the divisors and the residue numbers $\varepsilon(i, j) = k_j - k_i \frac{N_j}{N_i}$

Proposition II.3.1 [Loe90]

Let V_i be a vertex of the dual graph. Then, $-1 \leq \varepsilon(i, j) < 1$ for all $j \in \mathcal{N}(i)$. Moreover, equality holds if, and only if, it has valence $v_i = 1$.

Plane curves — Topological zeta function

Observation

To study the final poles of the zeta function, we must study the quotient k/N over the divisors and the residue numbers $\varepsilon(i, j) = k_j - k_i \frac{N_j}{N_i}$

Proposition II.3.1 [Loe90]

Let V_i be a vertex of the dual graph. Then, $-1 \leq \varepsilon(i, j) < 1$ for all $j \in \mathcal{N}(i)$. Moreover, equality holds if, and only if, it has valence $v_i = 1$.

The vertices where the minimum of k/N can be attained form a connected chain, which is relevant to show

Theorem

The topological zeta function has at most one double pole, and the candidate value is $\min\{k_i/N_i\}$.

Plane curves — Topological zeta function

As for simple poles, we fix an exceptional divisor E_i and with partial fraction decomposition find its contribution to the zeta function as

$$\frac{1}{k_i + sN_i} \left(2 - v_i + \sum_{j \in \mathcal{N}(i)} \frac{1}{\varepsilon(i,j)} \right)$$

By the previous lemma and the recurrences over the dual graph, this quantity turns out to be zero for vertices V_i with valence $v_i = 1$ or 2 .

Plane curves — Topological zeta function

As for simple poles, we fix an exceptional divisor E_i and with partial fraction decomposition find its contribution to the zeta function as

$$\frac{1}{k_i + sN_i} \left(2 - v_i + \sum_{j \in \mathcal{N}(i)} \frac{1}{\varepsilon(i,j)} \right)$$


By the previous lemma and the recurrences over the dual graph, this quantity turns out to be zero for vertices V_i with valence $v_i = 1$ or 2 .

Theorem 8.4.5 [Wal04]

Any pole of the topological zeta function is of the form $s = -k_i/N_i$ for some vertex V_i which is either an arrowhead vertex or a rupture point.

Plane curves — Topological zeta function


Consider again the polynomial $f = y^4 - 2y^2x^3 - 4yx^5 + x^6 - x^7 = 0$, with numerical data



Divisor	k	N	σ	v_i
E_0	2	4	$\frac{1}{2}$	1
E_1	3	6	$\frac{1}{3}$	1
E_2	5	12	$\frac{5}{12}$	3
E_3	6	13	$\frac{6}{13}$	1
E_4	11	26	$\frac{11}{26}$	3
ST	1	1	1	1

Plane curves — Topological zeta function

Consider again the polynomial $f = y^4 - 2y^2x^3 - 4yx^5 + x^6 - x^7 = 0$, with numerical data



Divisor	k	N	σ	v_i
E_0	2	4	$\frac{1}{2}$	1
E_1	3	6	$\frac{1}{3}$	1
E_2	5	12	$\frac{5}{12}$	3
E_3	6	13	$\frac{6}{13}$	1
E_4	11	26	$\frac{11}{26}$	3
ST	1	1	1	1


$$E_2: \frac{1}{5+12s} \left(2 - 3 + \frac{1}{\varepsilon(2,0)} + \frac{1}{\varepsilon(2,1)} + \frac{1}{\varepsilon(2,4)} \right) = \frac{1}{5+12s} (-1 + 3 + 2 + 6) = \frac{5/6}{s + 5/12}$$

$$E_4: \frac{1}{11+26s} \left(2 - 3 + \frac{1}{\varepsilon(4,2)} + \frac{1}{\varepsilon(4,3)} + \frac{1}{\varepsilon(4,ST)} \right) = \frac{1}{11+26s} \left(-1 - 13 + 2 + \frac{26}{15} \right) = -\frac{77/195}{s + 11/26}$$

$$ST: \frac{1}{1+s} \left(2 - 1 + \frac{1}{\varepsilon(ST,4)} \right) = \frac{1}{1+s} \left(1 - \frac{1}{15} \right) = \frac{14/15}{s+1}$$

Plane curves — Topological zeta function

Consider again the polynomial $f = y^4 - 2y^2x^3 - 4yx^5 + x^6 - x^7 = 0$, with numerical data



Divisor	k	N	σ	ν_i
E_0	2	4	$\frac{1}{2}$	1
E_1	3	6	$\frac{1}{3}$	1
E_2	5	12	$\frac{5}{12}$	3
E_3	6	13	$\frac{6}{13}$	1
E_4	11	26	$\frac{11}{26}$	3
ST	1	1	1	1

Altogether

$$Z_0(s) = \frac{5/6}{s + 5/12} - \frac{77/195}{s + 11/26} + \frac{14/15}{s + 1}$$

Plane curves — Sketch of the proof

Proposition III.3.2 [Loe88]

If E_i is a rupture divisor, there exists a multivalued horizontal family $\gamma(t)$ of cycles in $H_1(X_t, \mathbb{C})$, $t \neq 0$, such that

$$\int_{\gamma(t)} \frac{dx \wedge dy}{df} = Ct^{-1+\frac{k_i}{N_i}} + o\left(t^{-1+\frac{k_i}{N_i}}\right)$$

with $C \neq 0$ a constant.

Theorem III.3.1 [Loe88]

Let $f: (\mathbb{C}^2, 0) \rightarrow (\mathbb{C}, 0)$ be the germ of an analytic function, and E_i a rupture divisor of the resolution of $\{f = 0\}$. Then

- ① $-k_i/N_i$ is a root of $\tilde{b}_{f,0}$ the local reduced Bernstein-Sato polynomial of f .
- ② If there exists $j \in \mathcal{N}(i)$ (a neighbor of V_i in the dual graph) such that $\varepsilon(i, j) = 0$, then $-k_i/N_i$ is a root of multiplicity two of $\tilde{b}_{f,0}$.

Contents

- 1 Introduction
- 2 Preliminaries
 - Resolution of singularities
 - Zeta function
 - Bernstein-Sato polynomial
 - Strong Monodromy Conjecture
 - Periods of integrals
- 3 Plane curves
- 4 Newton non-degenerate
- 5 Examples
 - Example 1
 - Example 2
 - Example 3
 - Example 4

Newton non-degenerate — Definition

We consider a polynomial $f(x_1, \dots, x_n) = \sum_{p \in \mathbb{N}^n} a_p x_1^p \dots x_n^p$ such that $f(0) = 0$, and define the support $\text{supp}(f) = \{p \in \mathbb{N}^n \mid a_p \neq 0\}$.

Definition (Newton polyhedron)


Let $f = \sum_{p \in \mathbb{N}^n} a_p x^p \in \mathbb{C}[x]$ with $f(0) = 0$. We define the *global Newton polyhedron* $\Gamma_{gl}(f)$ of f as the convex hull of $\text{supp}(f)$. Also, we define the *local Newton polyhedron* $\Gamma(f)$ as the convex hull of the set

$$\bigcup_{p \in \text{supp}(f)} p + (\mathbb{R}_{\geq 0})^n$$

In particular, it is immediate that $\Gamma(f) = \Gamma_{gl}(f) + (\mathbb{R}_{\geq 0})^n$.

Newton non-degenerate — Definition

Consider the polynomial $f = x^3 - y^2 + 4xy + 3x^2y$.



Newton non-degenerate — Definition


Consider the polynomial $f = x^3 - y^2 + 4xy + 3x^2y$.

- Faces of dimension $\dim \tau = 0$:

$$\tau_1 = \{(0, 2)\} \quad f^{\tau_1} = -y^2$$

$$\tau_2 = \{(1, 1)\} \quad f^{\tau_2} = 4xy$$

$$\tau_3 = \{(3, 0)\} \quad f^{\tau_3} = x^3$$



- Faces of dimension $\dim \tau = 1$:

$$\tau_4 = \{(0, 2) + \mathbb{R}_{>0}(0, 1)\}$$

$$f^{\tau_4} = -y^2$$

$$\tau_5 = \{(1 - \lambda)(0, 2) + \lambda(1, 1) \mid 0 < \lambda < 1\}$$

$$f^{\tau_5} = -y^2 + 4xy$$

$$\tau_6 = \{(1 - \lambda)(1, 1) + \lambda(3, 0) \mid 0 < \lambda < 1\}$$

$$f^{\tau_6} = x^3 + 4xy$$

$$\tau_7 = \{(3, 0) + \mathbb{R}_{>0}(1, 0)\}$$

$$f^{\tau_7} = x^3$$

Newton non-degenerate — Definition

Definition (Newton non-degenerate)

We say that f is Newton non-degenerate at 0 if for any face $\tau \subset \Gamma(f)$, the hypersurface $f^\tau = 0$ satisfies the condition

$$x_1 \frac{\partial f^\tau}{\partial x_1} = \cdots = x_n \frac{\partial f^\tau}{\partial x_n} = 0 \quad \implies \quad x_1 \cdots x_n = 0$$

that is, the polynomials $x_i \frac{\partial f^\tau}{\partial x_i}$ do not vanish at the same time in $(\mathbb{C} \setminus 0)^n$.

Newton non-degenerate — Definition

Definition (Newton non-degenerate)

We say that f is Newton non-degenerate at 0 if for any face $\tau \subset \Gamma(f)$, the hypersurface $f^\tau = 0$ satisfies the condition

$$x_1 \frac{\partial f^\tau}{\partial x_1} = \cdots = x_n \frac{\partial f^\tau}{\partial x_n} = 0 \quad \implies \quad x_1 \cdots x_n = 0$$

that is, the polynomials $x_i \frac{\partial f^\tau}{\partial x_i}$ do not vanish at the same time in $(\mathbb{C} \setminus 0)^n$.

For example, $f = y^4 - 2y^2x^3 - 4yx^5 + x^6 - x^7 = (y^2 - x^3)^2 - 4yx^5 - x^7$ is degenerate: the truncation on the face (edge) τ with endpoints $(0, 4)$ and $(6, 0)$ is

$$f^\tau = (y^2 - x^3)^2 \quad \implies \quad \begin{cases} x \cdot 2(y^2 - x^3) \cdot 3x^2 = 0 \\ y \cdot 2(y^2 - x^3) \cdot 2y = 0 \end{cases}$$

and the system considered has solutions outside $(\mathbb{C} \setminus 0)^2$.

Newton non-degenerate — Definition

Definition (N,k)

Let $\Gamma(f)$ be the Newton diagram of f as defined. For $a \in (\mathbb{R}^+)^n$, we define

$$N(a) := \inf_{x \in \Gamma(f)} \{\langle a, x \rangle\} = \min_{\substack{x \text{ vertex} \\ \text{of } \Gamma(f)}} \{\langle a, x \rangle\}$$

We may recover the face by considering $F(a) := \{x \in \Gamma(f) \mid \langle a, x \rangle = N(a)\}$ the first meet locus. Lastly, denote $k(a) := \sum_{i=1}^n a_i$.

Newton non-degenerate — Definition

Definition (N,k)

Let $\Gamma(f)$ be the Newton diagram of f as defined. For $a \in (\mathbb{R}^+)^n$, we define

$$N(a) := \inf_{x \in \Gamma(f)} \{\langle a, x \rangle\} = \min_{\substack{x \text{ vertex} \\ \text{of } \Gamma(f)}} \{\langle a, x \rangle\}$$

We may recover the face by considering $F(a) := \{x \in \Gamma(f) \mid \langle a, x \rangle = N(a)\}$ the first meet locus. Lastly, denote $k(a) := \sum_{i=1}^n a_i$.

Definition (Dual cone)

For τ a face of $\Gamma(f)$, we define the *cone associated* to τ as

$$\Delta_\tau := \{a \in (\mathbb{R}_{>0})^n \mid F(a) = \tau\} / \sim$$

where the equivalence relation is given by $a \sim a' \iff F(a) = F(a')$.

Additionally, we will refer to the collection of these cones for all faces of the Newton polytope as the *dual fan*.

Newton non-degenerate — Definition

Notice that for every proper face τ we have

$$\tau = \bigcap_{\substack{\tau \subset \gamma \\ \dim \gamma = n-1}} \gamma$$

Moreover, for every face of $\Gamma(f)$ of codimension 1, there exists a unique integral primitive vector (meaning that all of its coordinates are relatively coprime) perpendicular to the face.

Newton non-degenerate — Definition

Notice that for every proper face τ we have

$$\tau = \bigcap_{\substack{\tau \subset \gamma \\ \dim \gamma = n-1}} \gamma$$

Moreover, for every face of $\Gamma(f)$ of codimension 1, there exists a unique integral primitive vector (meaning that all of its coordinates are relatively coprime) perpendicular to the face.

Lemma

Let τ be a proper face of $\Gamma(f)$ and $\gamma_1, \dots, \gamma_r$ the faces of $\Gamma(f)$ of dimension $n - 1$ that contain it. Let a_1, \dots, a_r be the unique primitive normal vectors to $\gamma_1, \dots, \gamma_r$, respectively. Then,

$$\Delta_\tau = \{ \lambda_1 a_1 + \dots + \lambda_r a_r \mid \lambda_i \in \mathbb{R}_{>0} \}$$

with $\dim \Delta_\tau = n - \dim \tau$.

Newton non-degenerate — Definition

Consider again $f = x^3 - y^2 + 4xy + 3x^2y$.

- $\tau_4 = \{(0, 2) + \mathbb{R}_{>0}(0, 1)\} \implies a_4 = (1, 0)$
- $\tau_5 = \{(1 - \lambda)(0, 2) + \lambda(1, 1) \mid 0 < \lambda < 1\} \implies a_5 = (1, 1)$
- $\tau_6 = \{(1 - \lambda)(1, 1) + \lambda(3, 0) \mid 0 < \lambda < 1\} \implies a_6 = (1, 2)$
- $\tau_7 = \{(3, 0) + \mathbb{R}_{>0}(1, 0)\} \implies a_7 = (0, 1)$

Newton non-degenerate — Definition



Consider again $f = x^3 - y^2 + 4xy + 3x^2y$.

- $\tau_4 = \{(0, 2) + \mathbb{R}_{>0}(0, 1)\} \implies a_4 = (1, 0)$
- $\tau_5 = \{(1 - \lambda)(0, 2) + \lambda(1, 1) \mid 0 < \lambda < 1\} \implies a_5 = (1, 1)$
- $\tau_6 = \{(1 - \lambda)(1, 1) + \lambda(3, 0) \mid 0 < \lambda < 1\} \implies a_6 = (1, 2)$
- $\tau_7 = \{(3, 0) + \mathbb{R}_{>0}(1, 0)\} \implies a_7 = (0, 1)$

Then, the cones associated to each face

$$\Delta_{\tau_1} = \mathbb{R}_{>0}(1, 0) + \mathbb{R}_{>0}(1, 1), \quad \Delta_{\tau_4} = \mathbb{R}_{>0}(1, 0), \quad \dots$$

Altogether, the associated dual fan is (right)



Newton non-degenerate — Resolution of singularities

Definition (Cone)

A *convex polyhedral cone* is a set

$$C = \{\lambda_1 v_1 + \cdots + \lambda_r v_r \in V \mid \lambda_i \geq 0\} = \mathbb{R}_{\geq 0} v_1 + \cdots + \mathbb{R}_{\geq 0} v_r$$

where V is an n -dimensional vector space over \mathbb{R} , and the vectors $\{v_i\}$ are called the *generators* of the cone.

We will say that a cone is...

- *simplicial* if its generating vectors v_1, \dots, v_r are linearly independent over \mathbb{R} .
- *regular* if $\{v_1, \dots, v_r\}$ is a subset of a base of the \mathbb{Z} -module \mathbb{Z}^n .

Newton non-degenerate — Resolution of singularities

Definition (Cone)

A *convex polyhedral cone* is a set

$$C = \{\lambda_1 v_1 + \cdots + \lambda_r v_r \in V \mid \lambda_i \geq 0\} = \mathbb{R}_{\geq 0} v_1 + \cdots + \mathbb{R}_{\geq 0} v_r$$

where V is an n -dimensional vector space over \mathbb{R} , and the vectors $\{v_i\}$ are called the *generators* of the cone.

We will say that a cone is...

- *simplicial* if its generating vectors v_1, \dots, v_r are linearly independent over \mathbb{R} .
- *regular* if $\{v_1, \dots, v_r\}$ is a subset of a base of the \mathbb{Z} -module \mathbb{Z}^n .

Theorem

Each cone admits a partition in simplicial subcones. Moreover, they can be made regular cones by introducing suitable new generating rays.

Newton non-degenerate — Resolution of singularities

Definition (Cone)

A *convex polyhedral cone* is a set

$$C = \{\lambda_1 v_1 + \cdots + \lambda_r v_r \in V \mid \lambda_i \geq 0\} = \mathbb{R}_{\geq 0} v_1 + \cdots + \mathbb{R}_{\geq 0} v_r$$

where V is an n -dimensional vector space over \mathbb{R} , and the vectors $\{v_i\}$ are called the *generators* of the cone.

We will say that a cone is...

- *simplicial* if its generating vectors v_1, \dots, v_r are linearly independent over \mathbb{R} .
- *regular* if $\{v_1, \dots, v_r\}$ is a subset of a base of the \mathbb{Z} -module \mathbb{Z}^n .

Theorem

Each cone admits a partition in simplicial subcones. Moreover, they can be made regular cones by introducing suitable new generating rays.

Lemma 8.7 [AVG12]

There exists a regular fan subordinate to a Newton polyhedron.

Newton non-degenerate — Resolution of singularities

Definition (Toric blowup)

Consider a unimodular integral $n \times n$ matrix σ

$$\sigma = \begin{pmatrix} \sigma_{1,1} & \sigma_{1,2} & \cdots & \sigma_{1,n} \\ \sigma_{2,1} & \sigma_{2,2} & \cdots & \sigma_{2,n} \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{n,1} & \sigma_{n,2} & \cdots & \sigma_{n,n} \end{pmatrix}$$

We define the *toric blowup* associated to σ as the birational morphism

$$\begin{aligned} \pi_\sigma: (\mathbb{C}^*)^n &\rightarrow (\mathbb{C}^*)^n \\ (x_1, \dots, x_n) &\mapsto (x_1^{\sigma_{1,1}} x_2^{\sigma_{1,2}} \cdots x_n^{\sigma_{1,n}}, \dots, x_1^{\sigma_{n,1}} x_2^{\sigma_{n,2}} \cdots x_n^{\sigma_{n,n}}) \end{aligned}$$

Newton non-degenerate — Resolution of singularities

Definition (Toric blowup)

Consider a unimodular integral $n \times n$ matrix σ

$$\sigma = \begin{pmatrix} \sigma_{1,1} & \sigma_{1,2} & \cdots & \sigma_{1,n} \\ \sigma_{2,1} & \sigma_{2,2} & \cdots & \sigma_{2,n} \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{n,1} & \sigma_{n,2} & \cdots & \sigma_{n,n} \end{pmatrix}$$

We define the *toric blowup* associated to σ as the birational morphism

$$\begin{aligned} \pi_\sigma: (\mathbb{C}^*)^n &\rightarrow (\mathbb{C}^*)^n \\ (x_1, \dots, x_n) &\mapsto (x_1^{\sigma_{1,1}} x_2^{\sigma_{1,2}} \cdots x_n^{\sigma_{1,n}}, \dots, x_1^{\sigma_{n,1}} x_2^{\sigma_{n,2}} \cdots x_n^{\sigma_{n,n}}) \end{aligned}$$

For a regular simplicial cone of maximum dimension in the subdivided dual fan $\Sigma^*(f)$ given by vectors $\{r_1, \dots, r_n\}$, we can consider the matrix $\sigma = (r_1 \cdots r_n)$.


Gluing adequately these charts π_σ , we obtain a non-singular variety X , and a proper analytic map $\pi: X \rightarrow \mathbb{C}^n$: the toric blowup associated with $\Sigma^*(f)$.

Newton non-degenerate — Resolution of singularities

Theorem [Oka96, p. 101]

If f is Newton non-degenerate, then the associated toric blowup $\pi: X \rightarrow \mathbb{C}^n$ is a good resolution of the f as a germ at the origin.

Consider again $f = x^4 - y^5 + x^2y^2$.



Newton non-degenerate — Resolution of singularities

Theorem [Oka96, p. 101]

If f is Newton non-degenerate, then the associated toric blowup $\pi: X \rightarrow \mathbb{C}^n$ is a good resolution of the f as a germ at the origin.

Consider again $f = x^4 - y^5 + x^2y^2$.

The rays of the regular subdivision are

$$\begin{pmatrix} 1 & 2 & 3 & 1 & 0 \\ 0 & 1 & 2 & 1 & 1 \end{pmatrix}$$

taking one of the charts, we perform the toric blowup as

$$\begin{pmatrix} 3 & 1 \\ 2 & 1 \end{pmatrix} \implies (x, y) \mapsto (z^3w, z^2w)$$
$$\pi^*f = z^{10}w^4(z^2 - w + 1)$$



Newton non-degenerate — Topological zeta function

Let τ be a face in $\Gamma(f)$, and consider a decomposition of the associated cone $\Delta_\tau = \cup_{i=1}^r \Delta_i$ in simplicial cones of dimension $\dim \Delta_\tau = l$ such that $\dim(\Delta_i \cap \Delta_j) < l$, for all $i \neq j$. Then, define

$$J(\tau, s) := \sum_{i=1}^r J_{\Delta_i}(s), \quad \text{with} \quad J_{\Delta_i}(s) = \frac{\text{mult}(\Delta_i)}{(N(a_{i_1})s + k(a_{i_1})) \cdots (N(a_{i_l})s + k(a_{i_l}))}$$

being $a_{i_1}, \dots, a_{i_l} \in \mathbb{N}^n$ the linearly independent primitive integral vectors that generate Δ_i . Lastly, if $\tau = \Gamma(f)$, we rather take $J(\tau, s) = 1$.

Newton non-degenerate — Topological zeta function

Let τ be a face in $\Gamma(f)$, and consider a decomposition of the associated cone $\Delta_\tau = \cup_{i=1}^r \Delta_i$ in simplicial cones of dimension $\dim \Delta_\tau = l$ such that $\dim(\Delta_i \cap \Delta_j) < l$, for all $i \neq j$. Then, define

$$J(\tau, s) := \sum_{i=1}^r J_{\Delta_i}(s), \quad \text{with} \quad J_{\Delta_i}(s) = \frac{\text{mult}(\Delta_i)}{(N(a_{i_1})s + k(a_{i_1})) \cdots (N(a_{i_l})s + k(a_{i_l}))}$$

being $a_{i_1}, \dots, a_{i_l} \in \mathbb{N}^n$ the linearly independent primitive integral vectors that generate Δ_i . Lastly, if $\tau = \Gamma(f)$, we rather take $J(\tau, s) = 1$.

Theorem 5.3 [DL92]

Let $f: (\mathbb{C}^n, 0) \rightarrow (\mathbb{C}, 0)$ be a germ of a holomorphic function Newton non-degenerate for $\Gamma_0(f)$, then

$$Z_0(s) = \sum_{\substack{\tau \text{ vertex} \\ \text{of } \Gamma(f)}} J(\tau, s) + \left(\frac{s}{s+1} \right) \sum_{\substack{\tau \text{ compact} \\ \text{face of } \Gamma(f) \\ \dim \tau \geq 1}} (-1)^{\dim \tau} (\dim \tau)! \text{Vol}(\tau) J(\tau, s)$$

Newton non-degenerate — Sketch of the proof

We define the toric residue numbers, introduced by Loeser.

Definition (Toric residue numbers)

If τ, τ' are two distinct faces of codimension 1 of $\Gamma(f)$, we denote by $\beta(\tau, \tau')$ the greatest common divisor of the minors of order 2 of the matrix $(a(\tau), a(\tau'))$.

Additionally, one defines

$$\lambda(\tau, \tau') = k(\tau') - \frac{k(\tau)}{N(\tau)} N(\tau'), \quad \varepsilon(\tau, \tau') = \lambda(\tau, \tau') / \beta(\tau, \tau')$$

whenever $N(\tau) \neq 0$, which is the case if τ is a compact face.

- We are working with the original (not regularly subdivided) dual fan.
- The resulting variety might be singular.
- We lose some symmetry properties, but we gain 'uniqueness'.

Newton non-degenerate — Sketch of the proof

Theorem 4.2 [Loe90]

Let f be a germ of an analytic function, Newton non-degenerate at the origin. Let τ_0 be a compact face of codimension 1 of $\Gamma(f)$. Suppose that the following two conditions are verified

- ① $\frac{k(\tau_0)}{N(\tau_0)} \notin \mathbb{Z}$,
- ② For every face τ of codimension 1 of $\Gamma(f, 0)$, distinct of τ_0 and having non-empty intersection with τ_0 , we have $\varepsilon(\tau_0, \tau) \notin \mathbb{Z}$.

Then, there exists a horizontal multiform section $\gamma(t)$ of the fibration H_n over T' such that

$$\lim_{t \rightarrow 0} t^{1 - \frac{k(\tau_0)}{N(\tau_0)}} \int_{\gamma(t)} \frac{dx_1 \wedge \cdots \wedge dx_n}{df} = C$$

with C a non-zero constant.

The proof consists of constructing a non-zero multivalued differential form ω , which requires that the monodromies are not the identity, hence the hypothesis.

Newton non-degenerate — Sketch of the proof

Theorem 5.5.1 [Loe90]

Let f be a comfortable polynomial verifying $f(0) = 0$, with Newton diagram $\Gamma(f)$, and Newton non-degenerate. Suppose that all compact faces τ_0 verify

- i) $\frac{k(\tau_0)}{N(\tau_0)} < 1$,
- ii) For every face τ of codimension 1 of $\Gamma(f)$, distinct of τ_0 and having non-empty intersection with τ_0 , we have $\varepsilon(\tau_0, \tau) \notin \mathbb{Z}$.

Then, the real parts of the poles of the zeta function of f are roots of the Bernstein-Sato polynomial of f .

Remark 5.5.2.1 [Loe90]

If one replaces the condition $\frac{k(\tau_0)}{N(\tau_0)} < 1$ with $\frac{k(\tau_0)}{N(\tau_0)} \notin \mathbb{N}$, this is enough to prove the weak version of the conjecture.

Newton non-degenerate — Sketch of the proof

Question

Can the second hypothesis be relaxed?

Newton non-degenerate — Sketch of the proof

Question

Can the second hypothesis be relaxed?

It can be expected that non-positive residue numbers could be allowed to happen.

The idea for the negative integers is that we can argue the existence of a non-zero cohomology class, by calculating the degree of certain line bundles. For example, in the case of plane curves, a result in this spirit

Proposition 11.1 [Bla20]

Let $\omega \in \Gamma(\mathbb{P}, \Omega^1(\sum \mu_s s - \sum \delta_x x)(L))$. Assume that $\sum_{s \in S} \mu_s \leq r - 1$ and that $\alpha_s \neq 1$ for all $s \in S$. Then, ω defines a non-zero cohomology class in $H^1(\mathbb{P} \setminus S, L)$.



Contents

- 1 Introduction
- 2 Preliminaries
 - Resolution of singularities
 - Zeta function
 - Bernstein-Sato polynomial
 - Strong Monodromy Conjecture
 - Periods of integrals
- 3 Plane curves
- 4 Newton non-degenerate
- 5 Examples
 - Example 1
 - Example 2
 - Example 3
 - Example 4

Example 1

Consider the polynomial $f = xz^3 + y^3 \in \mathbb{C}[x, y, z]$.

- Found via brute force search
- Positive integer residue numbers already with low degree polynomial
- SMConjecture holds



Example 1

Associated residue numbers to divisor...

- $(0, 1, 1): \{ST: -2/3 + 1, (1, 1, 0): 4/3, (0, 1, 0): 1, (2, 1, 0): 5/3, (3, 1, 0): 2, (0, 0, 1): 1, (1, 1, 1): 1\}$

Example 1

Associated residue numbers to divisor...

- $(0, 1, 1): \{-2/3 + 1, (1, 1, 0): 4/3, (0, 1, 0): 1, (2, 1, 0): 5/3, (3, 1, 0): 2, (0, 0, 1): 1, (1, 1, 1): 1\}$

Observation

Now, we would like that the bad divisors with positive integer residue numbers didn't appear as poles in the zeta function.

Example 1

Associated residue numbers to divisor...

- $(0, 1, 1)$: $\{ST: -2/3 + 1, (1, 1, 0): 4/3, (0, 1, 0): 1, (2, 1, 0): 5/3, (3, 1, 0): 2, (0, 0, 1): 1, (1, 1, 1): 1\}$

Observation

Now, we would like that the bad divisors with positive integer residue numbers didn't appear as poles in the zeta function. However...

Local topological zeta function has poles $\{-1, -2/3\}$.

$$Z_0(s) = \frac{4/3}{s + 2/3} - \frac{1}{s + 1}$$

Example 1

Associated residue numbers to divisor...

- $(0, 1, 1)$: $\{ST: -2/3 + 1, (1, 1, 0): 4/3, (0, 1, 0): 1, (2, 1, 0): 5/3, (3, 1, 0): 2, (0, 0, 1): 1, (1, 1, 1): 1\}$

Observation

Now, we would like that the bad divisors with positive integer residue numbers didn't appear as poles in the zeta function. However...

Local topological zeta function has poles $\{-1, -2/3\}$.

$$Z_0(s) = \frac{4/3}{s + 2/3} - \frac{1}{s + 1}$$



Associated toric residue numbers to divisor...

- $(0, 1, 1)$: $\{ST: -2/3 + 1, (1, 0, 0): 1, (3, 1, 0): 2, (0, 0, 1): 1, (0, 1, 0): 1\}$

Example 2

Consider the convenient polynomial $f = x^5 + y^6 + z^4 + x^2yz + xy^2z \in \mathbb{C}[x, y, z]$.

- Built with geometric intuition.
- Two different integer residue numbers for a same divisor.
- SMConjecture holds.
- Toric residue numbers do satisfy the hypothesis.



Example 2

Regular subdivision requires over 60 new rays.

Observation

Computations for a regular subdivision rapidly increase in complexity, as the number of added rays increases largely.

Example 2

Regular subdivision requires over 60 new rays.

Observation

Computations for a regular subdivision rapidly increase in complexity, as the number of added rays increases largely.

Associated residue numbers to divisor...

- $(4, 7, 5)$: $\{\text{ST: } -4/5 + 1, \dots\}$
- $(1, 1, 2)$: $\{\text{ST: } -4/5 + 1, \dots\}$
- $(6, 5, 14)$: $\{\text{ST: } -5/6 + 1, \dots\}$
- $(5, 2, 3)$: $\{\text{ST: } -5/6 + 1, \dots\}$
- $(1, 1, 1)$: $\{\text{ST: } -3/4 + 1, \dots, (8, 5, 10): 2, (6, 5, 12): 2, \dots\}$



Local topological zeta function has poles $\{-3/4, -4/5, -5/6, -1\}$.

$$Z_0(s) = \frac{9}{s + 3/4} - \frac{48/5}{s + 4/5} - \frac{35/6}{s + 5/6} + \frac{8}{s + 1}$$

Example 3

Consider the convenient polynomial $f = x^5 + y^6 + z^7 + x^2yz + xyz^2 + xy^2z$.

- Built with geometric intuition.
- Integer residue numbers and integer toric residue numbers.
- Bad divisors in each case are different, and arise from compact faces.
- SMConjecture holds
- Contributions of bad divisors to the zeta function are non-zero.



Example 3

Regular subdivision requires almost 400 new rays.

Associated residue numbers to divisor...

- $(1, 1, 1): \{ST: -3/4 + 1, (16, 6, 5): 3, (17, 11, 6): 4, (9, 7, 4): 2, (21, 15, 8): 5, (13, 15, 6): 4, (11, 5, 4): 2, (21, 7, 6): 4, (4, 10, 3): 2, (19, 17, 8): 5, (6, 5, 12): 2, \dots\}$

Local topological zeta function has poles $\{-3/4, -4/5, -5/6, -6/7, -1\}$.

$$Z_0(s) = \frac{81/4}{s + 3/4} - \frac{72/5}{s + 4/5} - \frac{70/6}{s + 5/6} - \frac{48/7}{s + 6/7} + \frac{14}{s + 1}$$

Example 3

Regular subdivision requires almost 400 new rays.

Associated residue numbers to divisor...

- $(1, 1, 1)$: {ST: $-3/4 + 1$, $(16, 6, 5)$: 3, $(17, 11, 6)$: 4, $(9, 7, 4)$: 2, $(21, 15, 8)$: 5, $(13, 15, 6)$: 4, $(11, 5, 4)$: 2, $(21, 7, 6)$: 4, $(4, 10, 3)$: 2, $(19, 17, 8)$: 5, $(6, 5, 12)$: 2, ...}

Local topological zeta function has poles $\{-3/4, -4/5, -5/6, -6/7, -1\}$.

$$Z_0(s) = \frac{81/4}{s + 3/4} - \frac{72/5}{s + 4/5} - \frac{70/6}{s + 5/6} - \frac{48/7}{s + 6/7} + \frac{14}{s + 1}$$

Associated toric residue numbers to divisor...

- $(1, 2, 1)$: {ST: $-4/5 + 1$, ...}
- $(1, 1, 2)$: {ST: $-4/5 + 1$, $(7, 18, 5)$: 2, ...}

Example 3

Regular subdivision requires almost 400 new rays.

Associated residue numbers to divisor...

- $(1, 1, 1)$: {ST: $-3/4 + 1$, $(16, 6, 5)$: 3, $(17, 11, 6)$: 4, $(9, 7, 4)$: 2, $(21, 15, 8)$: 5, $(13, 15, 6)$: 4, $(11, 5, 4)$: 2, $(21, 7, 6)$: 4, $(4, 10, 3)$: 2, $(19, 17, 8)$: 5, $(6, 5, 12)$: 2, ...}

Local topological zeta function has poles $\{-3/4, -4/5, -5/6, -6/7, -1\}$.

$$Z_0(s) = \frac{81/4}{s + 3/4} - \frac{72/5}{s + 4/5} - \frac{70/6}{s + 5/6} - \frac{48/7}{s + 6/7} + \frac{14}{s + 1}$$

Associated toric residue numbers to divisor...

- $(1, 2, 1)$: {ST: $-4/5 + 1$, ...}
- $(1, 1, 2)$: {ST: $-4/5 + 1$, $(7, 18, 5)$: 2, ...}

Question

Is it possible that the pole arises from the good divisor? That is, the contribution of the bad divisor to $Z_0(s)$ has residue 0 at $s = -4/5$.

Example 3

Recall the expression for the local topological zeta function for NND polynomials

$$Z_0(s) = \sum_{\substack{\tau \text{ vertex} \\ \text{of } \Gamma(f)}} J(\tau, s) + \left(\frac{s}{s+1} \right) \sum_{\substack{\tau \text{ compact} \\ \text{face of } \Gamma(f) \\ \dim \tau \geq 1}} (-1)^{\dim \tau} (\dim \tau)! \text{Vol}(\tau) J(\tau, s)$$

where

$$J(\tau, s) := \sum_{i=1}^r J_{\Delta_i}(s), \quad \text{with} \quad J_{\Delta_i}(s) = \frac{\text{mult}(\Delta_i)}{(N(a_{i_1})s + k(a_{i_1})) \cdots (N(a_{i_l})s + k(a_{i_l}))}$$

and $\Delta_\tau = \cup_{i=1}^r \Delta_i$ is a decomposition in simplicial cones of dimension $\dim \Delta_\tau = l$ such that $\dim(\Delta_i \cap \Delta_j) < l$, for all $i \neq j$.

Example 3

Recall the expression for the local topological zeta function for NND polynomials

$$Z_0(s) = \sum_{\substack{\tau \text{ vertex} \\ \text{of } \Gamma(f)}} J(\tau, s) + \left(\frac{s}{s+1} \right) \sum_{\substack{\tau \text{ compact} \\ \text{face of } \Gamma(f) \\ \dim \tau \geq 1}} (-1)^{\dim \tau} (\dim \tau)! \text{Vol}(\tau) J(\tau, s)$$

where

$$J(\tau, s) := \sum_{i=1}^r J_{\Delta_i}(s), \quad \text{with} \quad J_{\Delta_i}(s) = \frac{\text{mult}(\Delta_i)}{(N(a_{i_1})s + k(a_{i_1})) \cdots (N(a_{i_l})s + k(a_{i_l}))}$$

and $\Delta_\tau = \cup_{i=1}^r \Delta_i$ is a decomposition in simplicial cones of dimension $\dim \Delta_\tau = l$ such that $\dim(\Delta_i \cap \Delta_j) < l$, for all $i \neq j$.

To extract the contribution of a single divisor $a = a(\tau)$, add up only the terms from cones Δ_i that contain a as one of its generating rays. Notice that this is not the same as simply taking the terms where a fraction of $\frac{1}{N(a)s+k(a)}$ appears.

Example 3

- Divisor $a_1 = (1, 1, 2)$, with $(k(a_1), N(a_1)) = (4, 5)$.
- For $a_2 = (1, 2, 1)$ also $\sigma(a_2) = 4/5$, since $(k(a_2), N(a_2)) = (4, 5)$.

We compute the residues of the contributions at the corresponding point

$$\operatorname{Res}_{s=-4/5} Z_{0;(1,2,1)}(s) = -\frac{47}{5}, \quad \operatorname{Res}_{s=-4/5} Z_{0;(1,1,2)}(s) = -\frac{47}{5}$$

Example 3

- Divisor $a_1 = (1, 1, 2)$, with $(k(a_1), N(a_1)) = (4, 5)$.
- For $a_2 = (1, 2, 1)$ also $\sigma(a_2) = 4/5$, since $(k(a_2), N(a_2)) = (4, 5)$.

We compute the residues of the contributions at the corresponding point

$$\operatorname{Res}_{s=-4/5} Z_{0;(1,2,1)}(s) = -\frac{47}{5}, \quad \operatorname{Res}_{s=-4/5} Z_{0;(1,1,2)}(s) = -\frac{47}{5}$$

Remark

One should be careful not to conclude that the total residue is simply the sum of the residues given by all divisors with the given candidate value.

Rays $(1, 1, 2)$ and $(1, 2, 1)$ appear in a same cone: we are double counting a term!

$$\begin{aligned} \operatorname{Res}_{s=-4/5} Z_0(s) &= \frac{-72}{5} = \frac{-47}{5} + \frac{-47}{5} - \frac{-22}{5} \\ &= \operatorname{Res}_{s=-4/5} Z_{0;(1,1,2)}(s) + \operatorname{Res}_{s=-4/5} Z_{0;(1,2,1)}(s) - \operatorname{Res}_{s=-4/5} Z_{0;(1,1,2),(1,2,1)}(s) \end{aligned}$$

Example 3

- Divisor $a_1 = (1, 1, 2)$, with $(k(a_1), N(a_1)) = (4, 5)$.
- For $a_2 = (1, 2, 1)$ also $\sigma(a_2) = 4/5$, since $(k(a_2), N(a_2)) = (4, 5)$.

We compute the residues of the contributions at the corresponding point

$$\operatorname{Res}_{s=-4/5} Z_{0;(1,2,1)}(s) = -\frac{47}{5}, \quad \operatorname{Res}_{s=-4/5} Z_{0;(1,1,2)}(s) = -\frac{47}{5}$$

Remark

One should be careful not to conclude that the total residue is simply the sum of the residues given by all divisors with the given candidate value.

Rays $(1, 1, 2)$ and $(1, 2, 1)$ appear in a same cone: we are double counting a term!

$$\begin{aligned} \operatorname{Res}_{s=-4/5} Z_0(s) &= \frac{-72}{5} = \frac{-47}{5} + \frac{-47}{5} - \frac{-22}{5} \\ &= \operatorname{Res}_{s=-4/5} Z_{0;(1,1,2)}(s) + \operatorname{Res}_{s=-4/5} Z_{0;(1,2,1)}(s) - \operatorname{Res}_{s=-4/5} Z_{0;(1,1,2),(1,2,1)}(s) \end{aligned}$$


Observation

In a more general situation, where there are more divisors with the same candidate σ , an inclusion-exclusion expression should be used in order to equate the residues.

Example 4

Consider the convenient polynomial $f = x^9 + y^8 + z^{11} + xyz^2 + xy^2z$.

- Built with geometric intuition (same as Example 3).
- Integer residue numbers and integer toric residue numbers.
- Bad divisors in each case are different, and arise from compact faces.
- SMConjecture holds
- Contributions of bad divisors to the zeta function are non-zero and different.



Example 4

Associated toric residue numbers to divisor...

- $(8, 9, 46)$: $\{\text{ST: } -7/8 + 1, (61, 11, 8): 3, \dots\}$
- $(5, 1, 1)$: $\{\text{ST: } -7/8 + 1, \dots\}$

Local topological zeta function has poles $\{-1, -7/8, -10/11, -19/27\}$.

$$Z_0(s) = \frac{18}{s+1} - \frac{63/4}{s+7/8} - \frac{120/11}{s+10/11} + \frac{247/27}{s+19/27}$$

Now, we compute the individual contributions

$$\begin{aligned} \operatorname{Res}_{s=-7/8} Z_0(s) &= \frac{-63}{4} = \frac{-535}{36} + \frac{-31}{4} - \frac{-247}{36} \\ &= \operatorname{Res}_{s=-7/8} Z_{0;(5,1,1)}(s) + \operatorname{Res}_{s=-7/8} Z_{0;(8,9,46)}(s) - \operatorname{Res}_{s=-7/8} Z_{0;(5,1,1),(8,9,46)}(s) \end{aligned}$$

Monodromy conjecture for Newton non-degenerate hypersurfaces

`https://github.com/baezaguasch/MonodromyNND`
`oriol.baeza (at) estudiantat.upc.edu`

References I



Vladimir Igorevich Arnold, Aleksandr Nikolaevich Varchenko, and Sabir Medzhidovich Gusein-Zade. *Singularities of differentiable maps: Volume II Monodromy and asymptotic integrals*. Vol. 83. Springer Science & Business Media, 2012 (cit. on pp. 74–76).



Joseph Bernstein. “The analytic continuation of generalized functions with respect to a parameter”. In: *Functional Analysis and its applications* 6.4 (1972), pp. 273–285 (cit. on pp. 19–21).



Jan-Erik Björk. *Dimensions over algebras of differential operators*. Département de mathématiques, 1973 (cit. on pp. 19–21).



Guillem Blanco. “Bernstein-Sato polynomial of plane curves and Yano’s conjecture”. PhD thesis. Universitat Politècnica de Catalunya, 2020 (cit. on pp. 86, 87).



Jan Denef and François Loeser. “Caractéristiques d’Euler-Poincaré, fonctions zêta locales et modifications analytiques”. In: *Journal of the American Mathematical Society* 5.4 (1992), pp. 705–720 (cit. on pp. 18, 81, 82).



Pierre Deligne and George Daniel Mostow. “Monodromy of hypergeometric functions and non-lattice integral monodromy”. In: *Publications Mathématiques de l’IHÉS* 63 (1986), pp. 5–89 (cit. on pp. 37–39).

References II



Hélène Esnault and Eckart Viehweg. *Lectures on vanishing theorems*. Vol. 20. Springer, 1992 (cit. on pp. 37–39).



Heisuke Hironaka. “Resolution of singularities of an algebraic variety over a field of characteristic zero: II”. In: *Annals of Mathematics* (1964), pp. 205–326 (cit. on p. 11).



Masaki Kashiwara. “b-functions and holonomic systems”. In: *Inventiones mathematicae* 38.1 (1976), pp. 33–53 (cit. on pp. 19–21, 24–26).



János Kollár. “Singularities of pairs”. In: *Proceedings of Symposia in Pure Mathematics*. Vol. 62. American Mathematical Society. 1997, pp. 221–288 (cit. on pp. 24–26).



Ben Lichtin. “Poles of $|f(z, w)|^{2s}$ and roots of the b-function”. In: *Arkiv för Matematik* 27.1 (1989), pp. 283–304 (cit. on pp. 24–26).



François Loeser. “Fonctions d’Igusa p -adiques et polynômes de Bernstein”. In: *American Journal of Mathematics* 110.1 (1988), pp. 1–21 (cit. on p. 60).



François Loeser. “Fonctions d’Igusa p -adiques, polynômes de Bernstein, et polyèdres de Newton.”. In: (1990) (cit. on pp. 6, 7, 37–39, 52–54, 84, 85).

References III



Bernard Malgrange. "Sur les polynômes de IN Bernstein". In: *Séminaire Équations aux dérivées partielles (Polytechnique) dit aussi "Séminaire Goulaouic-Schwartz"* (1973), pp. 1–10 (cit. on pp. 37–39).



Bernard Malgrange. "Intégrales asymptotiques et monodromie". In: *Annales scientifiques de l'École normale supérieure*. Vol. 7. 3. 1974, pp. 405–430 (cit. on pp. 34–36).



Bernard Malgrange. "Le polynôme de Bernstein d'une singularité isolée". In: *Fourier Integral Operators and Partial Differential Equations*. Berlin, Heidelberg: Springer Berlin Heidelberg, 1975, pp. 98–119. ISBN: 978-3-540-37521-0 (cit. on pp. 24–26).



Bernard Malgrange. "Polynômes de Bernstein-Sato et cohomologie évanescence". In: *Astérisque* 101.102 (1983), pp. 243–267 (cit. on pp. 31, 32).



John Milnor. *Singular Points of Complex Hypersurfaces (AM-61)*. Vol. 61. Princeton University Press, 2016 (cit. on pp. 27, 28).



Toshinori Oka. "Algorithms for the b-function and D-modules associated with a polynomial". In: *Journal of Pure and Applied Algebra* 117 (1997), pp. 495–518 (cit. on p. 21).



Mutsuo Oka. "Geometry of plane curves via toroidal resolution". In: *Algebraic geometry and singularities*. Springer, 1996, pp. 95–121 (cit. on pp. 79, 80).

References IV



Michèle Raynaud and Dock Sang Rim. *Groupes de Monodromie en Geometrie Algebrique: Seminaire de Geometrie Algebrique du Bois-Marie 1967-1969. (SGA 7 I)*. Vol. 288. Springer, 2006 (cit. on pp. 29, 30).



Willem Veys. *Introduction to the monodromy conjecture*. 2024. arXiv: 2403.03343 [math.AG]. URL: <https://arxiv.org/abs/2403.03343> (cit. on pp. 17, 22, 23).



Juan Viu-Sos. “An introduction to p -adic and motivic integration, zeta functions and invariants of singularities”. In: *Contemp. Math* 778 (2021) (cit. on pp. 27, 28).



Charles Terence Clegg Wall. *Singular points of plane curves*. 63. Cambridge University Press, 2004 (cit. on pp. 55, 56).