



and fore defination
$$F: u_2 - u_1 : U_1^{p_1} cos b + U_1^{p_2} cos b + U_2^{p_3} cos b + U_3^{p_4} cos b + U_3^{p_4} cos b + U_3^{p_5} cos b + U_3^{p_6} c$$

Dalam bentur matriks

$$\begin{bmatrix} k^{(1)}c^2\theta_1 & k^{(1)}s\theta_1c\theta_1 & 0 & 0 & -k^{(1)}c^2\theta_1 & -k^{(1)}s\theta_1c\theta_1 \\ k^{(1)}s\theta_1c\theta_1 & k^{(1)}s^2\theta_1 & 0 & 0 & -k^{(1)}s\theta_1c\theta_1 & -k^{(1)}s^2\theta_1 \\ 0 & 0 & k^{(2)}c^2\theta_2 & k^{(2)}s\theta_2c\theta_2 & -k^{(2)}c^2\theta_2 & -k^{(2)}s\theta_2c\theta_2 \\ 0 & 0 & k^{(2)}s\theta_2c\theta_2 & k^{(2)}s^2\theta_2 & -k^{(2)}s\theta_2c\theta_2 & -k^{(2)}s\theta_2c\theta_2 \\ -k^{(1)}c^2\theta_{12} & -k_1s\theta_1c\theta_1 & -k^{(2)}c^2\theta_2 & -k^{(2)}s\theta_2c\theta_2 & k^{(1)}c^2\theta_1 + & k^{(1)}s\theta_1c\theta_1 + \\ -k_1s\theta_1c\theta_1 & -k^{(1)}s^2\theta_1 & -k^{(2)}s\theta_2c\theta_2 & -k^{(2)}s^2\theta_2 & k^{(1)}s\theta_1c\theta_1 + \\ -k_1s\theta_1c\theta_1 & -k^{(1)}s^2\theta_1 & -k^{(2)}s\theta_2c\theta_2 & -k^{(2)}s^2\theta_2 & k^{(1)}s\theta_1c\theta_1 + & k^{(1)}s^2\theta_1 + \\ -k_1s\theta_1c\theta_1 & -k^{(1)}s^2\theta_1 & -k^{(2)}s\theta_2c\theta_2 & -k^{(2)}s^2\theta_2 & k^{(2)}s\theta_2c\theta_2 \end{bmatrix}$$

EXT LUS = LEY

GISbar stythen manz

hoda dipie clivent Vodal forces







