




# Linear Last-iterate Convergence in Constrained Saddle-point Optimization

Chung-Wei Lee



joint with **Haipeng Luo**, **Chen-Yu Wei** and **Mengxiao Zhang**



## A One-sentence (Informal) Summary

We prove that the last-iterate of  
**Optimistic Gradient Descent Ascent (OGDA)** and  
**Optimistic Multiplicative Weights Update (OMWU)**  
converges to the Nash equilibrium **exponentially** fast,  
in various constrained settings including matrix games and  
strongly-convex-strongly-concave functions.

# Saddle-point Optimization

- Consider constrained saddle-point optimization in the form

$$\min_{\mathbf{x} \in \mathcal{X}} \max_{\mathbf{y} \in \mathcal{Y}} f(\mathbf{x}, \mathbf{y}),$$

where  $\mathcal{X}$  and  $\mathcal{Y}$  are closed convex sets, and  $f$  is a continuous differentiable function that is convex in  $\mathbf{x}$  and concave in  $\mathbf{y}$ .

# Saddle-point Optimization

- Consider constrained saddle-point optimization in the form

$$\min_{\mathbf{x} \in \mathcal{X}} \max_{\mathbf{y} \in \mathcal{Y}} f(\mathbf{x}, \mathbf{y}),$$

where  $\mathcal{X}$  and  $\mathcal{Y}$  are closed convex sets, and  $f$  is a continuous differentiable function that is convex in  $\mathbf{x}$  and concave in  $\mathbf{y}$ .

- Goal: find a *Nash equilibrium*  $(\mathbf{x}^*, \mathbf{y}^*) \in \mathcal{X}^* \times \mathcal{Y}^*$  satisfying

$$f(\mathbf{x}^*, \mathbf{y}) \leq f(\mathbf{x}^*, \mathbf{y}^*) \leq f(\mathbf{x}, \mathbf{y}^*)$$

for any  $(\mathbf{x}, \mathbf{y}) \in \mathcal{X} \times \mathcal{Y}$ .

# “No-Regret” Algorithms

- (Projected) Gradient Descent Ascent (GDA):

$$\mathbf{x}_{t+1} = \Pi_{\mathcal{X}} (\mathbf{x}_t - \eta \nabla_{\mathbf{x}} f(\mathbf{x}_t, \mathbf{y}_t)), \quad \mathbf{y}_{t+1} = \Pi_{\mathcal{Y}} (\mathbf{y}_t + \eta \nabla_{\mathbf{y}} f(\mathbf{x}_t, \mathbf{y}_t))$$

# “No-Regret” Algorithms

- (Projected) Gradient Descent Ascent (GDA):

$$\mathbf{x}_{t+1} = \Pi_{\mathcal{X}}(\mathbf{x}_t - \eta \nabla_{\mathbf{x}} f(\mathbf{x}_t, \mathbf{y}_t)), \quad \mathbf{y}_{t+1} = \Pi_{\mathcal{Y}}(\mathbf{y}_t + \eta \nabla_{\mathbf{y}} f(\mathbf{x}_t, \mathbf{y}_t))$$

- Multiplicative Weights Update (MWU):

$$\mathbf{x}_{t+1} \propto (\mathbf{x}_t \odot \exp(-\eta \nabla_{\mathbf{x}} f(\mathbf{x}_t, \mathbf{y}_t))), \quad \mathbf{y}_{t+1} \propto (\mathbf{y}_t \odot \exp(\eta \nabla_{\mathbf{y}} f(\mathbf{x}_t, \mathbf{y}_t))),$$

when  $\mathcal{X}$  and  $\mathcal{Y}$  are simplex, and  $\odot$  denotes the element-wise product.

# Optimistic No-Regret Algorithms

- Optimistic Gradient Descent Ascent (OGDA):

$$\begin{aligned}\hat{\mathbf{x}}_{t+1} &= \Pi_{\mathcal{X}}(\hat{\mathbf{x}}_t - \eta \nabla_{\mathbf{x}} f(\mathbf{x}_t, \mathbf{y}_t)), & \hat{\mathbf{y}}_{t+1} &= \Pi_{\mathcal{Y}}(\hat{\mathbf{y}}_t + \eta \nabla_{\mathbf{y}} f(\mathbf{x}_t, \mathbf{y}_t)) \\ \mathbf{x}_{t+1} &= \Pi_{\mathcal{X}}(\hat{\mathbf{x}}_{t+1} - \eta \nabla_{\mathbf{x}} f(\mathbf{x}_t, \mathbf{y}_t)), & \mathbf{y}_{t+1} &= \Pi_{\mathcal{Y}}(\hat{\mathbf{y}}_{t+1} + \eta \nabla_{\mathbf{y}} f(\mathbf{x}_t, \mathbf{y}_t))\end{aligned}$$

# Optimistic No-Regret Algorithms

- Optimistic Gradient Descent Ascent (OGDA):

$$\begin{aligned}\hat{\mathbf{x}}_{t+1} &= \Pi_{\mathcal{X}}(\hat{\mathbf{x}}_t - \eta \nabla_{\mathbf{x}} f(\mathbf{x}_t, \mathbf{y}_t)), & \hat{\mathbf{y}}_{t+1} &= \Pi_{\mathcal{Y}}(\hat{\mathbf{y}}_t + \eta \nabla_{\mathbf{y}} f(\mathbf{x}_t, \mathbf{y}_t)) \\ \mathbf{x}_{t+1} &= \Pi_{\mathcal{X}}(\hat{\mathbf{x}}_{t+1} - \eta \nabla_{\mathbf{x}} f(\mathbf{x}_t, \mathbf{y}_t)), & \mathbf{y}_{t+1} &= \Pi_{\mathcal{Y}}(\hat{\mathbf{y}}_{t+1} + \eta \nabla_{\mathbf{y}} f(\mathbf{x}_t, \mathbf{y}_t))\end{aligned}$$

- Optimistic Multiplicative Weights Update (OMWU):

$$\begin{aligned}\hat{\mathbf{x}}_{t+1} &\propto (\hat{\mathbf{x}}_t \odot \exp(-\eta \nabla_{\mathbf{x}} f(\mathbf{x}_t, \mathbf{y}_t))), & \hat{\mathbf{y}}_{t+1} &\propto (\hat{\mathbf{y}}_t \odot \exp(\eta \nabla_{\mathbf{y}} f(\mathbf{x}_t, \mathbf{y}_t))) \\ \mathbf{x}_{t+1} &\propto (\hat{\mathbf{x}}_{t+1} \odot \exp(-\eta \nabla_{\mathbf{x}} f(\mathbf{x}_t, \mathbf{y}_t))), & \mathbf{y}_{t+1} &\propto (\hat{\mathbf{y}}_{t+1} \odot \exp(\eta \nabla_{\mathbf{y}} f(\mathbf{x}_t, \mathbf{y}_t)))\end{aligned}$$



# Average-iterate Convergence

- Convergence of average-iterate  $(\frac{1}{T} \sum_{t=1}^T \mathbf{x}_t, \frac{1}{T} \sum_{t=1}^T \mathbf{y}_t)$  is well known in many settings.

# Average-iterate Convergence

- Convergence of average-iterate  $(\frac{1}{T} \sum_{t=1}^T \mathbf{x}_t, \frac{1}{T} \sum_{t=1}^T \mathbf{y}_t)$  is well known in many settings.
- GDA and MWU are known to enjoy a converging duality gap of  $\mathcal{O}(1/\sqrt{T})$ . [FS99]

# Average-iterate Convergence

- Convergence of average-iterate  $(\frac{1}{T} \sum_{t=1}^T \mathbf{x}_t, \frac{1}{T} \sum_{t=1}^T \mathbf{y}_t)$  is well known in many settings.
- GDA and MWU are known to enjoy a converging duality gap of  $\mathcal{O}(1/\sqrt{T})$ . [FS99]
- Optimistic algorithms such as OGDA and OMWU improve the converging rate to  $\mathcal{O}(1/T)$ . [RS13,DDK15,SALS15]

# Average-iterate Convergence

- Convergence of average-iterate  $(\frac{1}{T} \sum_{t=1}^T \mathbf{x}_t, \frac{1}{T} \sum_{t=1}^T \mathbf{y}_t)$  is well known in many settings.
- GDA and MWU are known to enjoy a converging duality gap of  $\mathcal{O}(1/\sqrt{T})$ . [FS99]
- Optimistic algorithms such as OGDA and OMWU improve the converging rate to  $\mathcal{O}(1/T)$ . [RS13,DDK15,SALS15]
- However, averaging large neural networks is usually prohibited.

# Average-iterate Convergence

- Convergence of average-iterate  $(\frac{1}{T} \sum_{t=1}^T \mathbf{x}_t, \frac{1}{T} \sum_{t=1}^T \mathbf{y}_t)$  is well known in many settings.
- GDA and MWU are known to enjoy a converging duality gap of  $\mathcal{O}(1/\sqrt{T})$ . [FS99]
- Optimistic algorithms such as OGDA and OMWU improve the converging rate to  $\mathcal{O}(1/T)$ . [RS13,DDK15,SALS15]
- However, averaging large neural networks is usually prohibited.
- This motivates us to consider the last-iterate  $(\mathbf{x}_T, \mathbf{y}_T)$  convergence.

# Last-iterate Convergence

- For MWU and GDA, last-iterate **diverges**.

[BP18,CP19]

# Last-iterate Convergence

- For MWU and GDA, last-iterate **diverges**. [BP18,CP19]
- On the contrary, for *Extra-Gradient*, a standard algorithm for saddle-point optimization, last-iterate convergence has been shown in various settings. [T95,LS19,MOP20]

# Last-iterate Convergence

- For MWU and GDA, last-iterate **diverges**. [BP18,CP19]
- On the contrary, for *Extra-Gradient*, a standard algorithm for saddle-point optimization, last-iterate convergence has been shown in various settings. [T95,LS19,MOP20]
- OMWU achieves last-iterate convergence when  $f$  is bilinear on simplex (i.e. matrix game) when Nash Equilibrium is unique. [DP19]



# Last-iterate Convergence

- For MWU and GDA, last-iterate **diverges**. [BP18,CP19]
- On the contrary, for *Extra-Gradient*, a standard algorithm for saddle-point optimization, last-iterate convergence has been shown in various settings. [T95,LS19,MOP20]
- OMWU achieves last-iterate convergence when  $f$  is bilinear on simplex (i.e. matrix game) when Nash Equilibrium is unique. [DP19]
  - ▶ No concrete convergence rate.

# Last-iterate Convergence

- For MWU and GDA, last-iterate **diverges**. [BP18,CP19]
- On the contrary, for *Extra-Gradient*, a standard algorithm for saddle-point optimization, last-iterate convergence has been shown in various settings. [T95,LS19,MOP20]
- OMWU achieves last-iterate convergence when  $f$  is bilinear on simplex (i.e. matrix game) when Nash Equilibrium is unique. [DP19]
  - ▶ No concrete convergence rate.
  - ▶ Learning rate is exponentially small, which is inconsistent with practice.

# Last-iterate Convergence

- For MWU and GDA, last-iterate **diverges**. [BP18,CP19]
- On the contrary, for *Extra-Gradient*, a standard algorithm for saddle-point optimization, last-iterate convergence has been shown in various settings. [T95,LS19,MOP20]
- OMWU achieves last-iterate convergence when  $f$  is bilinear on simplex (i.e. matrix game) when Nash Equilibrium is unique. [DP19]
  - ▶ No concrete convergence rate.
  - ▶ Learning rate is exponentially small, which is inconsistent with practice.
- OGDA achieves last-iterate convergence

# Last-iterate Convergence

- For MWU and GDA, last-iterate **diverges**. [BP18,CP19]
- On the contrary, for *Extra-Gradient*, a standard algorithm for saddle-point optimization, last-iterate convergence has been shown in various settings. [T95,LS19,MOP20]
- OMWU achieves last-iterate convergence when  $f$  is bilinear on simplex (i.e. matrix game) when Nash Equilibrium is unique. [DP19]
  - ▶ No concrete convergence rate.
  - ▶ Learning rate is exponentially small, which is inconsistent with practice.
- OGDA achieves last-iterate convergence
  - ▶ when  $\mathcal{X}$  and  $\mathcal{Y}$  are unconstrained. [DISZ18,DP18,LS19,MOP19]

# Last-iterate Convergence

- For MWU and GDA, last-iterate **diverges**. [BP18,CP19]
- On the contrary, for *Extra-Gradient*, a standard algorithm for saddle-point optimization, last-iterate convergence has been shown in various settings. [T95,LS19,MOP20]
- OMWU achieves last-iterate convergence when  $f$  is bilinear on simplex (i.e. matrix game) when Nash Equilibrium is unique. [DP19]
  - ▶ No concrete convergence rate.
  - ▶ Learning rate is exponentially small, which is inconsistent with practice.
- OGDA achieves last-iterate convergence
  - ▶ when  $\mathcal{X}$  and  $\mathcal{Y}$  are unconstrained. [DISZ18,DP18,LS19,MOP19]

# Last-iterate Convergence

- For MWU and GDA, last-iterate **diverges**. [BP18,CP19]
- On the contrary, for *Extra-Gradient*, a standard algorithm for saddle-point optimization, last-iterate convergence has been shown in various settings. [T95,LS19,MOP20]
- OMWU achieves last-iterate convergence when  $f$  is bilinear on simplex (i.e. matrix game) when Nash Equilibrium is unique. [DP19]
  - ▶ No concrete convergence rate.
  - ▶ Learning rate is exponentially small, which is inconsistent with practice.
- OGDA achieves last-iterate convergence
  - ▶ when  $\mathcal{X}$  and  $\mathcal{Y}$  are unconstrained. [DISZ18,DP18,LS19,MOP19]

Question: Whether OGDA and OMWU can achieve last-iterate convergence in **constrained** saddle-point optimization with **concrete** convergence rate?

# Our Contributions

- Under uniqueness assumption made by Daskalakis and Panageas (2019), we show that OMWU with **constant** learning rate has **exponential** convergence rate.

# Our Contributions

- Under uniqueness assumption made by Daskalakis and Panageas (2019), we show that OMWU with **constant** learning rate has **exponential** convergence rate.
- For OGDA, we get more general results: under a sufficient condition called SP-MS, OGDA with constant learning rate converges exponentially fast.



# Our Contributions

- Under uniqueness assumption made by Daskalakis and Panageas (2019), we show that OMWU with **constant** learning rate has **exponential** convergence rate.
- For OGDA, we get more general results: under a sufficient condition called SP-MS, OGDA with constant learning rate converges exponentially fast.
- The SP-MS condition includes many settings such as bilinear games over any polytope and strongly-convex-strongly-concave functions **without** uniqueness assumption.

# Experiments




Figure: Experiments of OGDA and OMWU with different learning rates for a matrix game.

# Future directions

- One future direction is to get rid of the uniqueness assumption for OMWU.

## Future directions

- One future direction is to get rid of the uniqueness assumption for OMWU.
- It is also interesting to generalize the results to Markov/Stochastic Games.

## Future directions

- One future direction is to get rid of the uniqueness assumption for OMWU.
- It is also interesting to generalize the results to Markov/Stochastic Games.
- For this direction, see our new paper *Last-iterate Convergence of Decentralized Optimistic Gradient Descent/Ascent in Infinite-horizon Competitive Markov Games* on arXiv.