

1. Problem 1-1 (p15)

Table 1: Comparison of Running Times

$f(n)$	1 second	1 hour
$\log n$	$2^{1,000,000}$	$2^{3,600,000,000}$
\sqrt{n}	$1 \cdot 10^{12}$	$1.296 \cdot 10^{19}$
n	$1 \cdot 10^6$	$3.6 \cdot 10^9$
$n \log n$	62,746	$1.334 \cdot 10^8$
n^2	1,000	60,000
n^3	100	1,532
2^n	20	31
$n!$	9	12

2. Exercise 2.3-4 (p44)

Prove that when $n \geq 2$ is an exact power of 2, the solution of the recurrence

$$T(n) = \begin{cases} 2 & n = 2, \\ 2T(n/2) + n & n > 2 \end{cases}$$

is $T(n) = n \log n$.

3. Problem 2-3 (p46)

(a) $\Theta(n)$

(b) `naiveHorner(A,n,x)`

```
p = 0
for i = 0 to n
    item = A[i]
    for j = 1 downto i
        item *= x
    p += item
return p
```

The running time of `naiveHorner` is $\Theta(n^2)$, and is beat by `Horner` asymptotically.

(c)

4. Exercise 3.2-2 (p62)

The statement “The running time of algorithm A is at least $O(n^2)$ ” is meaningless, because $O(n^2)$ defines an upper bound on the asymptotic behavior of the algorithm, so it will never have a time complexity greater than $O(n^2)$.

5. Exercise 3.2-6 (p63)

Prove that $o(g(n)) \cap \omega(g(n))$ is the empty set.

o -notation denotes an asymptotically loose upper bound, formally defined as the set

$$o(g(n)) = \{f(n) : \text{for any positive constant } c > 0, \text{ there exists a constant } n_0 \text{ such that } 0 \leq f(n) < cg(n) \text{ for all } n \geq n_0.\}$$

Additionally, ω -notation denotes an asymptotically loose lower bound, formally defined as the set

$$\omega(g(n)) = \{f(n) : \text{for any positive constant } c > 0, \text{ there exists a constant } n_0 \text{ such that } 0 \leq cg(n) < f(n) \text{ for all } n \geq n_0.\}$$

So, we clearly see that the intersection of these two sets is the empty set, because $f(n)$ cannot be both exclusively less than $cg(n)$ and exclusively greater than $cg(n)$ simultaneously.

6. Using the substitution method, show that the solution of $T(n) = T(\lceil n/2 \rceil) + 1$ is $O(\log n)$.

Show: $T(n)$ is $O(\log n) \rightarrow T(n) \leq c \log n$

$$\forall n, \lceil n/2 \rceil \leq \frac{n+1}{2}$$

$$\log(n+1) < \log n + \frac{1}{2} \text{ for } n > 1$$

Assume: $T(\frac{n+1}{2}) \leq c \log \frac{n+1}{2}$

$$\begin{aligned} T(n) &= T(\lceil n/2 \rceil) + 1 \\ &\leq T(\frac{n+1}{2}) + 1 \\ &= c \log \frac{n+1}{2} + 1 \\ &= c \log \frac{n+1}{2} + \log 2 \\ &= c \log(n+1) \\ &= c \log n + \frac{c}{2} \text{ for } c \geq 2 \end{aligned}$$

Picking $c = \max\{T(2), T(3)\}$ yields $T(2) \leq c < (\log 2 + 1/2)c$ and $T(3) \leq c < (\log 3 + 1/2)c$, establishing the inductive hypothesis for the base cases.

Thus, we have $T(n) \leq c \log n + \frac{c}{2}$ for all $n \geq 2$, which implies that the solution to the recurrence is $T(n) = O(\log n)$.

7. Exercise 4.5-1 (a, b, d, e) (p106)

- (a) $T(n) = 2T(n/4) + 1$
 Choosing $\epsilon = 1 > 0$, $f(n) = O(n^{\log_4 2^{-1}})$.
 So, $T(n) = \Theta(\sqrt{n})$.
- (b) $T(n) = 2T(n/4) + \sqrt{n}$
 Because $f(n) = \sqrt{n} = \Theta(\sqrt{n})$, $T(n) = \Theta(\sqrt{n} \log n)$.
- (c) N/A
- (d) $T(n) = 2T(n/4) + n$
 Choosing $\epsilon = 2$, $f(n) = \Omega(n)$.
 Additionally, for $c = 1/2$ and $\forall n$, $cf(n) \geq af(n/b) \equiv n/2 \geq n/2$.
 So, $T(n) = \Theta(n)$.
- (e) $T(n) = 2T(n/4) + n^2$
 Choosing $\epsilon = 12$, $f(n) = \Omega(n^2)$.
 Additionally, for $c = 1/2$ and $\forall n$, $cf(n) \geq af(n/b) \equiv n^2/2 \geq n^2/2$.
 So, $T(n) = \Theta(n^2)$.

8. Exercise 4.5-2 (p106)

Given that Strassen's algorithm follows $T(n) = 7T(n/2) + \Theta(n^2)$ and that Caesar's algorithm follows $T(n) = aT(n/4) + \Theta(n^2)$, we must find $\log_4 a$ such that $a < \log_2 7 \cong 2.81$. This is true for $a \leq 48$, so $a = 48$ is the largest integer for which his algorithm could run faster than Strassen's.

9. Exercise 5.2-1 (p133)

The probability of hiring exactly one time is $\frac{1}{n}$ (this occurs on the best-case scenario of a forward-sorted list). The probability of hiring exactly n times is $\frac{1}{n!}$ (this occurs on the worst-case scenario of a reverse-sorted list).

10. Exercise 5.2-2 (p133)

The probability of hiring exactly twice is $\frac{x}{x}$.

11. (EC) Problem 4-6 (p122)