


Disease Knowledge Transfer across Neurodegenerative Diseases



Răzvan V. Marinescu, Marco Lorenzi, Stefano B. Blumberg, Alexandra L. Young, Pere Morell, Neil Oxtoby, Arman Eshaghi, Keir Yong, Sebastian Crutch, Polina Golland, Daniel C. Alexander


Aim Infer progression of multimodal biomarkers in rare neurodegenerative diseases (NDs) by leveraging larger datasets of common NDs.

Why Posterior Cortical Atrophy (PCA): progression of multimodal biomarkers not known → Identify outcome measures and suitable subjects for PCA clinical trials



1. Intuition

- Diseases affect different brain regions un-equally, but underlying mechanisms are the same (amyloid cascade)
- **Idea:** each brain region follows “its own disease course”, common across diseases → dysfunction score




2. Method

1. Each disease characterised by region-specific dysfunction trajectories

$$\gamma_{ij}^l = f(\beta_i + m_{ij}; \lambda_{d_i}^l)$$
2. Dysfunction trajectory modelled using region-specific biomarkers

$$y_{ijk} = g(\gamma_{ij}^k; \theta_k) + N(0, \epsilon_k)$$



3. Inference with belief propagation

while θ, λ, β not converged do

```

; // Estimate biomarker trajectories (disease agnostic)

$$\theta_k^{(u)} = \arg \min_{\theta_k} \sum_{(i,j) \in \Omega_k} \left[ y_{ijk} - g \left( f(\beta_i^{(u-1)} + m_{ij}; \lambda_{d_i}^{k,(u-1)}); \theta_k \right) \right]^2$$

; // Estimate dysfunction trajectories (disease specific)

$$\lambda_d^{l,(u)} = \arg \min_{\lambda_d^l} \sum_{(i,j,k) \in \Omega_{d,l}} \left[ y_{ijk} - g \left( f(\beta_i^{(u-1)} + m_{ij}; \lambda_d^l); \theta_k^{(u)} \right) \right]^2$$

; // Estimate subject-specific time shifts

$$\beta_i^{(u)} = \arg \min_{\beta_i} \sum_{(j,k) \in \Omega_i} \left[ y_{ijk} - g \left( f(\beta_i + m_{ij}; \lambda_{d_i}^{k,(u)}); \theta_k^{(u)} \right) \right]^2$$



```

4. Datasets

- Dementia Research Center cohort: MRI scans from 76 PCA, 67 tAD, 87 controls for training, 10 PCA with DTI for validation.
- TADPOLE dataset (ADNI) split into three subgroups with different progressions: 21 hippocampal, 35 cortical, 27 subcortical.

5. Results

- Inferred multimodal trajectories for PCA in lack of such data.
- Results are plausible, suggesting late-stage posterior damage.



- Our model has favourable performance compared to other models, on two different datasets.

Model	Cingulate	Frontal	Hippocam.	Occipital	Parietal	Temporal
TADPOLE: Hippocampal subgroup to Cortical subgroup						
DKT (ours)	0.56 ± 0.23	0.35 ± 0.17	0.58 ± 0.14	-0.10 ± 0.29	0.71 ± 0.11	0.34 ± 0.26
AD model	0.44 ± 0.25	0.34 ± 0.21	0.34 ± 0.24*	-0.07 ± 0.22	0.64 ± 0.16	0.08 ± 0.24*
Multivariate	0.60 ± 0.18	0.11 ± 0.22*	0.12 ± 0.29*	-0.22 ± 0.22	-0.44 ± 0.14*	-0.32 ± 0.29*
Spline	-0.24 ± 0.25*	-0.06 ± 0.27*	0.58 ± 0.17	-0.16 ± 0.27	0.23 ± 0.25*	0.10 ± 0.25*
Linear	-0.24 ± 0.25*	0.20 ± 0.25*	0.58 ± 0.17	-0.16 ± 0.27	0.23 ± 0.25*	0.13 ± 0.23*
typical Alzheimer's to Posterior Cortical Atrophy						
DKT (ours)	0.77 ± 0.11	0.39 ± 0.26	0.75 ± 0.09	0.60 ± 0.14	0.55 ± 0.24	0.35 ± 0.22
AD model	0.80 ± 0.09	0.53 ± 0.17	0.80 ± 0.12	0.56 ± 0.18	0.50 ± 0.21	0.32 ± 0.24
Multivariate	0.73 ± 0.09	0.45 ± 0.22	0.71 ± 0.08	-0.28 ± 0.21*	0.53 ± 0.22	0.25 ± 0.23*
Spline	0.52 ± 0.20*	-0.03 ± 0.35*	0.66 ± 0.11*	0.09 ± 0.25*	0.53 ± 0.20	0.30 ± 0.21*
Linear	0.52 ± 0.20*	0.34 ± 0.27	0.66 ± 0.11*	0.64 ± 0.17	0.54 ± 0.22	0.30 ± 0.21*

6. Conclusion

- Developed a novel methodology and model for transfer learning across different diseases
- Inferred multimodal trajectories for Posterior Cortical Atrophy

Weblinks

- Source code: <https://github.com/mrazvan22/dkt>
- Website: <https://people.csail.mit.edu/razvan/>

Funders

