

Neural correlates of the deployment of spatial attention, and their modulation by repetitive movements

Cameron Smith¹ Daniel H. Baker^{1,2}

¹Department of Psychology, University of York, UK

²Corresponding author, email: daniel.baker@york.ac.uk

Abstract

The deployment of spatial attention generates distinct neural signatures that can be detected at the scalp. Here, we use multivariate pattern analysis of EEG data to decode the deployment of spatial attention, and ask if this is modulated by repetitive movements. ‘Stimming’ movements (also known as repetitive stereotypies), are widely reported in autism, but also present in some neurotypical individuals. Stimming has historically been viewed as a problematic behaviour, but many individuals claim that stimming benefits attention. We first validated our paradigm (a Posner-style cueing design), demonstrating above-chance classification of cue direction from around 300ms post-cue onset. We then investigated whether stimming modulates decoding accuracy and task performance. Our results, consisting of data primarily from neurotypical participants, do not suggest that stimming has a negative impact on an individual’s ability to attend, unless the individual does not typically engage in stimming behaviours. This suggests interventions aiming to reduce stimming behaviours are not necessarily warranted and highlights the need for further research into the potential benefits of stimming specifically within the autistic population. Future research might also consider the potential overlap between autistic stimming and the fidgeting behaviours which are characteristic of ADHD, to help understand the significant overlaps between the characteristics of the two conditions.

Keywords: attention; MVPA; stimming; EEG

25 Introduction

26 The ability to orient and control the deployment of our attention to spatial locations or features
27 of our environment is essential to our ability to perform tasks, focusing on relevant stimuli
28 and disregarding those that are unnecessary for the task at hand [1–3]. A classic technique for
29 studying the deployment of attention in space is the Posner cueing paradigm, which demon-
30 strates that it is possible for attention to be deployed to spatial locations independently of eye
31 movements [4].

32 Individuals with neurodevelopmental disorders such as attention deficit hyperactivity disorder
33 (ADHD) and autism present with deficits in attention [5]. It has been suggested that repetitive
34 stereotypies, or ‘stimming’ behaviours, are linked to these deficits in attention – in the few
35 studies where they have been asked, individuals who stim report that it benefits their ability
36 to focus, and that the behaviours aid in emotional regulation [6,7]. It has been proposed that
37 an improved ability to focus through stimming may have a neural basis [8]. Building on these
38 theories and research, we tested the effect of stimming on performance in a Posner cueing
39 task.

40 The Posner cueing paradigm [4] involves the participant attending to a central fixation point,
41 then receiving a cue (which may be exogenous, where attention is captured through bottom-
42 up processes; or endogenous, where attention is controlled by top-down processes, such as
43 attending selectively to cues of the correct colour [9]) indicating that a stimulus is about to
44 appear either to the left or the right of the fixation point [10]. The assumption underlying
45 the task is that if participants successfully orient their attention to the cued location, then
46 their reaction times will be shorter and responses more accurate for congruent trials (where
47 the stimulus appears at the cued location) than for incongruent trials (where the stimulus
48 appears at the uncued location); this has been demonstrated extensively [10]. The reaction
49 time difference was found by Posner [11] to be 25ms faster for congruent trials and 40ms slower
50 for incongruent trials, compared to a neutral cue condition.

51 This paradigm has been combined with EEG to investigate the neural basis of spatial atten-
52 tion. For example, Landau et al. [12] investigated gamma-band activity during a Posner cueing
53 task, comparing congruent and incongruent trials. Gamma range responses were associated
54 with voluntary shifts in attention, e.g., attending to the left of the fixation point following
55 a left-pointing arrow cue, but were absent when attention was involuntarily captured by the
56 appearance of a stimulus in the uncued location on incongruent trials; this suggests that volun-
57 tary and involuntary processes of attention deployment are controlled by separate mechanisms.
58 Thierry et al. [10] investigated two specific ERPs, the N2pc component (a negative response
59 typically seen at around 200ms post stimulus, in the posterior-contralateral region) and the
60 SPCN component (sustained posterior contralateral negativity). They used multivariate pat-
61 tern analysis and used the classifier accuracy to determine that these two components were
62 able to predict the focus of participants’ spatial attention. This was modulated by the varied
63 distance of the stimulus from the central fixation (further being better decoded).

64 A combined EEG and TMS study by Capotosto et al. [13] demonstrated that the applica-
65 tion of TMS to the right intraparietal sulcus (IPS) impairs target detection, particularly on
66 incongruent trials. They additionally noted that the P3 response amplitude was affected by
67 this; on incongruent trials its amplitude was significantly lower, and on congruent trials it was
68 significantly higher. The IPS has been previously identified as an important region in atten-
69 tional deployment by Vossel et al. [14] in an fMRI study identifying regions that are selectively
70 active on congruent and incongruent trials. The right IPS and, additionally, the right inferior
71 frontal gyrus were implicated in attentional processes related to both types of trial. Peelen
72 et al. [15] concluded that both endogenous and exogenous orienting cues involved the same
73 network in their fMRI study, implicating frontal and parietal regions (the premotor cortex,
74 posterior parietal cortex, medial frontal cortex, and right inferior frontal cortex). Fitzgerald
75 et al. [16] conducted an MRI study using the Posner cueing task and found that activation in
76 the ventral attention network differed between individuals with autism and controls; individ-
77 uals with autism also showed weaker functional connectivity in the dorsal attentional network
78 compared to their neurotypical counterparts.

79 A number of developmental disorders involve deficits in attention, notably ADHD and autism.
80 In the case of ADHD, attention deficits are a defining feature of the condition (American
81 Psychological Association, 2013). For autism, attention deficits are not an explicit part of the
82 diagnostic criteria (although restricted interests, which are included, are somewhat related),
83 but both attentional deficits and strengths are widely reported in autistic individuals. They
84 are typically more distractible, demonstrating ‘underselective’ attention, but also demonstrate
85 ‘overselective’ attention where they attend intensely to a more limited subset of stimuli. This
86 ‘overselective’ attention has been suggested to be linked to restricted interests, which is also
87 sometimes referred to as ‘hyperfocus’, where a task is the subject of intense focus for the
88 individual [e.g., 17]. These differences appear to have a neural basis – they have been linked to
89 unusually wide sulci in the parietal lobe [5]. Hyperfocus is also widely self-reported in ADHD,
90 despite a lack of clinical research into the phenomenon [18].

91 These marked differences in attentional deployment seen in both conditions are worth consider-
92 ing in the framework of the transdiagnostic approach, which seeks to identify core deficiencies
93 that result in the presentation of traits from multiple disorders, instead of approaching differ-
94 ent disorders as entirely separate even within the same individual [19]. ADHD is the most
95 common comorbidity of autism, and even without a dual diagnosis, 30-80% of individuals with
96 autism present symptoms of ADHD, and 20-60% of children with ADHD exhibit autism-like
97 traits [20]. Some shared deficiencies that have already been identified in the case of autism and
98 ADHD include fine motor function and verbal fluency, alongside some structural differences
99 in the brain [20].

100 Stimming behaviours are commonly seen in individuals with diagnoses of autism and/or
101 ADHD, and involve repetitive movements, typically featuring the arms, hands, or entire body
102 [21], for example hand flapping [6]. They are also observed in neurotypical individuals, but at
103 much lower rates than in neurodiverse individuals [21]. The behaviours are typically defined
104 as involuntary, but pleasant [6,7,21], and are associated with emotional states such as stress,

boredom, concentration, or excitement [22]. They are often defined as purposeless [e.g., 23] but self-reports from individuals with autism claim that stimming aids the regulation of their emotions [6,7].

A type of vocal stimming called echolalia has been consistently demonstrated to have benefits for autistic children in the acquisition of language [24,25]. This is one of the few instances that stimming has been researched in terms of its potential function. Echolalic utterances, which typically involve repeating words spoken by another, have also been identified as having predominantly communicative functions [26]. This opens up an avenue to research potential benefits that other types of stimming may provide.

Current research into stimming is largely focussed on designing interventions to eliminate the behaviours, for example the use of weighted vests [27]. Taking this approach without first attempting to understand if there are benefits to the behaviour seems ill-advised. McCarty and Brumback [8], for example, proposed based on self-reports of the benefits of stimming (focus, coping with overwhelming sensory input, and relaxation) that regular motor movements may generate (or be a by-product of) rhythmic oscillations in the motor cortex, which entrain oscillations in the sensory cortex. This entrainment might improve information transfer in the sensory cortex, effectively normalising the atypical motor and sensory oscillations observed in the brains of autistic individuals both during tasks [28–30] and at rest [31,32].

There is heavy stigma around stimming behaviours, with many individuals reporting they have been explicitly instructed not to stim, despite attempts at repression causing them some discomfort [7]. Individuals with autism state that they believe stimming should not be stigmatised in this way, and that they should be allowed to act in the way that feels natural to them [6]. Research into potential benefits of stimming is important as a step towards dispelling this stigma, and discouraging interventions that aim to eliminate the behaviour.

This study involved two experiments. In the first experiment, participants completed a Posner cueing task while EEG recordings were taken; a pattern classifier then attempted to differentiate, using the participants’ brain activity, whether they attended to the left vs. to the right following the cue onset. This is similar to the methodology used by Thiery et al. [10]: by measuring the timecourse of classifier accuracy, we are provided with an index of the deployment of spatial attention. The second experiment extended this paradigm by adding a stimming condition. The accuracy of the classifier was compared between conditions where the participants were permitted to stim during the trials, or instructed to keep still. In this case, we used decoding accuracy to assess whether stimming provided a benefit in terms of the deployment of spatial attention; if there was a benefit, the classifier accuracy would be higher, as the EEG signal would differ more between conditions due to the greater focus of attention in different spatial locations. For the purposes of our analyses, due to an insufficiently sized and heterogeneous clinical group, we grouped the participants based on whether or not they reported stimming in their everyday lives; as discussed previously, neurotypicals often engage in stimming [21], and 64% of our participants reported in the pre-experiment questionnaire that they typically engaged in stimming.

Methods

Participants

41 participants took part in Experiment 1, but four produced very noisy data so were excluded, leaving a sample of 37 participants (20 male, 17 female). A total of 22 participants took part in Experiment 2 (8 male, 14 female). Participants in Experiment 2 were asked to report diagnoses of neurodevelopmental disorders – 17 had no diagnosis, 2 had a diagnosis of autism, and 3 had a diagnosis of ADHD. 15/22 participants further reported they typically engaged in stimming behaviours, while 7 reported that they did not. Both experiments were granted ethical approval by the Ethics Committee of the Department of Psychology at the University of York. The identification numbers were 356 for Experiment 1, and 173 for Experiment 2. Participants provided written informed consent before data collection began.

Apparatus and stimuli

Stimuli were displayed on a ViewPixx 3D display (VPixx Technologies Inc., Quebec, Canada) with a diagonal extent of 24 inches, a refresh rate of 120Hz, and a resolution of 1920x1080 pixels. The screen was viewed by the participants from a distance of 57cm, resulting in a spatial resolution of 36 pixels per degree of visual angle. The display was gamma corrected using a Minolta LS110 photometer to ensure a linear luminance response. Stimuli were generated and displayed using a Mac Pro computer running Matlab, with the experiment script making use of functions from Psychtoolbox 3 [33–35].

In both experiments, EEG recordings were acquired using an ANT Neuroscan 64-channel EEG system (ANT Neuro, Hengelo, Netherlands). Electrodes were mounted in a waveguard cap, arranged according to the 10/20 system, and were referenced to the whole head average. The ground electrode was located at position *AFz*, and EEG signals were recorded at 1kHz. In Experiment 2, participants' eye movements were also recorded using an EyeLink 1000 (SR Research Ltd., Ottawa, Canada) eye tracker, also sampling at 1kHz. Low-latency digital triggers were sent from the stimulus computer to the EEG system using an 8-bit parallel cable.

Target stimuli in both experiments were horizontal sine-wave gratings with a diameter of 4 degrees, and a spatial frequency of 2c/deg. The target appeared offset by 8 degrees to either the left or the right of the central fixation point. The base contrast of the target was 50%, and either the upper or lower half of the grating contained a contrast increment of a further 10%. Participants were asked to indicate whether the upper or lower half of the grating contained the increment using arrow keys on the computer keyboard. The fixation cross was 0.25×0.25 degrees, and was displayed throughout.

Each trial was preceded by a cue. In Experiment 1 the cue was a face with eyes pointing left or right. In Experiment 2 the cue was an arrow pointing left or right. The face cues used in

Experiment 1 were the average of 22 female faces taken from the NimStim database [36] and were digitally altered so that the eyes were directed to the left or the right. The face had a width of 3 degrees and a height of 4 degrees and was displayed in the centre of the screen. The arrow cue used in Experiment 2 was 1.4 degrees long and was displayed in the centre of the screen.

Procedure

Participants in both experiments completed four blocks of the Posner cueing task, with each block consisting of 200 trials. They were instructed to keep their eyes focused on the central fixation cross, and to attend to the gratings without shifting their gaze. During each trial, participants viewed a fixation cross, which remained on the screen throughout the trial (excluding times when the cue appeared on the screen). The cue stimulus appeared for 200ms, followed by a randomly determined gap of 400-600ms before the grating appeared on the screen. The target grating was then displayed for 200ms. Participants were required to report whether the upper or lower half of the grating appeared higher in contrast; following their response, a randomly determined interval drawn from a normal distribution (average 1000ms, with a standard deviation of ± 200 ms) preceded the next appearance of the cue. This procedure is illustrated in Figure 1 for the face cue (panel a) and the arrow cue (panel b). Cue congruence was 75% in both experiments, meaning that each participant contributed a total of 600 congruent trials and 200 incongruent trials across the experiment.




Figure 1: Illustration of trial procedure for both experiments. In panel (a), the cue was a face with eyes pointing to either the left (as pictured) or the right. In panel (b), the cue was an arrow. Target gratings had a contrast of 50%, plus a 10% increment in either the upper or lower half. Note that images are not drawn to scale in this schematic to aid visibility - see text for actual dimensions.

200 In Experiment 2, participants were additionally instructed to stim, using a movement restricted
201 to one hand only, in half of the blocks. This was to ensure minimal distortion of the EEG
202 signal resulting from movement. The order in which participants completed stimming or non-
203 stimming blocks was counterbalanced; half the participants were instructed to stim on blocks
204 1 and 3, while the other half were instructed to stim on blocks 2 and 4.

205 Data analysis

206 Raw EEG data were converted into EEGLab .set format using Matlab and the EEGLab toolbox
207 [37]. We then used MNE python [38] for the main analysis. Data were bandpass filtered be-
208 tween 0.2 and 30Hz, and then epoched using the event timestamps. We performed classification
209 using a linear classifier with 10-fold cross-validation on the voltage data across all electrodes
210 (excluding electrodes *M1* and *M2*), independently at each time point and for each participant.
211 We then averaged classification accuracy across participants to generate the timecourses used
212 in the main analysis. We calculate Bayes factors [39] to compare classification accuracy to
213 chance performance (50% correct) and to make comparisons between conditions and groups,
214 but also report frequentist tests for reference. We express Bayes factors in logarithmic units
215 to account for extreme values ($\log_{10}BF_{10}$), such that a Bayes factor of 3 is a $\log_{10}BF_{10} = 0.48$,
216 and constitutes convincing evidence for the alternative hypothesis, a Bayes factor of 10 is a
217 $\log_{10}BF_{10} = 1$, and constitutes strong evidence for the alternative hypothesis, and a Bayes
218 factor of 30 is a $\log_{10}BF_{10} = 1.48$, and constitutes overwhelming evidence for the alternative
219 hypothesis.

220 Data and code availability

221 Raw data, processed data, and analysis scripts are freely available through the project reposi-
222 tory at: <http://doi.org/10.17605/OSF.IO/YUHZ4>.

223 Results

224 Experiment 1

225 We first analysed the behavioural responses from the experiment, comparing accuracy and
226 reaction times between stimuli shown in the cued (i.e. a congruent target) and uncued (an
227 incongruent target) location. There was a highly significant effect of cue congruency on both
228 accuracy ($\log_{10}BF_{10} = 1.77$, $t = 3.82$, $p < 0.001$, $d = 0.61$) and reaction time ($\log_{10}BF_{10} =$
229 6.08 , $t = -7.26$, $p < 0.001$, $d = 1.16$), as shown in Figure 2. Responses to targets in the cued
230 location were both faster ($M_{diff} = 51$ ms) and more accurate ($M_{diff} = 3\%$) than for those in
231 the uncued location, showing a classic Posner cueing effect.




Figure 2: Summary of reaction time and accuracy data for the cueing task in Experiment 1. Small points show data for individual participants ($N = 39$), and larger points give the group means. Error bars indicate 95% confidence intervals.

We next explored the EEG data in response to both the cue presentation and the target presentation. There were typical evoked potentials in response to both types of stimulus, as summarised in Figure 3b,c,e,f. Patterns of voltage across the scalp showed similar lateralisation effects both for cues pointing to either the left or right, and for targets presented to either the left or right. Multivariate pattern analysis was able to decode the side of cue presentation from around 300ms after cue onset (see Figure 3a), with a maximum accuracy of 69%, and a plateau of high performance from around 500-1000ms. This relatively late onset of above-chance classification likely reflects the deployment of covert spatial attention to one side of the visual field, and is therefore a neural index of attentional focus. The pattern classifier could decode the side of target presentation with much greater accuracy (maximum of 89%), from around 100ms after target onset (black curve in Figure 3d). Finally, we were able to decode cue congruency (i.e. whether the target presented was on the side indicated by the cue) with a maximum accuracy of 62%, and a similar timecourse to the decoding of target side (green curve in Figure 3d). Overall, these results indicate that our EEG data are sufficiently sensitive to provide an index of the deployment of spatial attention.

Experiment 2

In Experiment 2, we replicated the basic design of Experiment 1, but this time with an added ‘stimming’ condition, and with participants divided into groups based on their self-reported stimming behaviour. The results are summarised in Figure 4. Pooling across all participants, for response accuracy there was no effect of cue congruency ($\log_{10}BF_{10} = -0.66$, $F = 0.06$, $p = 0.803$, $\eta_G^2 = 0$), stimming condition ($\log_{10}BF_{10} = -0.66$, $F = 0$, $p = 0.977$, $\eta_G^2 = 0$), nor an interaction between the two ($\log_{10}BF_{10} = -0.47$, $F = 0.8$, $p = 0.383$, $\eta_G^2 = 0$). For reaction time, there were significant effects of cue congruency ($\log_{10}BF_{10} = 3.11$, $F = 29.28$, $p < 0.001$, $\eta_G^2 = 0.07$) and stimming condition ($\log_{10}BF_{10} = 1.09$, $F = 7.1$, $p = 0.01$, $\eta_G^2 = 0.04$), as well as an interaction between the two ($\log_{10}BF_{10} = -0.34$, $F = 12.03$, $p = 0.002$, $\eta_G^2 = 0.01$). Note that reactions were faster in the non-stimming condition (591 ms) than in the stimming condition (632 ms). These effects remained significant in the group who reported stimming in their daily lives (N=14) for the effects of cue congruency ($\log_{10}BF_{10} = 1.46$, $F = 11.42$, $p < 0.005$, $\eta_G^2 = 0.06$), stimming condition ($\log_{10}BF_{10} = 0.42$, $F = 4.89$, $p = 0.045$, $\eta_G^2 = 0.03$), and their interaction ($\log_{10}BF_{10} = -0.32$, $F = 8.71$, $p = 0.011$, $\eta_G^2 = 0.01$). For the group who did not report stimming (N=8), only the effect of cue congruency was significant ($\log_{10}BF_{10} = 0.93$, $F = 40.15$, $p < 0.001$, $\eta_G^2 = 0.11$).

The EEG results from Experiment 2 are summarised in Figure 5. Panels (a,b) demonstrate that multivariate pattern analysis is able to decode cue direction, target side, and target congruency with similar timecourses to Experiment 1 (see Figure 3), though with somewhat lower accuracy overall. We then compared decoding accuracy for cue side and cue congruency between the stimming group (N=14), and the non-stimming group (N=8). Although decoding accuracy appeared to be slightly higher for the non-stimming group for cue side (Figure 5c), this difference was not convincing. Accuracy for cue congruency was very similar for the




Figure 3: Summary of EEG data from Experiment 1. Panel (a) shows decoding accuracy of a pattern classification algorithm trained on responses to the cue stimulus (Time 0 is the cue onset). ERPs and scalp topographies are shown for cues pointing left (b) and right (c), with the shaded regions indicating the time period over which voltages were averaged to generate the scalp plots. Panel (d) shows decoding accuracy for classifying target location (black curve) or cue congruency (green curve) in response to the target (Time 0 is the target onset). ERPs and scalp topographies are shown for targets on the left (e) and right (f) side of fixation, with the shaded regions indicating the time period over which voltages were averaged to generate the scalp plots. Shaded regions in panels (a,d) indicate 95% confidence intervals, and lines above $y=1$ indicate Bayes factor scores above 3 (yellow >3 ; orange >10 ; red >30).




Figure 4: Summary of reaction time and accuracy data for the cueing task in Experiment 2. Small points show data for individual participants ($N = 22$), and larger points give the group means. Error bars indicate 95% confidence intervals.

two groups (Figure 5d). Finally, we compared the deployment of spatial attention between stimming and non-stimming conditions for each of the two groups. There appeared to be a slight decrease in decoding accuracy in the non-stimming group when they were asked to stim (Figure 5f), but no differences in the stimming group (Figure 5e).

We collected eye tracking data for a subset of the participants in Experiment 2 (16/22; for the remaining participants, the eye tracker was not able to calibrate successfully). We used these data to check if participants were able to follow the task instructions and fixate centrally throughout. The results are summarised in Figure 6. The grey distributions indicate horizontal fixation positions at the time of cue onset. For most participants these are close to the central fixation (0 deg, see vertical dashed line), though there are some notable consistent offsets that we attribute to calibration errors (e.g. P201, P223, P224). The coloured distributions indicate horizontal fixation positions at the time of target onset, following either a leftward (blue) or rightward (red) pointing cue. The majority of participants were able to follow the instructions and fixated centrally for both of these conditions, showing no systematic differences. One participant (P214) clearly made eye movements in the cued direction on most trials, and some participants (e.g. P209, P216, P218) showed evidence of sometimes making such eye movements. At a group level, there was a marginally significant difference between the mean horizontal positions for left vs right cues ($t = 2.15$, $p < 0.05$), but the overall evidence was not compelling ($\log_{10}BF_{10} = 0.19$). We therefore conclude that in general participants were able to follow instructions appropriately, and that eye movements are unlikely to be responsible for




Figure 5: Summary of EEG data from Experiment 2. Panel (a) shows decoding accuracy of a pattern classification algorithm trained on responses to the cue stimulus (Time 0 is the cue onset) for the full sample of participants ($N=22$). Panel (b) shows decoding accuracy for classifying target location (black curve) or cue congruency (green curve) in response to the target (Time 0 is the target onset), again for all participants. Panels (c,d) compare decoding between the stimulating ($N=14$) and non-stimulating ($N=8$) participants for cue side and cue congruency. Panels (e,f) compare decoding accuracy for cue side within each group, and between the stimulating and no stimulating conditions. Shaded regions in all panels indicate 95% confidence intervals, and lines above $y=1$ indicate Bayes factor scores above 3 (yellow >3 ; orange >10 ; red >30).

291 the EEG effects of cueing.

292 Discussion

293 Overall, our results from Experiment 1 support the assertion that EEG data can be used
294 as a metric of attentional deployment [10]. The classifier was able to differentiate both cue
295 and target side from EEG recordings in Experiment 1, indicating we have recorded a neural
296 correlate of attentional deployment, and that it is possible for a machine learning classifier
297 to determine the direction of spatial attention using EEG signals. Extending this method in
298 Experiment 2 by including the stimming manipulation suggested that stimming does not have
299 a detrimental effect on attentional deployment, with our only negative impact of stimming
300 being on reaction times. We will now discuss these results in greater detail, with a focus on
301 the implications for research into stimming in the context of the neurodiversity movement.

302 Behavioural data from Experiment 1 revealed a significant effect of congruency on both ac-
303 curacy and reaction time, a typical Posner cueing effect wherein higher accuracy and shorter
304 reaction times are associated with congruent relative to incongruent conditions [10,11]. Ex-
305 periment 1 further involved testing the classifier, which revealed that cue side was reliably
306 decoded from 300ms post-stimulus onset onwards, reaching an accuracy level of 69%, and tar-
307 get side was reliably decoded from much earlier (100ms post-stimulus onset) and reached 89%
308 accuracy. This sufficiently validated the use of the measure in Experiment 2 and supported
309 prior research using similar techniques [10].

310 Experiment 2 replicated this methodology with the addition of the stimming manipulation.
311 Additionally, eye position data were recorded where possible, allowing us to confirm that in
312 the majority of cases participants understood and followed the instructions, keeping their eyes
313 fixated on the centre of the screen, and did not overtly move them following the cue. This
314 confirms that the neural correlates we measured do not reflect eye movements, suggesting they
315 are instead a genuine representation of the deployment of spatial attention.

316 Analysis of the behavioural data revealed no significant effect of either stimming or congruency
317 on accuracy. The effect of congruency on accuracy in Posner cueing tasks is robust [10,11] and
318 our version of the task generated this effect in Experiment 1, suggesting an external factor
319 resulted in this difference in Experiment 2. For example, the environmental conditions under
320 which participants completed the experiment were quite different, as data for Experiment
321 1 were collected over the winter, while Experiment 2 was conducted in the summer, when
322 the laboratory was very warm, potentially reducing participants' ability to focus on the task.
323 Accuracy was lower overall for Experiment 2 than Experiment 1; participants in Experiment
324 1 averaged 96% correct for congruent trials and 93% correct for incongruent trials, while
325 participants in Experiment 2 averaged 86% correct for both congruent trials and incongruent
326 trials. Difficulties resulting from external factors such as temperature and general comfort
327 may have influenced the participants' ability to attend to the experiment to the point where
328 differences in accuracy became non-significant. Our finding that stimming had no effect on




Figure 6: Distributions of horizontal eye positions for 16 participants during Experiment 2. Grey distributions represent eye positions at cue onset. Coloured distributions represent eye positions at target onset following a leftward pointing cue (blue) or a rightward pointing cue (red). Dotted vertical lines indicate the target locations (± 8 degrees).

accuracy would therefore need to be validated through future replication; should it hold, it is an important one, as it suggests that stimming does not have a negative impact on individuals' performance, countering the notion that stimming behaviours are detrimental and should be targeted through interventions [27,e.g., 40]. These findings do not fully align with self-reports from autistic individuals, however, who report that stimming benefits attention [6]; accuracy was no greater in stimming conditions than non-stimming conditions. This may result from the proportion of our participants with autism being very small, so future research should aim to further investigate this in a larger sample. It has been suggested that stimming acts are a compensatory mechanism [6] for the previously discussed attention problems associated with autism [5], so there is a possibility that positive effects of stimming are only seen in these individuals, which warrants further investigation.

Reaction times for participants in Experiment 2 were also longer than for those in Experiment 1, again suggesting the participants experienced greater difficulty when completing the task under the external conditions of the second experiment. In Experiment 1, the average reaction time was 484ms for congruent trials and 535ms for incongruent trials, while in Experiment 2, the average reaction time was 557ms/611ms for congruent trials and 627/654ms for incongruent trials (for non-stimming/stimming conditions, respectively). However, analysis of the reaction time data did reveal a typical Posner cueing effect in regard to congruency: congruent trials were associated with shorter reaction times. Analysis also suggested a negative effect of stimming on reaction times, with reaction times being significantly slower when participants were stimming. Given the lack of impact observed on accuracy (and on EEG decoding, discussed next) it is likely this finding relates to stimming interfering with the participants' ability to complete the task (they were required to respond to stimuli with buttons on the keyboard while moving their other hand) rather than stimming being inherently distracting. When we further divided participants into those who reported that they stim in their everyday lives and those who reported that they did not, we found the pattern only persisted in the former group; this may be due to the much smaller size of the latter group, as for the former group the Bayes factor and effect size were smaller for the effect of stimming than the effect of congruency, suggesting the relationship between stimming and reaction time is weaker than between congruency and reaction time. In a larger group of individuals who do not typically engage in stimming, this relationship might persist. Future research should aim to recruit larger sample sizes for various neurotypes and everyday stimming behaviours. It should also aim to adapt tasks so that there is no potential for the physical motion of stimming to interfere with the physical motion of completing a task, allowing the pure effects of stimming to be investigated.

Classifier accuracy in Experiment 2 was lower overall than in Experiment 1, which was not unexpected given the smaller sample size and the fact that participants in Experiment 2 were moving, potentially introducing extra noise to the EEG signal. Within Experiment 2, classifier accuracy was largely similar between individuals who reported typically stimming and those who did not, validating comparisons between the groups based on classifier accuracy.

A decrease in classifier accuracy during stimming was observed exclusively in the group who

reported that they did not typically stim. The same was not true for the typically stimming group. We consider classifier accuracy here to be a measure of the strength of an individual's ability to deploy their spatial attention, as stronger attentional deployment to the left or right should result in a greater condition-based difference in neural activity, making it easier for the classifier to differentiate the direction of their attention. Therefore, this suggests an individual's ability to deploy their spatial attention is only reduced when they are asked to stim if they do not typically engage in the behaviour, contesting the notion that stimming is an inherently harmful behaviour. Further, it may give insight into where this concept originated; for individuals who do not typically stim, it appears to have negative consequences for their attentional abilities, which may lead them to assume that the same is true for everyone (which our evidence suggests is not the case). For individuals who typically engage in stimming behaviours, there does not appear to be an impact of stimming or not stimming on their attentional deployment, indicating that the behaviour is not detrimental; greater inclusion of neurodiverse individuals may allow for observation of a positive effect, as stimming has been proposed as a compensatory mechanism [6].

It has long been claimed that stimming behaviours interrupt attention and learning, which has been used to justify therapies aiming to reduce these behaviours [e.g., 40]. Our investigation contests this view, given our lack of evidence to suggest a negative impact of stimming on attention. The evidence often used to support this view of stimming behaviours falls foul to a host of methodological issues, such as incredibly small groups of participants [41], stopping children stimming entirely during the learning phase [42], or not attempting to prevent stimming behaviours at all [43], thereby providing no real evidence for whether or not stimming affects learning abilities, or whether reducing them would have a beneficial impact. Indeed, self-reports from autistic individuals that the behaviours benefit their attention may mean this view is leading autism interventions in entirely the wrong direction when it comes to improving the quality of life of autistic individuals. Interventions that aim to reduce stimming are worryingly widespread, given their weak research base. Behavioural approaches, particularly common in North America [44], frequently target these behaviours [e.g., 6,45]. Future research should attempt to take a similar approach to ours, investigating and allowing for the potential that stimming behaviours may have positive effects, or even no effect at all, to ensure we capture the full picture of stimming and its potential functions before encouraging interventions that target the behaviour.

One factor that future research should take into consideration is that in our investigation, we directed participants to stim or refrain from stimming at certain times. It is unclear whether there is a subjective difference in stimming when it is genuinely self-generated compared to when it is externally prompted. This could explain the lack of a positive effect of stimming that we identified here, if only the truly self-generated form of stimming produces attentional effects. A careful balance needs to be struck between controlling experimental conditions and ensuring stimming behaviours are authentic to best understand them. Potentially, future research could benefit from a more naturalistic design, in which participants complete a task while stimming when it feels natural to them, with researchers recording start and end times of stimming periods to create stimming and non-stimming conditions. Likely the ideal method of

dealing with this potential conflict is for researchers across the field to conduct investigations into stimming in a number of different ways, exploiting the strengths and weaknesses of various methodologies to obtain a full picture of stimming. It would also be beneficial to ask individuals who stim directly about whether or not they believe externally prompted stimming benefits them in the same way as self-directed stimming.

Our investigation intended to lay the groundwork for an investigation into whether stimming behaviours in autism have the same underlying functions and mechanisms as fidgeting in ADHD. Given our limited sample size of participants with either disorder, we have not gathered sufficient evidence to suggest whether or not this is the case. However, we believe this route is a promising one for investigation under the framework of the transdiagnostic approach, especially given that there are overlaps in the benefits of stimming and fidgeting as reported by individuals with autism and ADHD. Kapp et al. [6] reported autistic individuals stated stimming benefitted their ability to focus, while Canela et al. [46] similarly reported that individuals with ADHD stated fidgeting aided in their productivity and concentration. A greater understanding of this behaviour across conditions may further our understanding of the similarities between autism and ADHD, a pressing research interest given the high comorbidity of the two conditions, with the current and lifetime prevalence rates of ADHD in the autistic population being estimated at 38.5% and 40.2% respectively [47].

Future research should additionally pursue other avenues of exploration into the potential benefits of stimming. In this case, we focused on attention, given that this is a commonly reported area in which stimming generates benefits in individuals with autism [6] and there is crossover with fidgeting in ADHD, which is also reported to benefit attention [46]. However, autistic individuals report a number of other benefits of stimming, for example emotional regulation [6,7]. In order to obtain a full picture of the mechanisms and functions of stimming, to fully understand the behaviour and thus inform interventions which aim to reduce the behaviour, future research should investigate all reported avenues in which stimming may benefit.

Given that our investigation did not uncover a positive effect of stimming on attention, it could be suggested that interventions aiming to reduce stimming behaviours are not causing harm. Often these interventions are additionally justified on the basis that children who stim may be socially excluded or bullied as a result of their stimming behaviours [e.g., 48], and engaging in these behaviours can increase the likelihood of autistic children experiencing bullying [49], which is a serious problem. However, even if we assume stimming has no positive effect (despite self-reports from autistic individuals saying it does), efforts may be better focused on educating an autistic child's neurotypical teachers and peers on autism and stimming, cracking down on bullying, and teaching autistic children how to handle bullying. It has been demonstrated that describing and explaining autistic behaviours to children can improve their attitudes towards an autistic child [50], and a recent intervention teaching autistic children how to stand up to bullies achieved a reasonable level of success [49]. Autistic children may experience bullying as a result of a wide variety of their autistic traits [51]; surely, then, the best approach is to address the bullying, rather than encouraging autistic children to alter a wide variety of their own

behaviours to avoid it. Particularly in the case of stimming, autistic adults protest against the elimination of these behaviours [6]. Even though we have been unable to demonstrate in a controlled setting that these behaviours have benefits, it is important to listen to the community we aim to help and respect their wishes.

In summary, our results do not support prior literature suggesting stimming is a behaviour with negative consequences for attention. Rather, we find that in individuals who typically engage in stimming behaviours there is no particular detriment associated with doing so (excluding delayed reaction times, which potentially are an artefact of the experimental design). Negative impacts of stimming on attentional deployment were identified in individuals who do not typically engage in stimming behaviours, which may help explain why the perception has previously been that the behaviours are detrimental. This research had a predominantly neurotypical sample, and as such it is vital for future research to include a greater sample of autistic individuals to investigate whether stimming functions in the same way for this group as it appears to for neurotypicals who stim. Furthermore, the inclusion of a greater sample of individuals with ADHD may help uncover whether the fidgeting behaviours seen in ADHD may be similar to, or the same as, autistic stimming. Future research should also aim to investigate the potential benefits of stimming in a wider range of areas, given that individuals with autism also report benefits to emotional regulation [6] in addition to attention.

Declarations

Funding

CS was supported by a Bradshaw-Eagle Undergraduate Research Scholarship from the Applied Vision Association. DHB was supported by BBSRC grant BB/V007580/1.

Competing interests

The authors have declared that no competing interests exist.

Data availability statement

Raw data, processed data, and analysis scripts are freely available through the project repository at: <http://doi.org/10.17605/OSF.IO/YUHZ4>.

Author contributions

CS: conceived the study, collected the data, performed the analysis, wrote the paper.

DHB: conceived the study, contributed data, performed the analysis, wrote the paper.

References

1. Knudsen EI. Fundamental components of attention. *Annu Rev Neurosci.* 2007;30: 57–78. doi:[10.1146/annurev.neuro.30.051606.094256](https://doi.org/10.1146/annurev.neuro.30.051606.094256)
2. Hopfinger JB, Buonocore MH, Mangun GR. The neural mechanisms of top-down attentional control. *Nat Neurosci.* 2000;3: 284–91. doi:[10.1038/72999](https://doi.org/10.1038/72999)
3. Rueda MR, Posner MI, Rothbart MK. 15: Attentional control and self-regulation. 2nd ed. In: Vohs KD, Baumeister RF, editors. *Handbook of self-regulation*. 2nd ed. Guilford Press; 2011. pp. 284–299.
4. Posner MI. Orienting of attention. *Q J Exp Psychol.* 1980;32: 3–25. doi:[10.1080/00335558008248231](https://doi.org/10.1080/00335558008248231)
5. Allen G, Courchesne E. Attention function and dysfunction in autism. *Front Biosci.* 2001;6: D105–19. doi:[10.2741/allen](https://doi.org/10.2741/allen)
6. Kapp SK, Steward R, Crane L, Elliott D, Elphick C, Pellicano E, et al. 'People should be allowed to do what they like': Autistic adults' views and experiences of stimming. *Autism.* 2019;23: 1782–1792. doi:[10.1177/1362361319829628](https://doi.org/10.1177/1362361319829628)
7. Steward R. Repetitive stereotyped behaviour or “stimming”: An online survey of 100 people on the autism spectrum. *INSAR conference*; 2015. Available: <https://insar.confex.com/insar/2015/webprogram/Paper20115.html>
8. McCarty MJ, Brumback AC. Rethinking stereotypes in autism. *Seminars in Pediatric Neurology.* 2021;38: 100897. doi:<https://doi.org/10.1016/j.spen.2021.100897>
9. Leblanc E, Prime DJ, Jolicoeur P. Tracking the location of visuospatial attention in a contingent capture paradigm. *J Cogn Neurosci.* 2008;20: 657–71. doi:[10.1162/jocn.2008.20051](https://doi.org/10.1162/jocn.2008.20051)

- 493 10. Thiery T, Lajnef T, Jerbi K, Arguin M, Aubin M, Jolicoeur P. Decoding the locus of covert visuospatial attention from EEG signals. *PLoS One*. 2016;11: e0160304. doi:[10.1371/journal.pone.0160304](https://doi.org/10.1371/journal.pone.0160304)
- 494 11. Posner MI. *Chronometric explorations of mind*. Lawrence Erlbaum; 1978.
- 495 12. Landau AN, Esterman M, Robertson LC, Bentin S, Prinzmetal W. Different effects of voluntary and involuntary attention on EEG activity in the gamma band. *Journal of Neuroscience*. 2007;27: 11986–11990.
- 496 13. Capotosto P, Corbetta M, Romani GL, Babiloni C. Electrophysiological correlates of stimulus-driven reorienting deficits after interference with right parietal cortex during a spatial attention task: A TMS-EEG study. *J Cogn Neurosci*. 2012;24: 2363–71. doi:[10.1162/jocn_a_00287](https://doi.org/10.1162/jocn_a_00287)
- 497 14. Vossel S, Weidner R, Thiel CM, Fink GR. What is "odd" in posner's location-cueing paradigm? Neural responses to unexpected location and feature changes compared. *J Cogn Neurosci*. 2009;21: 30–41. doi:[10.1162/jocn.2009.21003](https://doi.org/10.1162/jocn.2009.21003)
- 498 15. Peelen MV, Heslenfeld DJ, Theeuwes J. Endogenous and exogenous attention shifts are mediated by the same large-scale neural network. *NeuroImage*. 2004;22: 822–830. doi:<https://doi.org/10.1016/j.neuroimage.2004.01.044>
- 499 16. Fitzgerald J, Johnson K, Kehoe E, Bokde ALW, Garavan H, Gallagher L, et al. Disrupted functional connectivity in dorsal and ventral attention networks during attention orienting in autism spectrum disorders. *Autism Res*. 2015;8: 136–52. doi:[10.1002/aur.1430](https://doi.org/10.1002/aur.1430)
- 500 17. Dupuis A, Mudiyansele P, Burton CL, Arnold PD, Crosbie J, Schachar RJ. Hyperfocus or flow? Attentional strengths in autism spectrum disorder. *Front Psychiatry*. 2022;13: 886692. doi:[10.3389/fpsy.2022.886692](https://doi.org/10.3389/fpsy.2022.886692)
- 501 18. Hupfeld KE, Abagis TR, Shah P. Living "in the zone": Hyperfocus in adult ADHD. *Atten Defic Hyperact Disord*. 2019;11: 191–208. doi:[10.1007/s12402-018-0272-y](https://doi.org/10.1007/s12402-018-0272-y)
- 502 19. Astle DE, Holmes J, Kievit R, Gathercole SE. Annual research review: The transdiagnostic revolution in neurodevelopmental disorders. *J Child Psychol Psychiatry*. 2022;63: 397–417. doi:[10.1111/jcpp.13481](https://doi.org/10.1111/jcpp.13481)

- 503 20. Zhang M, Huang Y, Jiao J, Yuan D, Hu X, Yang P, et al. Transdiagnostic symptom subtypes across autism spectrum disorders and attention deficit hyperactivity disorder: Validated by measures of neurocognition and structural connectivity. *BMC Psychiatry*. 2022;22: 102. doi:[10.1186/s12888-022-03734-4](https://doi.org/10.1186/s12888-022-03734-4)
- 504 21. Chadehumbe M. A comprehensive review of motor stereotypies in children. *Current Pediatrics Reports*. 2018;6: 30–33. doi:[10.1007/s40124-018-0153-z](https://doi.org/10.1007/s40124-018-0153-z)
- 505 22. Mackenzie K. Stereotypic movement disorders. *Seminars in Pediatric Neurology*. 2018;25: 19–24. doi:<https://doi.org/10.1016/j.spen.2017.12.004>
- 506 23. Tan A, Salgado M, Fahn S. The characterization and outcome of stereotypical movements in nonautistic children. *Mov Disord*. 1997;12: 47–52. doi:[10.1002/mds.870120109](https://doi.org/10.1002/mds.870120109)
- 507 24. Charlop MH. The effects of echolalia on acquisition and generalization of receptive labeling in autistic children. *J Appl Behav Anal*. 1983;16: 111–26. doi:[10.1901/jaba.1983.16-111](https://doi.org/10.1901/jaba.1983.16-111)
- 508 25. Pruccoli J, Spadoni C, Orsenigo A, Parmeggiani A. Should echolalia be considered a phonic stereotypy? A narrative review. *Brain Sci*. 2021;11. doi:[10.3390/brainsci11070862](https://doi.org/10.3390/brainsci11070862)
- 509 26. Prizant BM, Duchan JF. The functions of immediate echolalia in autistic children. *J Speech Hear Disord*. 1981;46: 241–9. doi:[10.1044/jshd.4603.241](https://doi.org/10.1044/jshd.4603.241)
- 510 27. Fertel-Daly D, Bedell G, Hinojosa J. Effects of a weighted vest on attention to task and self-stimulatory behaviors in preschoolers with pervasive developmental disorders. *The American Journal of Occupational Therapy*. 2001;55: 629–640. doi:[10.5014/ajot.55.6.629](https://doi.org/10.5014/ajot.55.6.629)
- 511 28. Milne E, Scope A, Pascalis O, Buckley D, Makeig S. Independent component analysis reveals atypical electroencephalographic activity during visual perception in individuals with autism. *Biol Psychiatry*. 2009;65: 22–30. doi:[10.1016/j.biopsych.2008.07.017](https://doi.org/10.1016/j.biopsych.2008.07.017)
- 512 29. Snijders TM, Milivojevic B, Kemner C. Atypical excitation-inhibition balance in autism captured by the gamma response to contextual modulation. *Neuroimage Clin*. 2013;3: 65–72. doi:[10.1016/j.nicl.2013.06.015](https://doi.org/10.1016/j.nicl.2013.06.015)
- 513 30. Murphy E, Benítez-Burraco A. Language deficits in schizophrenia and autism as related oscillatory connectopathies: An evolutionary account. *Neurosci Biobehav Rev*. 2017;83: 742–764. doi:[10.1016/j.neubiorev.2016.07.029](https://doi.org/10.1016/j.neubiorev.2016.07.029)

- 514 31. Wang J, Barstein J, Ethridge LE, Mosconi MW, Takarae Y, Sweeney JA. Resting state EEG abnormalities in autism spectrum disorders. *J Neurodev Disord.* 2013;5: 24. doi:[10.1186/1866-1955-5-24](https://doi.org/10.1186/1866-1955-5-24)
- 515 32. Berman JI, Liu S, Bloy L, Blaskey L, Roberts TPL, Edgar JC. Alpha-to-gamma phase-amplitude coupling methods and application to autism spectrum disorder. *Brain Connect.* 2015;5: 80–90. doi:[10.1089/brain.2014.0242](https://doi.org/10.1089/brain.2014.0242)
- 516 33. Kleiner M, Brainard DH, Pelli DG. What’s new in psychtoolbox-3? *Perception.* 2007;36S: 14.
- 517 34. Brainard DH. [The psychophysics toolbox](#). *Spat Vis.* 1997;10: 433–6.
- 518 35. Pelli DG. [The VideoToolbox software for visual psychophysics: Transforming numbers into movies](#). *Spat Vis.* 1997;10: 437–42.
- 519 36. Tottenham N, Tanaka JW, Leon AC, McCarry T, Nurse M, Hare TA, et al. The NimStim set of facial expressions: Judgments from untrained research participants. *Psychiatry Res.* 2009;168: 242–9. doi:[10.1016/j.psychres.2008.05.006](https://doi.org/10.1016/j.psychres.2008.05.006)
- 520 37. Delorme A, Makeig S. EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J Neurosci Methods.* 2004;134: 9–21. doi:[10.1016/j.jneumeth.2003.10.009](https://doi.org/10.1016/j.jneumeth.2003.10.009)
- 521 38. Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, et al. MEG and EEG data analysis with MNE-python. *Front Neurosci.* 2013;7: 267. doi:[10.3389/fnins.2013.00267](https://doi.org/10.3389/fnins.2013.00267)
- 522 39. Rouder JN, Speckman PL, Sun D, Morey RD, Iverson G. Bayesian t tests for accepting and rejecting the null hypothesis. *Psychon Bull Rev.* 2009;16: 225–37. doi:[10.3758/PBR.16.2.225](https://doi.org/10.3758/PBR.16.2.225)
- 523 40. Boyd BA, McDonough SG, Bodfish JW. Evidence-based behavioral interventions for repetitive behaviors in autism. *J Autism Dev Disord.* 2012;42: 1236–48. doi:[10.1007/s10803-011-1284-z](https://doi.org/10.1007/s10803-011-1284-z)
- 524 41. Koegel RL, Covert A. The relationship of self-stimulation to learning in autistic children. *J Appl Behav Anal.* 1972;5: 381–7. doi:[10.1901/jaba.1972.5-381](https://doi.org/10.1901/jaba.1972.5-381)
- 525 42. Varni JW, Lovaas OI, Koegel RL, Everett NL. An analysis of observational learning in autistic and normal children. *J Abnorm Child Psychol.* 1979;7: 31–43. doi:[10.1007/BF00924508](https://doi.org/10.1007/BF00924508)

- 526 43. Pierce K, Courchesne E. Evidence for a cerebellar role in reduced exploration and stereotyped behavior in autism. *Biol Psychiatry*. 2001;49: 655–64. doi:[10.1016/s0006-3223\(00\)01008-8](https://doi.org/10.1016/s0006-3223(00)01008-8)
- 527 44. Keenan M, Dillenburger K, Röttgers HR, Dounavi K, Jónsdóttir SL, Moderato P, et al. Autism and ABA: The gulf between north america and europe. *Review Journal of Autism and Developmental Disorders*. 2015;2: 167–183. doi:[10.1007/s40489-014-0045-2](https://doi.org/10.1007/s40489-014-0045-2)
- 528 45. Raulston TJ, Hansen SG, Machalicek W, McIntyre LL, Carnett A. Interventions for repetitive behavior in young children with autism: A survey of behavioral practices. *J Autism Dev Disord*. 2019;49: 3047–3059. doi:[10.1007/s10803-019-04023-y](https://doi.org/10.1007/s10803-019-04023-y)
- 529 46. Canela C, Buadze A, Dube A, Eich D, Liebreinz M. Skills and compensation strategies in adult ADHD - a qualitative study. *PLoS One*. 2017;12: e0184964. doi:[10.1371/journal.pone.0184964](https://doi.org/10.1371/journal.pone.0184964)
- 530 47. Rong Y, Yang C-J, Jin Y, Wang Y. Prevalence of attention-deficit/hyperactivity disorder in individuals with autism spectrum disorder: A meta-analysis. *Research in Autism Spectrum Disorders*. 2021;83: 101759. doi:<https://doi.org/10.1016/j.rasd.2021.101759>
- 531 48. Tse CYA, Pang CL, Lee PH. Choosing an appropriate physical exercise to reduce stereotypic behavior in children with autism spectrum disorders: A non-randomized crossover study. *J Autism Dev Disord*. 2018;48: 1666–1672. doi:[10.1007/s10803-017-3419-3](https://doi.org/10.1007/s10803-017-3419-3)
- 532 49. Rex C, Charlop MH, Spector V. Using video modeling as an anti-bullying intervention for children with autism spectrum disorder. *J Autism Dev Disord*. 2018;48: 2701–2713. doi:[10.1007/s10803-018-3527-8](https://doi.org/10.1007/s10803-018-3527-8)
- 533 50. Campbell JM, Ferguson JE, Herzinger CV, Jackson JN, Marino CA. Combined descriptive and explanatory information improves peers’ perceptions of autism. *Res Dev Disabil*. 2004;25: 321–39. doi:[10.1016/j.ridd.2004.01.005](https://doi.org/10.1016/j.ridd.2004.01.005)
- 534 51. Chen P-Y, Schwartz IS. Bullying and victimization experiences of students with autism spectrum disorders in elementary schools. *Focus on Autism and Other Developmental Disabilities*. 2012;27: 200–212. doi:[10.1177/1088357612459556](https://doi.org/10.1177/1088357612459556)