

Today: *

- * boundary of chains
- * cycles, boundaries, homology groups

Recall boundary of p -simplex σ (over \mathbb{Z}_2 or \mathbb{Z}).

Note that \mathcal{Z}_p^σ is a $(p-1)$ -chain (in both cases).

The boundary of a p -chain is the sum of the boundaries of its p -simplices.

$$\bar{c} = \sum_{i=1}^m a_i \sigma_i \Rightarrow \partial_p \bar{c} = \sum_{i=1}^m a_i (\partial_p \sigma_i), \text{ which is also a } (p-1)\text{-chain.}$$


Examples

Note that vertices shared by two edges in the chain do not appear in its boundary.


$$\partial_2 \left(\begin{array}{c} \bar{d}_1 \\ \begin{array}{ccccc} v_2 & & v_3 & & v_4 \\ & \swarrow e_2 & \searrow e_6 & & \\ v_0 & & v_1 & & \end{array} \end{array} \right) = \begin{array}{c} v_2 \quad \quad \quad v_3 \quad \quad \quad v_4 \\ \quad \swarrow e_2 \quad \searrow e_6 \quad \quad \\ v_0 \quad \quad \quad v_1 \end{array}$$

The boundary of a triangle is made of its edges.

Consider $\partial_2 \bar{d}_1$ over \mathbb{Z} now:



Notice that the induced orientation on \bar{e}_1 is opposite to its own orientation.



The dimension is often omitted, and we just talk about $\partial \bar{c}$ of a p -chain \bar{c} , with the dimension understood.

Taking the boundary maps a p -chain to a $(p-1)$ -chain. Equivalently, we can talk about the map $\partial_p: C_p \rightarrow C_{p-1}$. Notice that such a map is defined for each p in the range $1 \leq p \leq \dim K$.

Also, $\partial_p(\bar{c} \pm \bar{c}') = \partial_p \bar{c} \pm \partial_p \bar{c}'$ for 2 p -chains \bar{c}, \bar{c}' . Hence ∂_p is a homomorphism, referred to as the **p -th boundary map** or homomorphism. *commutes with \pm , which is the definition of a homomorphism*

Similarly, we have ∂_{p-1} , which is the $(p-1)$ -st boundary homomorphism, and ∂_{p-2} , and so on. $\partial_{p-1}: C_{p-1} \rightarrow C_{p-2}$.

∂_p is a homomorphism over \mathbb{Z} (or \mathbb{Q}, \mathbb{R}) as well, not just over \mathbb{Z}_2 .


A **chain complex** is a sequence of chain groups connected by boundary homomorphisms

$$\dots \xrightarrow{\partial_{p+2}} C_{p+1} \xrightarrow{\partial_{p+1}} C_p \xrightarrow{\partial_p} C_{p-1} \rightarrow \dots$$

The word complex as used here is different from a simplicial complex. At the same time, the chain complex is indeed an abstract simplicial complex, with elements connected by boundary homomorphisms.

Cycles A **p-cycle** is a p-chain with empty boundary.

For instance, consider the 1-chain \bar{z}_1 of K :



Hence \bar{z}_1 is a 1-cycle.

Alternatively, a p-chain \bar{c} is a p-cycle if $\partial \bar{c} = 0$.



Since ∂ commutes with $+$, the p -cycles of K form a group, denoted by Z_p or $Z_p(K)$. Z_p is a subgroup of C_p . Also, $Z_p = \ker \partial_p$, i.e., Z_p is the kernel of the p th boundary homomorphism. Notice that $\partial_p \bar{z} = 0$ for $\bar{z} \in Z_p$. Also, just as C_p is, Z_p is an Abelian group.

The surface of a tetrahedron (made of four triangles) is a 2-cycle. Notice that its boundary is empty.


So, all cycles are also chains, but not the other way usually. But for $p=0$, $\partial_0 v_j = 0$, i.e., the boundary of a vertex is empty (by definition). Hence every 0-chain is also a 0-cycle, i.e., $Z_0 = C_0$.

But typically, $Z_p \subset C_p$ for $p \geq 1$.


Boundaries A p -boundary \bar{b} is a p -chain that is the boundary of some $(p+1)$ -chain \bar{d} , i.e., $\bar{b} = \partial_{p+1} \bar{d}$ for $\bar{d} \in C_{p+1}$.

Again, since ∂ commutes with $+$ (or $+$) we get a group of p -boundaries B_p (or $B_p(K)$). B_p is a subgroup of Z_p , and of C_p .

B_p is the image of $\partial_{p+1}: C_{p+1} \rightarrow C_p$. $B_p = \text{im } \partial_{p+1}$. B_p is abelian.



$\bar{z}_1 = \partial \bar{f}_0$, and hence is a 1-boundary.
→ 2-chain made of triangle f_0





But \bar{z}_2 is not a 1-boundary.

We have $B_p \subset Z_p \subset C_p$ (in general).



So, in summary, p -cycle \bar{z} : $\partial \bar{z} = 0$; $Z_p = \ker \partial_p$.
 p -boundary \bar{b} : $\bar{b} = \partial_{p+1} \bar{d}$; $B_p = \text{im } \partial_{p+1}$.

Examples

v_2 v_4 is a 0-boundary,
 as it is $\partial_1 \bar{c}_1$.



Now consider oriented K_1 (over \mathbb{Z}):



$$\bar{b} = \begin{bmatrix} 0 \\ -1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \end{bmatrix}$$

\bar{b}_1 is a 1-boundary, as $\bar{b}_1 = \partial_2 \bar{f}_1 \rightarrow$ elementary 2-chain of f_1

let's try to enumerate how many p -cycles and p -boundaries are there for general p . We want to study cycles that are not boundaries.

$p=0$ case (over \mathbb{Z}_2)

\bar{c} , a 1-chain, is a collection of edges. $\partial_1 \bar{c}$ gives end points of edges with duplicate end points canceled in pairs, leaving an even number of distinct v_j 's.

(156)

If K is connected, for every set of even number of vertices, we can find paths made of edges that connect the vertices such that the 1-chain made of these edges has as its boundary the collection of vertices.


e.g., $\partial_1(e_0 + e_2) = \{v_0, v_1, v_2, v_3\}$.




Thus, every set of even number of vertices (v_j 's) is a 0-boundary, while every odd set is not. Hence, if K is connected, exactly half of the 0-cycles are 0-boundaries.

But we typically cannot make similar statements about counts of p -cycles and p -boundaries for $p \geq 1$.

We want to characterize cycles that are not boundaries, as they capture holes, e.g., as \bar{z}_2 here.



An observation: Consider $\bar{b}_1 = \partial_2 \bar{f}_1$.



$\partial \bar{b}_1 = 0$ (as each v_j cancels in pairs).

In other words, $\partial_1 \partial_2 f_1 = 0$. This result holds in general!

Fundamental Lemma of Homology $\partial_p \partial_{p+1} \bar{d} = 0 \quad \forall p \in \mathbb{Z}$.

In words, boundary of a boundary is empty.

Proof (over \mathbb{Z}_2) For each $(p+1)$ -simplex τ , $\partial_p \partial_{p+1} \tau = 0$, as $\partial_{p+1} \tau$ consists of all p -faces of τ . Each $(p-1)$ -face of τ belongs to exactly two p -faces.

$$\partial_2 \left(\partial_3 \left(\begin{array}{c} v_3 \\ \diagup \quad \diagdown \\ v_2 \quad v_1 \\ \diagdown \quad \diagup \\ v_0 \end{array} \right) \right) = \partial_2 \left(\text{union of 4 triangles} \right) = 0,$$

tetrahedron

as each edge is shared by exactly two triangles.


$$\partial_p \left(\partial_{p+1} \tau = \partial_{p+1} [v_0 \dots v_{p+1}] \right) = \partial_p \left(\sum_{j=0}^{p+1} [v_0 \dots \hat{v}_j \dots v_{p+1}] \right)$$

WLOG, the $(p-1)$ -simplex $[v_0 \dots v_{p-1}]$ is a face of $[v_0 \dots v_{p-1}, v_p]$ and $[v_0 \dots v_{p-1}, v_{p+1}]$.

The result holds over \mathbb{Z} as well (over any ring, in fact). We must consider induced orientations when taking ∂_{p+1} and ∂_p .

A p boundary is also a p -cycle. Hence B_p is a subgroup of Z_p .

The groups C_p, Z_p, B_p for various p are related as follows:



Homology Groups

158

Since B_p is a subgroup of Z_p , we can take quotients.



Def The p -th homology group is the p -th cycle group modulo the p -boundary group, $H_p = Z_p / B_p$.

H_p has the classes of cycles that are not boundaries.


Each element of $H_p(K)$ is obtained by adding p -boundaries to given p -cycles (in their class), i.e., $\bar{z} + B_p$, where $\bar{z} \in Z_p$.

$\bar{z} + B_p$ is a coset of B_p in Z_p .

Example



\bar{z}_2 and \bar{z}_4 are cycles going around the same hole.



\bar{z}_5 also goes around the same hole as \bar{z}_2 (and \bar{z}_4).

We could use any one cycle going around the hole ($\bar{z}_2, \bar{z}_4, \bar{z}_5$) as a representative of $H_1(K)$.




In general, another cycle $\bar{z}' = \bar{z} + \bar{b}$ for \bar{z} in $H_p(K)$ and $\bar{b} \in B_p$ is in the same class as \bar{z} , i.e., $\bar{z}' + B_p = \bar{z} + B_p$ (as $\bar{b} + B_p = B_p$ itself). This is a class in $H_p(K)$, and any two cycles in this class are said to be **homologous**, written as $\bar{z} \sim \bar{z}'$.

In this setting, addition of classes is well-defined:

$$(\bar{z} + B_p) + (\bar{z}' + B_p) = (\bar{z} + \bar{z}' + B_p), \text{ independent of the particular representatives } \bar{z} \text{ and } \bar{z}'.$$



Thus H_p is indeed a group, and is abelian.

Another example




Notice $\bar{z}_6 \not\sim \bar{z}_2$, as we cannot get $\bar{z}_6 = \bar{z}_2 + \bar{b}$ for any $\bar{b} \in B_1$.


Now consider two more cycles \bar{z}_7 and \bar{z}_8 as shown.



$\bar{z}_7 \sim \bar{z}_8$ as



Now, consider \bar{z}_9 and \bar{z}_6 . Notice that $\bar{z}_9 \not\sim \bar{z}_6$. But



$\bar{z}_{10} = \bar{z}_9 + \bar{z}_6$. $\bar{z}_{10} \not\sim \bar{z}_9$, but $\bar{z}_{10} \sim \bar{z}_8$

homology class of \bar{z}_2

To describe $H_1(K_2)$ completely, we could present $[\bar{z}_2]$ and $[\bar{z}_6]$, or equivalently, $[\bar{z}_2]$ and $[\bar{z}_7]$, or $[\bar{z}_6]$ and $[\bar{z}_7]$.

Intuitively, since K_2 has two holes, we expect $H_1(K_2)$ to have two classes.