Optimal Cycles and LP/IP

Bala Krishnamoorthy
Washington State U.

Optimal ~~Cycles~~ and LP/IP Chains

Bala Krishnamoorthy
Washington State U.

OHCP

The Optimal Homologous Cycle Problem:
Given p-cycle c in simplicial complex K, find homologous cycle x that is "optimal."

OHCP

The Optimal Homologous Cycle Problem:

Given p-cycle c in simplicial complex K, find homologous cycle x that is "optimal."

$x \sim c$

$x - c = \partial_{p+1} s$

$\sum w_i |x_i|$ is min.

OHCP

The Optimal Homologous Cycle Problem:
Given p-cycle c in simplicial complex K,
find homologous cycle x that is "optimal."

$x \sim c$

$x - c = \partial_{p+1} s$

$\sum w_i |x_i|$ is min.





OHCP

The Optimal Homologous Chain Problem:

Given p-chain c in simplicial complex K, find "homologous" chain x that is "optimal."

$x \sim c$

$x - c = \partial_{p+1} s$

$\sum w_i |x_i|$ is min.





OHCP

The Optimal Homologous Chain Problem:

Given p-chain c in simplicial complex K, find "homologous" chain x that is "optimal."

$x \sim c$

$\sum w_i |x_i|$ is min.

$x - c = \partial_{p+1} s$

$$\min \quad w^T x$$

$$\text{s.t.} \quad x = c + [\partial_{p+1}] s$$

$$x \in \mathbb{Z}^m, \; s \in \mathbb{Z}^n$$

OBCP

The Optimal Bounding Chain Problem:

Given p-boundary c in simplicial complex K, find $(p+1)$-chain s bounding c that is "optimal."

$c = \partial_{p+1} s$

$\sum v_j |s_j|$ is min.

OBCP

The Optimal Bounding Chain Problem:
Given p-boundary c in simplicial complex K, find $(p+1)$-chain s bounding c that is "optimal."

$c = \partial_{p+1} s$

$\sum v_j |s_j|$ is min.

$$\min \; v^T s$$
$$\text{s.t.} \quad c = [\partial_{p+1}] s$$
$$s \in \mathbb{Z}^n$$

MSFN

Ibrahim, K, Vixie (2013)

Multiscale Simplicial Flat Norm:

Given p-chain c in simplicial complex K, find "homologous" chain x that is "optimal".

$x \sim c$

$x = c - \partial_{p+1} s$

$\sum w_i |x_i| + \lambda \sum v_j s_j$

$\lambda \geq 0$

MSFN

Ibrahim, K, Vixie (2013)

Multiscale Simplicial Flat Norm:

Given p-chain c in simplicial complex K, find "homologous" chain x that is "optimal."

$x \sim c$

$\sum w_i |x_i| + \lambda \sum v_j s_j$

$x = c - \partial_{p+1} s$

$\lambda \geq 0$

$$\begin{aligned} \min \quad & w^T x + \lambda v^T s \\ \text{s.t.} \quad & x = c - [\partial_{p+1}] s \\ & x \in \mathbb{Z}^m, \; s \in \mathbb{Z}^n \end{aligned}$$

MSFN Example



$X = C - \partial S$

MSFN Example



$X = C - \partial S$



$$w_i = \begin{cases} 1, & i = 1,3,4,6,7 \\ \sqrt{2}, & i = 2,5 \end{cases} \Bigg\} \text{ lengths}$$

$$v_j = \frac{1}{2},\ j = 1,2,3 \ \Big\} \text{ areas}$$

MSFN Example

boundary

$\partial\Big($  $\Big) =$ 



$- \partial\Big($  $\Big) =$ 

for $\lambda = 2.5$,

$$F_\lambda = 3 + \frac{\lambda}{2}$$

$$= 4.25$$

MSFN Example

boundary





for $\lambda = 2.5$,

$$F_\lambda = 3 + \frac{\lambda}{2} = 4.25$$

for $\lambda = 1$, $F_\lambda = 1 + \frac{3\lambda}{2} = 2.5$



MSFN: Applications

* protrusions in chemistry

E Alvarado, Z Liu, M Servis, BK, A Clark

JCTC 2020



MSFN: Applications

* protrusions in chemistry

E Alvarado, Z Liu, M Servi's, BK, A Clark

JCTC 2020



* validating synthetic power networks

K Lyman, R Meyur, BK, M Halappanavar

IJCGA 2025



OHCP under Persistence

* volume-optimal cycles: Obayashi, 2018
* minimal persistent cycles: Dey, Hou, Mandal, 2020 (over $\mathbb{Z}_2$)

OHCP under Persistence

* volume-optimal cycles: Obayashi, 2018
* minimal persistent cycles: Dey, Hou, Mandal, 2020 (over $\mathbb{Z}_2$)



↳ Example from Li et al., 2021

OHCP under Persistence

* volume-optimal cycles: Obayashi, 2018
* minimal persistent cycles: Dey, Hou, Mandal, 2020 (over $\mathbb{Z}_2$)



↳ Example from Li et al., 2021

ℓ_0-min per. cycle

vol-optimal cycle

OHCP under Persistence

* volume-optimal cycles: Obayashi, 2018
* minimal persistent cycles: Dey, Hou, Mandal, 2020 (over $\mathbb{Z}_2$)



↳ Example from Li et al., 2021

ℓ_0-min per. cycle

vol-optimal cycle

! Today: OHCP in a single simplicial complex with homology over $\mathbb{Z}$

Cheaper by the Dozen!



T A filling of T A filling of $2T$

(R. Young, 2013)

previous examples by
- L. Young, 1963
- F. Morgan, 1984
- B. White, 1984

Cheaper by the Dozen!



T A filling of T A filling of $2T$

(R. Young, 2013)

previous examples by
- L. Young, 1963
- F. Morgan, 1984
- B. White, 1984

min. surface for $2T$ < 2 (min surface for T) !

CHEAPER BY THE DOZEN!



T A filling of T A filling of $2T$

(R. Young, 2013)

previous examples by
- L. Young, 1963
- F. Morgan, 1984
- B. White, 1984

min. surface for $2T < 2$ (min surface for T) !

Q. If S is the min surface for curve T, when is $2S$ the min surface for $2T$?

OHCP as an Integer Program

$$\min_{x,y} \sum_i w_i |x_i| \quad \text{such that} \qquad \text{(OHCP)}$$

$$x = c + [\partial_{p+1}]s, \quad x \in \mathbb{Z}^m, \quad s \in \mathbb{Z}^n \quad (w_i \geq 0)$$

OHCP as an Integer Program

piecewise linear

$$\min_{x,y} \sum_i w_i |x_i| \quad \text{such that} \qquad \text{(OHCP)}$$

$$x = c + [\partial_{p+1}]s, \quad x \in \mathbb{Z}^m, \quad s \in \mathbb{Z}^n \quad (w_i \geq 0)$$

$$\min \sum_i w_i (x_i^+ + x_i^-)$$

$$\text{s.t.} \quad x^+ - x^- = c + [\partial_{p+1}](s^+ - s^-) \qquad \text{(IP)}$$

$$x^+, x^-, s^+, s^- \geq 0 \quad x^+, x^- \in \mathbb{Z}^m, \; s^+, s^- \in \mathbb{Z}^n$$

OHCP as an Integer Program

piecewise linear

$$\min_{x,y} \sum_i w_i |x_i| \quad \text{such that} \qquad \text{(OHCP)}$$

$$x = c + [\partial_{p+1}]s, \quad x \in \mathbb{Z}^m, \quad s \in \mathbb{Z}^n \quad (w_i \geq 0)$$

$$\min \sum_i w_i (x_i^+ + x_i^-)$$

$$\text{s.t.} \quad x^+ - x^- = c + [\partial_{p+1}](s^+ - s^-) \qquad \text{(IP)}$$

$$x^+, x^-, s^+, s^- \geq 0 \qquad \boxed{x^+, x^- \in \mathbb{Z}^m, \; s^+, s^- \in \mathbb{Z}^n}$$

ignore to get LP

LINEAR PROGRAMMING (LP)

Optimization in Calculus I

$$\min_{a \le x \le b} f(x)$$

$$f'(x) = 0, \quad f''(x) > 0$$



LINEAR PROGRAMMING (LP)

Optimization in Calculus I

$$\min_{a \le x \le b} f(x)$$

$f'(x) = 0$, $f''(x) > 0$

also check end points!

Here, minimum is at $x = a$



LINEAR PROGRAMMING (LP)

Optimization in Calculus I

$$\min_{a \le x \le b} f(x)$$

$f'(x) = 0$, $f''(x) > 0$

also check end points!

Here, minimum is at $x = a$



$$\left\{ \begin{array}{l} \min g(x) \\ a \le x \le b \end{array} \right\}$$ where $g(x)$ is linear?

Need to check ONLY the end points!

LINEAR PROGRAMMING (LP)

Extend linear case to high dimensions $\Rightarrow$ LP
(also called linear optimization)

$$\begin{aligned} \max \quad & x_1 + x_2 \\ \text{s.t.} \quad & 5x_1 - x_2 \leq 7 \\ & 3x_1 - 4x_2 \geq -3 \\ & x_1, x_2 \geq 0 \end{aligned}$$



Need to check ONLY corner points ($\equiv$ end pts.)

LINEAR PROGRAMMING (LP)

Extend linear case to high dimensions $\Rightarrow$ LP
(also called linear optimization)

$$\begin{aligned} \max \quad & x_1 + x_2 \\ \text{s.t.} \quad & 5x_1 - x_2 \leq 7 \\ & 3x_1 - 4x_2 \geq -3 \\ & x_1, x_2 \geq 0 \end{aligned}$$



Need to check ONLY corner points ($\equiv$ end pts.)

Require $x_j \in \mathbb{Z} \Rightarrow$ integer programming (IP)

IP and Total Unimodularity

$$\min \{ c^T x \mid Ax = b, x \geq 0, x \in \mathbb{Z}^n \} \quad \text{(IP)}$$

$$\min \{ c^T x \mid Ax = b, x \geq 0 \} \quad \text{(LP)}$$

IP and Total Unimodularity

$$\min \{ c^T x \mid Ax = b, x \geq 0, x \in \mathbb{Z}^n \} \quad \text{(IP)}$$

$$\min \{ c^T x \mid Ax = b, x \geq 0 \} \quad \text{(LP)}$$

<u>Result</u>: (IP) can always be solved in polynomial time by solving (LP) iff A is totally unimodular (TU).

IP and Total Unimodularity

$$\min \{ c^T x \mid Ax = b, x \geq 0, x \in \mathbb{Z}^n \} \quad \text{(IP)}$$

$$\min \{ c^T x \mid Ax = b, x \geq 0 \} \quad \text{(LP)}$$

<u>Result</u>: (IP) can always be solved in polynomial time by solving (LP) iff A is totally unimodular (TU).

A is TU if every square submatrix has determinant $-1, 0,$ or 1.

In particular, $A_{ij} \in \{-1, 0, 1\} \ \forall i, j$.

IP and Total Unimodularity

$$\min \{ c^T x \mid Ax = b, x \geq 0, x \in \mathbb{Z}^n \} \quad \text{(IP)}$$

$$\min \{ c^T x \mid Ax = b, x \geq 0 \} \quad \text{(LP)}$$

<u>Result</u>: (IP) can always be solved in polynomial time by solving (LP) iff A is totally unimodular (TU).

A is TU if every square submatrix has determinant $-1, 0,$ or 1.

In particular, $A_{ij} \in \{-1, 0, 1\} \ \forall i, j$.

e.g., node-arc incidence matrix of directed graph

OHCP AND TU OF $[\partial_{p+1}]$

total unimodularity

$$\min \sum_i w_i (x_i^+ + x_i^-)$$

$$\text{s.t.} \quad x^+ - x^- = c + [\partial_{p+1}](s^+ - s^-) \qquad \text{(LP)}$$

$$x^+, x^-, s^+, s^- \geq 0$$

OHCP AND TU OF $[\partial_{p+1}]$

total unimodularity

$$\min \sum_i w_i (x_i^+ + x_i^-)$$

$$\text{s.t.} \quad x^+ - x^- = c + [\partial_{p+1}](s^+ - s^-) \qquad \text{(LP)}$$

$$x^+, x^-, s^+, s^- \geq 0$$

The constraint matrix of above LP is TU iff $[\partial_{p+1}]$ is TU.

OHCP AND TU OF $[\partial_{p+1}]$

total unimodularity

$$\min \sum_i w_i (x_i^+ + x_i^-)$$

$$\text{s.t.} \quad x^+ - x^- = c + [\partial_{p+1}](s^+ - s^-) \qquad \text{(LP)}$$

$$x^+, x^-, s^+, s^- \geq 0$$

The constraint matrix of above LP is TU iff $[\partial_{p+1}]$ is TU.

$\Rightarrow$ OHCP is solvable in poly time when $[\partial_{p+1}]$ is TU.

OHCP AND TU OF $[\partial_{p+1}]$

total unimodularity

$$\min \sum_i w_i (x_i^+ + x_i^-)$$

$$\text{s.t.} \quad x^+ - x^- = c + [\partial_{p+1}](s^+ - s^-) \qquad \text{(LP)}$$

$$x^+, x^-, s^+, s^- \geq 0$$

The constraint matrix of above LP is TU iff $[\partial_{p+1}]$ is TU.

$\Rightarrow$ <u>OHCP</u> is solvable in poly time when $[\partial_{p+1}]$ is TU.

OBCP, MSFN as well

When is $[\partial_{p+1}]$ TU?

Dey, Hirani, K, 2010: $[\partial_{p+1}]$ is TU iff K has "no relative torsion."
(DHK'10)

When is $[\partial_{p+1}]$ TU?

Dey, Hirani, K, 2010: (DHK'10) $[\partial_{p+1}]$ is TU iff K has "no relative torsion."

Poincaré: TU in 1899!

When is $[\partial_{p+1}]$ TU?

Dey, Hirani, K, 2010: (DHK'10) $[\partial_{p+1}]$ is TU iff K has "no relative torsion."

Special cases

✓ K is an orientable manifold

Poincaré: TU in 1899!

When is $[\partial_{p+1}]$ TU?

Dey, Hirani, K, 2010: (DHK'10) $[\partial_{p+1}]$ is TU iff K has "no relative torsion."

Poincaré: TU in 1899!

Special cases

✓ K is an orientable manifold

✓ $\dim(K) = d$, K in $\mathbb{R}^d$

e.g., tetrahedra in $\mathbb{R}^3$

When is $[\partial_{p+1}]$ TU?

Dey, Hirani, K, 2010: (DHK '10) $[\partial_{p+1}]$ is TU iff K has "no relative torsion."

Poincaré: TU in 1899!

Special cases

✓ K is an orientable manifold

✓ $\dim(K) = d$, K in $\mathbb{R}^d$

e.g., tetrahedra in $\mathbb{R}^3$



✓ $[\partial_2]$ is TU $\Leftrightarrow$ K has "no Möbius strip"

(DHK '11)

Dual Graph: Max Flow

* Chen & Freedman, 2008 (OHCP, $\mathbb{Z}_2$)
* Dey, Hou, Mandal, 2020 (min. pers. cycle, $\mathbb{Z}_2$)

Dual Graph: Max Flow

* Chen & Freedman, 2008 (OHCP, $\mathbb{Z}_2$)
* Dey, Hou, Mandal, 2020 (min. pers. cycle, $\mathbb{Z}_2$)
* Borradaile, Maxwell, Nayyeri, 2020 (OBCP, $\mathbb{Z}_2$)

Dual Graph: Max Flow

* Chen & Freedman, 2008 (OHCP, $\mathbb{Z}_2$)
* Dey, Hou, Mandal, 2020 (min. pers. cycle, $\mathbb{Z}_2$)
* Borradaile, Maxwell, Nayyeri, 2020 (OBCP, $\mathbb{Z}_2$)
* Chambers, Erickson, Nayyeri, 2009, 2012 (OHCP, $\mathbb{R}$)

Dual Graph: Max Flow

* Chen & Freedman, 2008 (OHCP, $\mathbb{Z}_2$)
* Dey, Hou, Mandal, 2020 (min. pers. cycle, $\mathbb{Z}_2$)
* Borradaile, Maxwell, Nayyeri, 2020 (OBCP, $\mathbb{Z}_2$)
* Chambers, Erickson, Nayyeri, 2009, 2012 (OHCP, $\mathbb{R}$)
* Sullivan, 1990 (OBCP, $\mathbb{Z}$)

Dual Graph: Max Flow

* Chen & Freedman, 2008 (OHCP, $\mathbb{Z}_2$)
* Dey, Hou, Mandal, 2020 (min. pers. cycle, $\mathbb{Z}_2$)
* Borradaile, Maxwell, Nayyeri, 2020 (OBCP, $\mathbb{Z}_2$)
* Chambers, Erickson, Nayyeri, 2009, 2012 (OHCP, $\mathbb{R}$)
* Sullivan, 1990 (OBCP, $\mathbb{Z}$)

! dual graph: tricky when c is not a cycle, MSFN

Dual Graph: Max Flow

* Chen & Freedman, 2008 (OHCP, $\mathbb{Z}_2$)
* Dey, Hou, Mandal, 2020 (min. pers. cycle, $\mathbb{Z}_2$)
* Borradaile, Maxwell, Nayyeri, 2020 (OBCP, $\mathbb{Z}_2$)
* Chambers, Erickson, Nayyeri, 2009, 2012 (OHCP, $\mathbb{R}$)
* Sullivan, 1990 (OBCP, $\mathbb{Z}$)

! dual graph: tricky when c is not a cycle, MSFN

Q: What happens when boundary matrix is not TU?

MSFN in Codim. 2





C

X

K: 2-skeleton of solid cube

2-skeleton of solid torus

MSFN in Codim. 2





C

X

K: 2-skeleton of solid cube 2-skeleton of solid torus

$[\partial_2]$ is not TU, but MSFN LP gives integer soln!

MSFN in Codim. 2





c

x

K: 2-skeleton of solid cube 2-skeleton of solid torus

$[\partial_2]$ is not TU, but MSFN LP gives integer soln!

Q1: When/Why does MSFN LP in codim. 2 give integer solutions for free?

MSFN in Codim. 2





C

X

K: 2-skeleton of solid cube 2-skeleton of solid torus

$[\partial_2]$ is not TU, but MSFN LP gives integer soln!

Q1: When/Why does MSFN LP in codim. 2 give integer solutions for free?

— Li et al, 2021: mostly $\{-1,0,1\}$ sol.'s Why?

Non-TU Neutralized K

– K, G Smith, 2018

✓ Define a condition when simplicial complex K is non total-unimodularity neutralized (NTUN)

Non-TU Neutralized K

– K, G Smith, 2018

✓ Define a condition when simplicial complex K is non total-unimodularity neutralized (NTUN)

– boundary matrix is not TU

Non-TU Neutralized K

– K, G Smith, 2018

✓ Define a condition when simplicial complex K is non total-unimodularity neutralized (NTUN)

- boundary matrix is not TU, but
- every OHCP LP, i.e., for every input chain, has an integer optimal solution

Non-TU Neutralized K

– K, G Smith, 2018

✓ Define a condition when simplicial complex K is non total-unimodularity neutralized (NTUN)

– boundary matrix is not TU, but

– every OHCP LP, i.e., for every input chain, has an integer optimal solution

✓ property of K (independent of weights on simplices and input chain)

Non-TU Neutralized K

<u>Theorem</u> Every OHCP LP on K has an integer optimal solution if every elementary chain in each relative torsion in K has a "neutralizing chain" in K.

Non-TU Neutralized K

<u>Theorem</u> Every OHCP LP on K has an integer optimal solution if every elementary chain in each relative torsion in K has a "<u>neutralizing chain</u>" in K.

ensures that every fractional basic solution is a convex combination of two integral basic solutions.

Non-TU Neutralized K

<u>Theorem</u> Every OHCP LP on K has an integer optimal solution if every elementary chain in each relative torsion in K has a "<u>neutralizing chain</u>" in K.

ensures that every fractional basic solution is a convex combination of two integral basic solutions.

K is non-TU neutralized (NTUN)

TU or Not ?





TU or NOT ?



X



TU or Not ?



✗



TU ✓

Non-TU Neutralized





Neutralizing Chain



a disc whose boundary consists of an odd number of red (heavy/manifold) edges

Non-TU Neutralized K

✓ Property of K – holds for every choice of weights on simplices and input chain

Non-TU Neutralized K

✓ Property of K – holds for every choice of weights on simplices and input chain

* OHCP LP polytope is **not** integral, but

! an integral optimal solution exists for every right hand-side vector and every relevant objective function vector!

Non-TU Neutralized K

✓ Property of K – holds for every choice of weights on simplices and input chain

* OHCP LP polytope is **not** integral, but

! an integral optimal solution exists for every right hand-side vector and every <u>relevant</u> objective function vector!

↳ s^+, s^- do not appear in objective function

Non-TU Neutralized K

✓ Property of K – holds for every choice of weights on simplices and input chain

* OHCP LP polytope is **not** integral, but

! an integral optimal solution exists for every right hand-side vector and every <u>relevant</u> objective function vector!

↳ s^+, s^- do not appear in objective function

Q2: NTUN result for MSFN ?

Median Shape

Hu, Hudelson, K, Tumurbaatar, Vixie, 2019

* flat distance between T, T': $\mathbb{F}_\lambda(T, T') = \mathbb{F}_\lambda(T - T')$

flat norm w/ scale λ

Median Shape

Hu, Hudelson, K, Tumurbaatar, Vixie, 2019

* flat distance between T, T': $\mathbb{F}_\lambda(T, T') = \mathbb{F}_\lambda(T - T')$

  flat norm w/ scale λ

* median of $\{T_i\}$:

$$\hat{T} = \arg\min_T \sum_i \mathbb{F}_\lambda(T - T_i)$$

Median Shape : LP

$$\min \begin{bmatrix} & [\ \mathbf{w} \quad \mathbf{w} \quad \lambda\mathbf{v}\ \lambda\mathbf{v}] & [\ \mathbf{w} \quad \mathbf{w} \quad \lambda\mathbf{v} \quad \lambda\mathbf{v}] & \cdots & [\ \mathbf{w} \quad \mathbf{w} \quad \lambda\mathbf{v} \quad \lambda\mathbf{v}] \end{bmatrix} \mathbf{x}$$

s.t.

$$\begin{bmatrix} \begin{bmatrix} I & -I \end{bmatrix} & \begin{bmatrix} -I & I & -B & B \end{bmatrix} & & & \\ \begin{bmatrix} I & -I \end{bmatrix} & & \begin{bmatrix} -I & I & -B & B \end{bmatrix} & & \\ \vdots & & & \ddots & \\ \begin{bmatrix} I & -I \end{bmatrix} & & & & \begin{bmatrix} -I & I & -B & B \end{bmatrix} \end{bmatrix} \mathbf{x} = \begin{bmatrix} \mathbf{t}_1 \\ \mathbf{t}_2 \\ \vdots \\ \mathbf{t}_N \end{bmatrix}$$

$$\begin{bmatrix} \mathbf{t}^+ & \mathbf{t}^- & \mathbf{r}_1^+ & \mathbf{r}_1^- & \mathbf{s}_1^+ & \mathbf{s}_1^- & \mathbf{r}_2^+ & \mathbf{r}_2^- & \mathbf{s}_2^+ & \mathbf{s}_2^- & \cdots & \mathbf{r}_N^+ & \mathbf{r}_N^- & \mathbf{s}_N^+ & \mathbf{s}_N^- \end{bmatrix} \geq \mathbf{0}.$$

$B = [\partial_{p+n}]$

Median Shape : LP

$$\min \begin{bmatrix} & [\,\mathbf{w} \quad \mathbf{w} \quad \lambda\mathbf{v} \; \lambda\mathbf{v}] & [\,\mathbf{w} \quad \mathbf{w} \quad \lambda\mathbf{v} \quad \lambda\mathbf{v}] & \cdots & [\,\mathbf{w} \quad \mathbf{w} \quad \lambda\mathbf{v} \quad \lambda\mathbf{v}] \end{bmatrix} \mathbf{x}$$

s.t.

$$\begin{bmatrix} \begin{bmatrix} I & -I \end{bmatrix} & \begin{bmatrix} -I & I & -B & B \end{bmatrix} & & & \\ \begin{bmatrix} I & -I \end{bmatrix} & & \begin{bmatrix} -I & I & -B & B \end{bmatrix} & & \\ \vdots & & & \ddots & \\ \begin{bmatrix} I & -I \end{bmatrix} & & & & \begin{bmatrix} -I & I & -B & B \end{bmatrix} \end{bmatrix} \mathbf{x} = \begin{bmatrix} \mathbf{t}_1 \\ \mathbf{t}_2 \\ \vdots \\ \mathbf{t}_N \end{bmatrix}$$

$$\begin{bmatrix} \mathbf{t}^+ & \mathbf{t}^- & \mathbf{r}_1^+ & \mathbf{r}_1^- & \mathbf{s}_1^+ & \mathbf{s}_1^- & \mathbf{r}_2^+ & \mathbf{r}_2^- & \mathbf{s}_2^+ & \mathbf{s}_2^- & \cdots & \mathbf{r}_N^+ & \mathbf{r}_N^- & \mathbf{s}_N^+ & \mathbf{s}_N^- \end{bmatrix} \geq \mathbf{0}.$$

$B = [\partial_{p+1}]$

✗ constraint matrix not TU even if $[\partial_{p+1}]$ is

Median Shape: Examples



$[\partial_2]$ is TU, but LP matrix is not

Median Shape: Examples



$[\partial_2]$ is TU, but LP matrix is not





$[\partial_2]$ is not TU

Median Shape: Questions

Q3: When/Why does median shape LP give integer solutions for free?

Median Shape: Questions

Q3: When/Why does median shape LP give integer solutions for free?

Q4: Network flow (dual graph) model for median shape?

Median Shape: Questions

Q3: When/Why does median shape LP give integer solutions for free?

Q4: Network flow (dual graph) model for median shape?

Thank You!