

MATH230 - Lecture 16 (03/03/2011)

Offer to possibly get A in MATH 230:

- * If you score 95 or more in the Final, your final score will replace your mid-term score.
- * If you score ≥ 90 in the Final, the weights for mid-term and final would be changed to 15% and 40%, respectively.

Inverse of an $n \times n$ matrix

If $A \in \mathbb{R}^{n \times n}$, and if there exists $B \in \mathbb{R}^{n \times n}$ such that $AB = BA = I_n$, then $A^{-1} = B$. (A is invertible, B is the inverse of A ; A is the inverse of B).

If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then A is invertible if $\det A = ad - bc \neq 0$ (determinant). Then

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Theorem 4,
DL-LAA
page 119

If $A \in \mathbb{R}^{n \times n}$ is invertible, then $A\bar{x} = \bar{b}$ has a unique solution for any $\bar{b} \in \mathbb{R}^n$ given by $\bar{x} = A^{-1}\bar{b}$.

Theorem 5, DL-LAA page 120.

Prob 6, pg 126

$$\begin{aligned} 8x_1 + 5x_2 &= -9 \\ -7x_1 - 5x_2 &= 11 \end{aligned}$$

Use matrix inverse to solve the system.

$$A\bar{x} = \bar{b} \text{ where } A = \begin{bmatrix} 8 & 5 \\ -7 & -5 \end{bmatrix}. \quad \det A = 8 \times -5 - (-7) \times 5 = -40 + 35 = -5 \neq 0$$

So, A^{-1} exists. $A^{-1} = \frac{1}{-5} \begin{bmatrix} -5 & -5 \\ 7 & 8 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ -7/5 & -8/5 \end{bmatrix}.$

Hence the unique solution is $\bar{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = A^{-1}\bar{b} = \begin{bmatrix} 1 & 1 \\ -7/5 & -8/5 \end{bmatrix} \begin{bmatrix} -9 \\ 11 \end{bmatrix} = \begin{bmatrix} 2 \\ -5 \end{bmatrix}$

In MATLAB, the function `inv()` gives the inverse of a matrix.

e.g., $\Rightarrow A_{\text{inv}} = \text{inv}(A)$

Properties of matrix inverses ($A, B \in \mathbb{R}^{n \times n}$, invertible)

$$1. (A^{-1})^{-1} = A$$

$$2. (AB)^{-1} = B^{-1}A^{-1}$$

inverse of product = product of inverses in reverse order

$$3. (A^T)^{-1} = (A^{-1})^T$$

Proof sketch

$$2. (AB)^{-1} = B^{-1}A^{-1}$$

We want $C \in \mathbb{R}^{n \times n}$ such that

$$C(AB) = (AB)C = I.$$

Can just verify that $C = B^{-1}A^{-1}$ works.

$$(B^{-1}A^{-1})(AB) = B^{-1}(A^{-1}A)B = B^{-1}IB = B^{-1}B = I$$

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I$$

$$3. (A^T)^{-1} = (A^{-1})^T$$

We want C such that $CA^T = A^TC = I$.

$$C = (A^{-1})^T \text{ works!}$$

$$(A^{-1})^T A^T = (AA^{-1})^T = I^T = I$$

$$A^T (A^{-1})^T = (A^{-1}A)^T = I^T = I.$$

as $(AB)^T = B^T A^T$

How to invert an $n \times n$ matrix? ($n \geq 3$)

Say we want to find $B = A^{-1}$. We have $AB = I$.

We know how to solve $A\bar{x} = \bar{b}$

$$[A | \bar{b}] \xrightarrow{\text{EROs}} \text{rref}([A | \bar{b}]).$$

We could just solve for each column of B as a system of n equations in n variables.

$$\boxed{A} \quad \boxed{\begin{matrix} j \\ \bar{b}_j \\ j \end{matrix} B} = \boxed{\begin{matrix} j \\ \bar{e}_j \\ j \end{matrix} I}$$

For \bar{b}_j (j^{th} column of B), we solve $A\bar{b}_j = \bar{e}_j \rightarrow$ the j^{th} unit vector.

The augmented matrix for this system is $[A | \bar{e}_j]$. We could combine the augmented matrices for all j , and look at $[A | I]$, and perform EROs on it.

We'll get $[A | I] \xrightarrow{\text{EROs}} [I | A^{-1}]$ if A^{-1} exists.

Theorem 7 DL-LAA pg 123

$A \in \mathbb{R}^{n \times n}$ is invertible if and only if it is row-equivalent to I_n (i.e., you can reduce A to I_n using EROs). In this case, the same EROs reduce I_n to A^{-1} . So, $\text{rref}([A | I_n]) = [I_n | A^{-1}]$

If we do not get I_n in place of A , then A is not invertible.

Prob 32 pg 127

$$A = \begin{bmatrix} 1 & -2 & 1 \\ 4 & -7 & 3 \\ -2 & 6 & -4 \end{bmatrix}$$

Find A^{-1} if it exists.

$$[A | I] = \left[\begin{array}{ccc|ccc} 1 & -2 & 1 & 1 & 0 & 0 \\ 4 & -7 & 3 & 0 & 1 & 0 \\ -2 & 6 & -4 & 0 & 0 & 1 \end{array} \right] \xrightarrow[R_3+2R_1]{R_2-4R_1} \left[\begin{array}{ccc|ccc} 1 & -2 & 1 & 1 & 0 & 0 \\ 0 & 1 & -1 & -4 & 1 & 0 \\ 0 & 2 & -2 & 2 & 0 & 1 \end{array} \right] \xrightarrow[R_3-2R_2]{R_1+2R_2}$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & -1 & -7 & 2 & 0 \\ 0 & 1 & -1 & -4 & 1 & 0 \\ 0 & 0 & 0 & 10 & -2 & 1 \end{array} \right]$$

A is not invertible, as

$$A \not\sim I_3.$$

" \sim ": row-equivalent

\hookrightarrow not row-equivalent

Prob 14 pg 126

$B, C \in \mathbb{R}^{m \times n}$, D is invertible, and $(B-C)D = O$.

Show that $B = C$.

Since $(B-C)D$ is defined, D has n rows. But D is invertible, so D is $n \times n$.

$$\underbrace{(B-C)}_{m \times n} \underbrace{D}_{n \times n} = \underbrace{O}_{m \times n}$$

D^{-1} exists. Multiply the above equation on the right by D^{-1} .

$$\underbrace{(B-C)D}_{m \times n} \underbrace{D^{-1}}_{n \times n} = \underbrace{O}_{m \times n} \underbrace{D^{-1}}_{n \times n}$$

$$\Rightarrow (B-C)(DD^{-1}) = OD^{-1} = O$$

$$\Rightarrow (B-C)I = O, \text{ i.e., } B-C = O$$

$$\Rightarrow B = C.$$

Prob 18, pg 126

P is invertible, $A = PBP^{-1}$. Solve for B .

$$(A = PBP^{-1})P$$

$$AP = PBP^{-1}P = PBI = PB$$

Multiply on left with P^{-1} to get

$$P^{-1}AP = \underbrace{P^{-1}P}B = IB = B.$$

So $B = P^{-1}AP$

For $A, B \in \mathbb{R}^{n \times n}$, if there exists an invertible matrix $P \in \mathbb{R}^{n \times n}$ such that $A = P^{-1}BP$, then A and B are similar. Then $B = PAP^{-1}$.

Prob 20, pg 126

$A, B, X \in \mathbb{R}^{n \times n}$, A, X and $(A - AX)$ are invertible, and

$$(A - AX)^{-1} = X^{-1}B \quad \text{--- (1)}$$

(a) Is B invertible?

(b) Solve for X from (1).