# MATH 401: Lecture 1 (08/19/2025)

This is Introduction to Analysis I

I'm Bala Krishnamoorthy (Call me Bala).

Today: * syllabus, logistics → see the course web page for details
* proof techniques
  - contrapositive proof
  - proof by contradiction
  - proof by induction

---

Book: Lindstrøm: Spaces–An Intro to Real Analysis (LSIRA)

LSIRA 1.1

Logical statements and notation.

If A then B (or $A \Rightarrow B$) "implies"

$A \Rightarrow B$ typically does not mean $B \Rightarrow A$.

e.g., A: $p$ a natural number, is divisible by 6

B: $p$ is divisible by 3.

$A \Rightarrow B$ holds, but $B \not\Rightarrow A$ (B does not imply A),

e.g., $p = 9$.

But if $A \Rightarrow B$ and $B \Rightarrow A$ hold, we say A $\underbrace{\text{if and only if}}_{\text{iff}}$ B, or

$A \Leftrightarrow B$ (or A is equivalent to B).

To prove $A \Leftrightarrow B$, we often prove $A \Rightarrow B$ and $B \Rightarrow A$ ($A \Leftarrow B$) separately.

We start by reviewing certain standard techniques to construct proofs of mathematical statements.

# 1. Contrapositive Proof

To show $A \Rightarrow B$, equivalently show not $B \Rightarrow$ not $A$ $(\neg B \Rightarrow \neg A)$.

"negation" or "not"

"If A happened then B happened"
This statement is equivalent to
"If B did not happen then A did not happen."

<u>LSIRA 1.1 Prob 3.</u> Prove the following Lemma.

**<u>Lemma 1</u>** If $n$ is a natural number such that $n^2$ is divisible by 3, then $n$ is divisible by 3.

This is $A \Rightarrow B$ where $A$: $3 \mid n^2$ ($n^2$ is divisible by 3). or 3 divides $n^2$

$B$: $3 \mid n$ ($n$ is divisible by 3).

Let's try to prove $A \Rightarrow B$ directly:

$n^2 = 3k$ (as $n^2 \mid 3$) $\Rightarrow$ $n = \sqrt{3}\sqrt{k}$ (taking square root on both sides)

Hard to conclude that $n \mid 3$ ☹! Would have to argue $k \mid 3$, which is not obvious!

Let's try proving $\neg B \Rightarrow \neg A$.

$\neg B$: $n$ is not divisible by 3.

$\Rightarrow$ $n = 3p+1$ or $n = 3q+2$, for $p, q \in \mathbb{N}$. set of natural numbers

Case 1. $n = 3p+1$

$$\Rightarrow n^2 = (3p+1)^2$$
$$= 9p^2 + 6p + 1$$
$$= 3(3p^2+2p)+1$$
$$= 3k+1 \text{ for } k = 3p^2+2p$$

$\Rightarrow n^2$ is not divisible by 3

Case 2. $n = 3q+2$

$$\Rightarrow n^2 = (3q+2)^2$$
$$= 9q^2 + 12q + 4$$
$$= 9q^2 + 12q + 3 + 1$$
$$= 3\underbrace{(3q^2+4q+1)}_{=k'}+1$$
$$= 3k'+1$$

$\Rightarrow n^2$ is not divisible by 3.

Hence we have proved that if $n$ is not divisible by 3, then $n^2$ is not divisible by 3. Hence, by the contrapositive, we have $n^2 \mid 3 \Rightarrow n \mid 3$. $\square$

Should we always try to build a contrapositive proof?
Not necessarily! In cases where $A \Rightarrow B$ could be concluded directly, the contrapositive argument might make life harder!
It is one of the different proof approaches that you should be aware of.

## 2. Proof by Contradiction

Assume opposite of what you want to prove, and end up with a contradiction (or an obviously wrong statement). Hence the original assumption must be wrong, i.e., you have proved the statement.

<u>LSIRA 1.1 Prob 3 (continued)</u> Prove the following Theorem.

<u>Theorem 2</u> $\sqrt{3}$ is irrational.

Assume $\sqrt{3}$ is rational. → the opposite of what you want to prove

$\Rightarrow \left(\sqrt{3} = \frac{p}{q}\right)^2$, $p, q \in \mathbb{N}$ with no common factors. → by definition, any positive rational number can be written in the form $p/q$ as specified.

Let's square both sides, and cross multiply.

$\Rightarrow 3q^2 = p^2 \Rightarrow 3 \mid p^2$ ($p^2$ is divisible by 3).

Hence by Lemma 1, $3 \mid p$. Let $p = 3k$. ($k \in \mathbb{N}$). Plug $p=3k$ back in:

$\Rightarrow 3q^2 = (3k)^2 = 9k^2$ (divide both sides by 3)

$\Rightarrow q^2 = 3k^2$, i.e., $3 \mid q^2$ ($q^2$ is divisible by 3).

Again by Lemma 1, $3 \mid q$.

Since we started with the assumption that $p$ and $q$ have no common factors

Thus $p$ and $q$ have a common factor of 3, which is a contradiction.

Hence $\sqrt{3}$ is irrational. $\square$

# 3. Proof by Induction

To show a statement $P(n)$ holds for all $n \in \mathbb{N}$,

1. show $P(1)$ holds;
2. Assume $P(k)$ holds for some $k \in \mathbb{N}$.
3. Show $P(k+1)$ holds under Assumption 2.

<u>Example</u>

Show that $P(n) = 3 + 5 + \cdots + 2n+1 = n(n+2)$ $\forall n \in \mathbb{N}$. "for all"

1. $P(1) = 3 = 1(1+2)$ (so $P(1)$ is true).
2. Assume $P(k) = k(k+2)$ for some $k \in \mathbb{N}$.
3. $P(k+1) = P(k) + 2(k+1)+1 = P(k) + 2k+3$

$$= k(k+2) + 2k+3 \quad \text{by induction assumption.}$$

$$= k(k+2) + k + k+3$$

$$= k(k+3) + k+3$$

$$= (k+1)(k+3) = n(n+2) \text{ for } n = k+1.$$

$\Rightarrow P(n) = n(n+2)$ $\forall n \in \mathbb{N}$. $\square$

# MATH 401: Lecture 2 (08/21/2025)

Today: * sets and operations

## Sets and Operations (LS IRA 1.2)

Set: Collection of mathematical objects.

They can be finite, e.g., $\{2,5,9,1,6\}$, or infinite, e.g., $[0,1]$, the collection of all $x \in \mathbb{R}$ with $0 \le x \le 1$.

"element of" set of all real numbers

Given sets $A, B$ we have

$A \subseteq B$: $A$ is a subset of, or equal to, $B$.

$A \subset B$: $A$ is a strict subset of $B$, i.e., there is at least one $x \in B$ such that $x \notin A$.
But $\forall x \in A$, $x \in B$ holds.

To prove $A = B$, we often prove $A \subseteq B$ and $A \supseteq B$ (or $B \subseteq A$).

Here are some standard sets we will use regularly.

$\emptyset$: empty set.

$\mathbb{N} = \{1,2,3,\dots\}$, set of all natural numbers

$\mathbb{R}$ = set of all real numbers

$\mathbb{Z} = \{\dots,-2,-1,0,1,2,\dots\}$, set of all integers

$\mathbb{Q}$ = set of rational numbers, $\mathbb{C}$ = set of complex numbers.

$\mathbb{R}^n$: set of all real $n$-tuples, or $n$-vectors

Notation for sets: $[-2,1] = \{x \in \mathbb{R} \mid -2 \le x \le 1\}$.

closed interval from $-2$ to $1$

"such that" could also use ":" instead of "|".

More generally, $A = \{a \in B \mid P(a)\}$.

bigger set than $A$ property

# Union and Intersection

If $A_i$ are sets for $i=1,\dots,n$, i.e., $A_1, A_2, \dots, A_n$ are sets, then

$$\bigcup_{i=1}^{n} A_i = A_1 \cup A_2 \cup \cdots \cup A_n = \{a \mid a \in A_i \text{ for } \underline{\text{at least}} \text{ one } i\}$$ is their union,

"for all"

$$\bigcap_{i=1}^{n} A_i = A_1 \cap A_2 \cap \cdots \cap A_n = \{a \mid a \in A_i \ \forall i\}$$ is their intersection.

universe $\to \mathcal{U}$




$A \cup B$




$A \cap B$




**LSIRA 1.2 Prob 1** Show $[0,2] \cup [1,3] = [0,3]$.




We show $[0,2] \cup [1,3] \subseteq [0,3]$ and
$[0,2] \cup [1,3] \supseteq [0,3]$.

($\subseteq$) Let $x \in [0,2] \cup [1,3]$

$\Rightarrow x \in [0,2]$ or $x \in [1,3]$ (definition of $\cup$).

$x \in [0,2] \Rightarrow x \in [0,3]$ (as $[0,3]$ contains $[0,2]$)

$x \in [1,3] \Rightarrow x \in [0,3]$. In either case, $x \in [0,3]$.

Hence $[0,2] \cup [1,3] \subseteq [0,3]$.

The result is an obvious one. But we go through the steps of a formal proof more for practice!

($\supseteq$) Let $x \in [0,3]$. Hence $0 \le x \le 3$. Then we get that either $x \le 2$, and hence $x \in [0,2]$, or $x \in (2,3]$.

But if $x \in (2,3]$ then $x \in [1,3]$ (as $[1,3]$ includes $(2,3]$).

$\Rightarrow x \in [0,2] \cup [1,3]$.

Hence $[0,3] \subseteq [0,2] \cup [1,3]$. $\square$

# Distributive Laws of Union and Intersection

For all sets $B, A_1, \dots, A_n$, we have

LSIRA (1.2.1) $$B \cap (A_1 \cup A_2 \cup \dots \cup A_n) = (B \cap A_1) \cup (B \cap A_2) \cup \dots \cup (B \cap A_n).$$

Using more compact notation, we can write

$$B \cap \left( \bigcup_{i=1}^{n} A_i \right) = \bigcup_{i=1}^{n} (B \cap A_i)$$

<u>Proof</u>

We will prove

$$B \cap (A_1 \cup \dots \cup A_n) \subseteq (B \cap A_1) \cup \dots \cup (B \cap A_n), \text{ and}$$

$$B \cap (A_1 \cup \dots \cup A_n) \supseteq (B \cap A_1) \cup \dots \cup (B \cap A_n).$$

('$\subseteq$') Let $x \in B \cap (A_1 \cup \dots \cup A_n)$.

$\Rightarrow x \in B$ and $x \in (A_1 \cup \dots \cup A_n)$ (definition of $\cap$)

$\Rightarrow x \in B$ and $x \in A_i$ for at least one $A_i$. (defn. of $\cup$)

$\Rightarrow x \in B \cap A_i$ for at least one $A_i$.

$\Rightarrow x \in (B \cap A_1) \cup \dots \cup (B \cap A_n)$.

('$\supseteq$') Let $x \in (B \cap A_1) \cup \dots \cup (B \cap A_n)$.

$\Rightarrow x \in (B \cap A_i)$ for at least one $A_i$.

$\Rightarrow x \in B$ and $\underbrace{x \in A_i \text{ for at least one } A_i}$

$\Rightarrow x \in B$ and $x \in (A_1 \cup \dots \cup A_n)$

$\Rightarrow x \in B \cap (A_1 \cup \dots \cup A_n)$. $\square$

LSIRA (1.2.2) is assigned in Homework 1.

# Set Difference and Complement

We write $A \setminus B$ or $A - B$ "setminus"

Caution!
* $A \setminus B \neq B \setminus A$ !

"A setminus B" is $A \setminus B = \{a \mid a \in A, a \notin B\}$.

If $\mathcal{U}$ is the universe, i.e., $A \subseteq \mathcal{U}$ for all sets $A$, then

$$A^c = \mathcal{U} \setminus A = \{a \in \mathcal{U} \mid a \notin A\}$$ is the

complement of $A$ (or $A$-complement).




# De Morgan's Laws

LSIRA (1.2.3) $(A_1 \cup \cdots \cup A_n)^c = A_1^c \cap \cdots \cap A_n^c$. "complement of union = intersection of complements"

LSIRA (1.2.4) $(A_1 \cap \cdots \cap A_n)^c = A_1^c \cup A_2^c \cup \cdots \cup A_n^c$. complement of intersection = union of complements.

See LSIRA for the proof.

Let's illustrate (1.2.4) for $n=2$, i.e., with $A_1$ and $A_2$ first.



We will prove subset inclusion in both directions.

($\subseteq$) Let $x \in (A_1 \cap \cdots \cap A_n)^c$

$\Rightarrow x \notin A_1 \cap \cdots \cap A_n$ (definition of complement)

$\Rightarrow x \notin A_j$ for some $j$. (definition of $\cap$)

$\Rightarrow x \in A_j^c$ for some $j$

$\Rightarrow x \in A_1^c \cup \cdots \cup A_n^c$.

Hence $(A_1 \cap \cdots \cap A_n)^c \subseteq A_1^c \cup \cdots \cup A_n^c$.

($\supseteq$) Let $x \in A_1^c \cup \cdots \cup A_n^c$.

$\Rightarrow x \in A_j^c$ for some $j$

$\Rightarrow x \notin A_j$ for some $j$

$\Rightarrow x \notin A_1 \cap \cdots \cap A_n$. since $x \notin A_j$ for some $j$, it cannot be in the intersection of all $A_i$'s.

$\Rightarrow x \in (A_1 \cap \cdots \cap A_n)^c$.

Hence $A_1^c \cup \cdots \cup A_n^c \subseteq (A_1 \cap \cdots \cap A_n)^c$. $\square$

## Cartesian Products

For $A, B$: sets, we define

$$A \times B = \{(a,b) \mid a \in A, b \in B\}$$

cartesian product of A and B

Given $A_i$, $i = 1, \dots, n$ $(A_1, \dots, A_n)$, we define

compact notation

$\prod$ : product

$$A_1 \times A_2 \times \dots \times A_n = \prod_{i=1}^{n} A_i = \{(a_1, \dots, a_n) \mid \underbrace{a_i \in A_i \ \forall i}_{a_1 \in A_1, a_2 \in A_2, \dots, a_n \in A_n}\}.$$

e.g., $\mathbb{R}^n$: set of $n$-tuples of real numbers
(or set of real $n$-vectors)

<u>LSIRA 1.2 Prob 9 (Pg 11)</u> Prove that $(A \cup B) \times C = (A \times C) \cup (B \times C)$.

We'll finish the proof in the next lecture...

# MATH 401: Lecture 3 (08/26/2025)

Today: * families of sets, properties
* functions, images, pre images

---

We first do a problem on Cartesian products...

LSIRA 1.2 Prob 9 (Pg 11) Prove that $(A \cup B) \times C = (A \times C) \cup (B \times C)$.

'$\subseteq$' Let $(x,y) \in (A \cup B) \times C$.

$\Rightarrow x \in A \cup B,\ y \in C$ (Definition of cartesian product)

$\Rightarrow x \in A$ or $x \in B$, $y \in C$

If $x \in A$ then $(x,y) \in A \times C$, and

if $x \in B$ then $(x,y) \in B \times C$.

$\Rightarrow (x,y) \in A \times C$ or $(x,y) \in B \times C$

$\Rightarrow (x,y) \in (A \times C) \cup (B \times C)$.

'$\supseteq$' Let $(x,y) \in (A \times C) \cup (B \times C)$

$\Rightarrow (x,y) \in A \times C$ or $(x,y) \in B \times C$

$\Rightarrow x \in A, y \in C$ or $x \in B, y \in C \Rightarrow (x \in A$ or $x \in B), y \in C$.

$\Rightarrow x \in A \cup B, y \in C \Rightarrow (x,y) \in (A \cup B) \times C$.

$\square$

# LSIRA 1.3 Families of Sets

Recall: $B \cap \left(\bigcup_{i=1}^{n} A_i\right) = \bigcup_{i=1}^{n} (B \cap A_i)$. → compact notation for distributive law (from Lecture 2)

We could write, instead, $B \cap \left(\bigcup_{i \in \mathcal{I}} A_i\right) = \bigcup_{i \in \mathcal{I}} (B \cap A_i)$, where $\mathcal{I} = \{1, 2, \ldots, n\}$.

We now generalize $\mathcal{I}$ to be infinite in some cases, or indexing more general collections in general.

**Def** A collection of sets is a **family**.

e.g., $\mathcal{A} = \{[a,b] \mid a, b \in \mathbb{R}\}$ is the family of all closed intervals on $\mathbb{R}$.

## Union and Intersection

We extend union, intersection, as well as their distribution to families.

$$\bigcup_{A \in \mathcal{A}} A = \{a \mid a \in A \text{ for some } A \in \mathcal{A}\}.$$ → collection of elements that belong to at least one set in the family

$$\bigcap_{A \in \mathcal{A}} A = \{a \mid a \in A \text{ for all } A \in \mathcal{A}\}$$ → collection of elements that belong to every set in the family.

We naturally extend distributive and De Morgan's laws to families.

$$B \cap \left(\bigcup_{A \in \mathcal{A}} A\right) = \bigcup_{A \in \mathcal{A}} (B \cap A), \quad \left(\bigcap_{A \in \mathcal{A}} A\right)^c = \bigcup_{A \in \mathcal{A}} A^c, \text{ etc.}$$

We now work on some problems involving families of sets.

## LSIRA 1.3 Prob 1 (Pg 12)

Show that $\bigcup_{n\in\mathbb{N}} [-n,n] = \mathbb{R}$.

('$\subseteq$') $\mathbb{R}$ is the universe here, so $\bigcup_{n\in\mathbb{N}} [-n,n] \subseteq \mathbb{R}$.

Or, observe that $[-n,n] \in \mathbb{R}$ for each $n \in \mathbb{N}$, hence $\bigcup_{n\in\mathbb{N}} [-n,n] \subseteq \mathbb{R}$.

('$\supseteq$') Let $x \in \mathbb{R}$. → To be more careful, we could consider $x=0$ separately. Note that $x=0 \in [-n,n]$ $\forall n \in \mathbb{N}$.

Let $m = \lceil |x| \rceil$, ceiling of absolute value of $x$, i.e., the smallest natural number $\geq |x|$.

$\lceil x \rceil = \text{ceil}(x)$ = smallest integer $\geq x$.

Then $x \in [-m,m] = [-\lceil |x| \rceil, \lceil |x| \rceil]$, as

$x \leq |x| \leq \lceil |x| \rceil = m$, and $x \geq -|x| \geq -\lceil |x| \rceil$.

→ e.g., $x = -3.3 \Rightarrow x \geq -|-3.3| = -3.3 \geq -4$.

$\Rightarrow x \in \bigcup_{n\in\mathbb{N}} [-n,n]$.

$\square$

## LSIRA 1.3 Prob 4

Show $\bigcap_{n\in\mathbb{N}} (0,\frac{1}{n}] = \emptyset$ (empty set).

('$\supseteq$') $\emptyset \subseteq A$ for any set $A$ (trivially).

('$\subseteq$') We show $\bigcap_{n\in\mathbb{N}} (0,\frac{1}{n}] \subseteq \emptyset$. → $\emptyset^c = \mathbb{R}$. Hence we show $x \in \mathbb{R}$ is not in $\bigcap_{n\in\mathbb{N}} (0,\frac{1}{n}]$.

For $x \in \mathbb{R}$, we show $x \notin \bigcap_{n\in\mathbb{N}} (0,\frac{1}{n}]$.

If $x \leq 0$, then clearly, $x \notin (0,\frac{1}{n}]$ $\forall n \in \mathbb{N}$.

If $x \geq 1$, then $x \notin (0,\frac{1}{2}]$ for $n=2$, for instance.

Let $0<x<1$. Consider $m=\lceil \frac{1}{x} \rceil+1$.

Then $x \notin (0, \frac{1}{m}]$ as $x > \frac{1}{m} = \frac{1}{\lceil \frac{1}{x} \rceil+1}$. $\left( \lceil \frac{1}{x} \rceil+1 > \frac{1}{x} \right)$

$\Rightarrow \quad x \notin \bigcap_{n \in \mathbb{N}} (0, \frac{1}{n}]$.

**Q.** Why take $\lceil \frac{1}{x} \rceil+1$?

Consider $x=\frac{1}{5}$, for instance.
Then $\lceil \frac{1}{x} \rceil = \lceil 5 \rceil = 5$.
Hence $x \in (0, \frac{1}{m}]$ here!

$\square$

<u>LSIRA 1.3 Prob 5 (Pg 12)</u>

Prove that $B \cup \left( \bigcap_{A \in \mathcal{A}} A \right) = \bigcap_{A \in \mathcal{A}} (B \cup A)$.

($\subseteq$) Let $x \in B \cup \left( \bigcap_{A \in \mathcal{A}} A \right)$

$\Rightarrow x \in B$ or $x \in \bigcap_{A \in \mathcal{A}} A$

$\Rightarrow x \in B$ or $x \in A$ for each $A \in \mathcal{A}$.

$\Rightarrow x \in B \cup A$ for each $A \in \mathcal{A}$.

$\Rightarrow x \in \bigcap_{A \in \mathcal{A}} (B \cup A)$.

($\supseteq$) Let $x \in \bigcap_{A \in \mathcal{A}} (B \cup A)$

$\Rightarrow x \in B \cup A$ for every $A \in \mathcal{A}$.

$\Rightarrow x \in B$ or $\underbrace{x \in A \text{ for every } A \in \mathcal{A}}$.

$\Rightarrow x \in B$ or $x \in \bigcap_{A \in \mathcal{A}} A \Rightarrow x \in B \cup \left( \bigcap_{A \in \mathcal{A}} A \right)$.

$\square$

# LSIRA 1.4 Functions

A function $f: X \to Y$ for sets $X, Y$ is a rule that assigns for each $x \in X$ a **unique** $y \in Y$. We write $f(x) = y$, or

$x \mapsto y$ "maps to".




Rather than the graphs of functions you may have seen previously, we think of such visualizations for functions now.

$X$ is the domain and $Y$ the codomain of $f$.

## Compositions




Let $f: X \to Y$ and $g: Y \to Z$ be functions. Then their composition is specified as $h: X \to Z$ defined as $h(x) = g(f(x))$.
The composition is written as $g \circ f$, with $g \circ f(x) = g(f(x))$.

"composition of f and g"

$f_1(f_2(\cdots f_n(x))))\cdots)$ → composition of $n$ functions $f_1, f_2, \ldots, f_n$

Function: $f: X \to Y$. We now define images and preimages under $f$.




For $A \subseteq X$, $f(A) \subseteq Y$ is defined as

$$f(A) = \{f(a) \mid a \in A\},$$

and is called the **image** of $A$ under $f$.




For $B \subseteq Y$, the set $f^{-1}(B) \subseteq X$ defined as

$$f^{-1}(B) = \{x \in X \mid f(x) \in B\}$$

is the **inverse image** or **preimage** of $B$ under $f$.

In the next lecture, we consider how preimages and images commute with unions and intersections, or not...

# MATH 401: Lecture 4 (08/28/2025)

Today:
* images/preimages and unions/intersections
* injective/surjective functions
* relations, equivalence relations, partitions

---

We now consider how images and inverse images commute (or not) with unions and intersections of families of sets.

<u>Prop 1.4.1</u> Inverse images commute with arbitrary unions and intersections:

$$f^{-1}\Big(\bigcup_{B\in\mathcal{B}} B\Big) = \bigcup_{B\in\mathcal{B}} f^{-1}(B)$$ and

"inverse of union = union of inverses"

$$f^{-1}\Big(\bigcap_{B\in\mathcal{B}} B\Big) = \bigcap_{B\in\mathcal{B}} f^{-1}(B)$$

"inverse of intersection = intersection of inverses"

**Proof** (of the second statement) → See LSIRA for proof of first statement

($\subseteq$) Let $x \in f^{-1}\big(\bigcap_{B\in\mathcal{B}} B\big) \Rightarrow f(x) \in \bigcap_{B\in\mathcal{B}} B$

$\Rightarrow f(x) \in B$ for every $B \in \mathcal{B}$.

$\Rightarrow x \in f^{-1}(B)$ for every $B \in \mathcal{B}$.

$\Rightarrow x \in \bigcap_{B\in\mathcal{B}} f^{-1}(B)$.

($\supseteq$) Let $x \in \bigcap_{B\in\mathcal{B}} f^{-1}(B)$

$\Rightarrow x \in f^{-1}(B)$ for every $B \in \mathcal{B}$.

$\Rightarrow f(x) \in B$ for every $B \in \mathcal{B}$.

$\Rightarrow f(x) \in \bigcap_{B\in\mathcal{B}} B \Rightarrow x \in f^{-1}\big(\bigcap_{B\in\mathcal{B}} B\big)$.

$\square$

We saw that inverse images commute with unions and intersections. But forward images behave a bit differently.

<u>Prop 1.4.2</u> $f: X \to Y$ is a function, $\mathcal{A}$ is a family of subsets of $X$.

Then $f\left(\bigcup_{A\in\mathcal{A}} A\right) = \bigcup_{A\in\mathcal{A}} f(A)$, $f\left(\bigcap_{A\in\mathcal{A}} A\right) \subseteq \bigcap_{A\in\mathcal{A}} f(A)$.

<u>Proof</u>

('$\subseteq$') Let $y \in f\left(\bigcup_{A\in\mathcal{A}} A\right)$

"There exists"

$\Rightarrow \exists\, x \in \bigcup_{A\in\mathcal{A}} A$ such that $f(x) = y$.

$\Rightarrow x \in A$ for at least one $A \in \mathcal{A}$ such that $f(x) = y$

$\Rightarrow y \in f(A)$ for at least one $A \in \mathcal{A}$

$\Rightarrow y \in \bigcup_{A\in\mathcal{A}} f(A)$.

($\supseteq$) Let $y \in \bigcup_{A\in\mathcal{A}} f(A)$.

$\Rightarrow y \in f(A)$ for at least one $A \in \mathcal{A}$

$\Rightarrow \exists\, x \in A$ for at least one $A \in \mathcal{A}$ such that $f(x) = y$.

$\Rightarrow \exists\, x \in \bigcup_{A\in\mathcal{A}} A$ such that $f(x) = y$.

$\Rightarrow y \in f\left(\bigcup_{A\in\mathcal{A}} A\right)$

LSIRA gives a slightly different proof for ($\supseteq$):

"for all"

$A \subseteq \bigcup_{A\in\mathcal{A}} A \quad \forall A \in \mathcal{A}$

$\Rightarrow f(A) \subseteq f\left(\bigcup_{A\in\mathcal{A}} A\right) \quad \forall A \in \mathcal{A}$

Since this result holds for every $A \in \mathcal{A}$, we can write

$\bigcup f(A) \subseteq f\left(\bigcup_{A\in\mathcal{A}} A\right)$.

$\square$

We consider intersections now: $f\left(\bigcap_{A\in\mathcal{A}} A\right) \subseteq \bigcap_{A\in\mathcal{A}} f(A)$.

<u>Proof for ('$\subseteq$')</u>

$$\bigcap_{A\in\mathcal{A}} A \subseteq A \quad \forall A\in\mathcal{A}$$

$$\Rightarrow f\left(\bigcap_{A\in\mathcal{A}} A\right) \subseteq f(A) \ \forall A\in\mathcal{A}.$$

Since this inclusion holds for every $A\in\mathcal{A}$, we get

$$f\left(\bigcap_{A\in\mathcal{A}} A\right) \subseteq \bigcap_{A\in\mathcal{A}} f(A).$$

<u>Counterexample for ('$\supseteq$') for $\cap$</u>




For $x_1 \neq x_2$, $x_1, x_2 \in X$, let $f(x_i) = y$, $i=1,2$.

Let $A_i = \{x_i\}$, $i=1,2$. $\Rightarrow \bigcap_{i=1,2} A_i = \phi$ (empty set).

But note that $f(A_i) = \{y\}$, $i=1,2$.

$\Rightarrow f\left(\bigcap_{i=1,2} A_i\right) = \phi$. But $\bigcap_{i=1,2} f(A_i) = \{y\} \neq \phi$.

$$\Rightarrow \bigcap_{i=1,2} f(A_i) \not\subseteq f\left(\bigcap_{i=1,2} A_i\right).$$

But we get this reverse inclusion if we specify that $f$ is injective.

**Def** Let $f: X \to Y$ be a function.

$f$ is **injective** (1-to-1) if $f(x_1) \neq f(x_2)$ whenever $x_1 \neq x_2$.

Equivalent definition: For any $y \in Y$, there is **at most** one $x \in X$ s.t. $f(x) = y$.
→ there could be no $x \in X$

$f$ is **surjective** (onto) if for every $y \in Y$, there is **at least** one $x \in X$ such that $f(x) = y$.
→ there could be more than one

$f$ is **bijective** if it is both injective and surjective.

## LSIRA 1.4 prob 4 (Pg 17)

Let $f: \overset{X}{\mathbb{R}} \to \overset{Y}{\mathbb{R}}$ be a strictly increasing function, i.e.,

$$x_1 < x_2 \Rightarrow f(x_1) < f(x_2) \text{ for } x_i \in \mathbb{R},\ i = 1, 2.$$

1. Show that $f$ is injective.
2. Does it have to be surjective?

→ Either give a proof or a counterexample.

→ The same result holds when $x_2 < x_1$ as well.

1. We show $f(x_1) \neq f(x_2)$ whenever $x_1 \neq x_2$.
   Without loss of generality (WLOG), let $x_1 < x_2$.
   Then $f(x_1) < f(x_2)$, as $f$ is strictly increasing.
   Hence $f(x_1) \neq f(x_2)$, and so $f$ is injective.

2. No. $f = \arctan(x)$ is strictly increasing.
   $f: \mathbb{R} \to \mathbb{R}$, but $\arctan(\mathbb{R}) = (-\frac{\pi}{2}, \frac{\pi}{2}) \subset \mathbb{R}$.
   So $f$ need not be surjective.
   Another example is $f = e^x$.

# Relations (LSIRA 1.5)

We had seen functions, where a unique $y \in Y$ is assigned for each $x \in X$ by $f: X \to Y$. But entities are related in other ways — numbers are $>$ or $<$ each other, lines are parallel, etc. We define relations formally to describe such dependencies.

**Def** A relation $R$ on a set $X$ is a subset of $\underbrace{X \times X}_{\text{Cartesian product of } X \text{ with itself}}$.

We write $xRy$, $(x,y) \in R$, or $x \sim y$.

e.g.; $R = \{(x,y) \in \mathbb{R}^2 \mid x = y\}$.




Recall, $y=x$ is the $45°$ line through $(0,0)$. All points are "related" by them belonging to this line.

Here is another relation (on integers):

$P = \{(x,y) \in \mathbb{Z}^2 \mid x, y \text{ have same parity}\}$.

So, all odd integers are related, and so are all even integers.

Some relations have more structure than default – as defined below.

## Equivalence Relations

**Def** A relation $\sim$ on $X$ is an **equivalence relation** if it is

(i) reflexive, i.e., $x \sim x \ \forall x \in X$;

(ii) symmetric, i.e., $x \sim y \Rightarrow y \sim x \ \forall x, y \in X$; and

(iii) transitive, i.e., $x \sim y,\ y \sim z \Rightarrow x \sim z \ \forall x, y, z \in X$.

Note that $<$ is not reflexive, or symmetric, e.g., $5 \not< 5$, and $3 < 5 \not\Rightarrow 5 < 3$.

**Def** Given an equivalence relation $\sim$ on $X$, we define the **equivalence class** $[x]$ of $x \in X$ as

$$[x] = \{y \in X \mid x \sim y\}.$$

the set of all "relatives" of $x$

The collection of equivalence classes forms a <u>partition</u> of $X$.

**Def** A **partition** $\mathcal{P}$ of $X$ is a family of nonempty subsets of $X$ such that $x \in X$ satisfies $x \in P \in \mathcal{P}$ for exactly one $P$ in $\mathcal{P}$ (for every $x \in X$).

The elements $P$ of $\mathcal{P}$ are called **partition classes** of $\mathcal{P}$.

e.g., $\mathcal{P} = \{\underbrace{\{2k, k \in \mathbb{Z}\}}_{\text{even integers}}, \underbrace{\{2k+1, k \in \mathbb{Z}\}}_{\text{odd integers}}\}$ is a partition of $\mathbb{Z}$.

Here is a direct example of a partition of $\mathbb{R}^2$.

The collection of all lines with slope $= 1$ ($45^\circ$) is a partition of $\mathbb{R}^2$.




there are infinitely many lines with slope $m=1$.

Any point in $\mathbb{R}^2$ belongs to exactly one line with a slope of $m=1$ (i.e., $45^\circ$ degree slope).

We have not checked that the defining relation is an equivalence relation, but this can be done easily.

recall, the point-slope form of the equation of a line: $\frac{y-y_0}{x-x_0} = m$, given slope $m$ and one point $(x_0, y_0)$.

# MATH 401: Lecture 5 (09/02/2025)

Today: * equivalence relations and partitions
* countability

---

**Recall:** * $\sim$ is an equivalence relation on $X$: $x \sim x$, $x \sim y \Rightarrow y \sim x$, $x \sim y, y \sim z \Rightarrow x \sim z$
* partition of $X$ $\mathcal{P} = \{P\}$

We show that equivalence relations naturally define partitions.

**Prop 1.5.3** If $\sim$ is an equivalence relation on $X$, then the collection of equivalence classes $\mathcal{P} = \{[x] \mid x \in X\}$ is a partition of $X$.

**Proof** We show each $x \in X$ belongs to exactly one equivalence class.

$x \sim x$ $\sim$ is equivalence relation, so is reflexive ((i))

$\Rightarrow x \in [x]$ $\longrightarrow$ So, each $x \in X$ belongs to at least its own class.

We now show if $x \in [y]$ for $y \in X$, $y \neq x$, then $[x] = [y]$.

We show $[x] \subseteq [y]$ and $[x] \supseteq [y]$.

($\subseteq$) Let $z \in [x]$

$\Rightarrow x \sim z$ Definition of $[x]$

We assumed $x \in [y] \Rightarrow y \sim x$

$\sim$ is an equivalence relation, so $y \sim x, x \sim z \Rightarrow y \sim z$. ($\sim$ is transitive ((iii))

$\Rightarrow z \in [y]$.

($\supseteq$) Let $z \in [y]$ $\Rightarrow y \sim z$

Also, $x \in [y] \Rightarrow y \sim x$

$\sim$ is equivalence relation $\Rightarrow x \sim y$ ($\sim$ is symmetric (ii))

$\Rightarrow x \sim y, y \sim z \Rightarrow x \sim z$ ($\sim$ is transitive (iii))

$\Rightarrow z \in [x]$. $\square$

LSIRA 1.5 Prob 5 (Pg 20) Let $\sim$ be a relation on $\mathbb{R}^3$ defined as

$$(x,y,z) \sim (x',y',z') \iff 3x-y+2z = 3x'-y'+2z'.$$

Show that $\sim$ is an equivalence relation. Describe its equivalence classes.

We check that $\sim$ is reflexive, symmetric, and transitive.

Reflexive: $(x,y,z) \sim (x,y,z)$, as $3x-y+2z = 3x-y+2z$. ✓

Symmetric: $(x,y,z) \sim (x',y',z') \Rightarrow (x',y',z') \sim (x,y,z)$ holds as

$$3x-y+2z = 3x'-y'+2z' \Rightarrow$$
$$3x'-y'+2z' = 3x-y+2z. \checkmark$$

$a=b \Rightarrow b=a$ for $a,b \in \mathbb{R}$.

Transitive: $(x,y,z) \sim (x',y',z')$ and $(x',y',z') \sim (x'',y'',z'')$

$\Rightarrow (x,y,z) \sim (x'',y'',z'')$ also holds, as

$$3x-y+2z = 3x'-y'+2z' \text{ and } 3x'-y'+2z' = 3x''-y''+2z''$$
$$\Rightarrow 3x-y+2z = 3x''-y''+2z''. \checkmark$$

$$[(x,y,z)] = \{(x',y',z') \in \mathbb{R}^3 \mid 3x-y+2z = 3x'-y'+2z'\}$$

If we set $3x-y+2z = d \in \mathbb{R}$, then

$$[(x,y,z)] = \{\underline{(x',y',z') \in \mathbb{R}^3 \mid 3x'-y'+2z' = d}\}$$

plane with normal vector $(3,-1,2)$ (or $\begin{bmatrix}3\\-1\\2\end{bmatrix}$) through $(x,y,z)$.

We can describe the equivalence classes as follows.
The equivalence class of a point in $\mathbb{R}^3$ is the plane with normal $(3,-1,2)$ passing through that point.

We write $\mathbb{R}^3/\sim$ for the set of all equivalence classes of $\sim$.

**Def** If $\sim$ is an equivalence relation on $X$, then $X/\sim$ is the set of all equivalence classes under $\sim$. "X quotient $\sim$"

$\mathbb{R}^3/\sim$ here is the set of all planes with normal $(3,-1,2)$.

Note that any point $(x,y,z) \in \mathbb{R}^3$ belongs to exactly one plane with normal $(3,-1,2)$. Also, all such parallel planes together cover all of $\mathbb{R}^3$, i.e., $\mathbb{R}^3/\sim$ is indeed a partition of $\mathbb{R}^3$. Note the similarity to previous example of $45^\circ$ lines in $\mathbb{R}^2$.

<u>Another example on equivalence classes and Partitions</u>

Let $X$ be the set of all fruits in a grocery store. We can group them into fruit types (classes), e.g., apples, citrus, grapes, tomatoes, plums, etc. Note that apples could include honeycrisp, red delicious, etc. (varities of apples)

$\mathcal{P}$: A partition of $X$ into fruit classes may look like this $\longrightarrow$




$$\mathcal{P} = \{\underbrace{\text{apples}}_{P_1}, \underbrace{\text{grapes}}_{P_2}, \underbrace{\text{citrus}}_{P_3}, \underbrace{\text{plums}}_{P_4}, \ldots\}$$

Note that any individual fruit belongs to exactly one class. $\mathcal{P}$ is indeed a partition of $X$.

<u>Equivalence relation $\sim$ on $X$ associated with $\mathcal{P}$</u>

For fruits $x, y$, $x \sim y$ if $x$ and $y$ are the same fruit type.
$\sim$ is indeed an equivalence relation (can check its reflexive, symmetric, transitive).

What is the equivalence class $[x]$ of a fruit $x$?

$[x]$ is the set of all fruits of its type in the store.
e.g., $x$ = Valencia orange, $[x] = \{$set of all citrus fruits$\}$.

What is the quotient space $X/\sim$? $X/\sim$ is the set of all fruit types.

So $X/\sim = \{$apples, citrus, $\ldots\}$

Check all problems on equivalence relations from LSIRA.

# LSIRA 1.6 Countability

We typically count a set of objects as 1, 2, 3, ..., i.e., by numbering or indexing the first element, then the second one, etc. We can talk about sets being countable (or not) in general.

**Def** A set $A$ is **countable** if it is possible to list all elements of $A$ as $a_1, a_2, \ldots, a_n, \ldots$

e.g., $\mathbb{N}$ is countable — just list the elements as $1, 2, 3, \ldots$.

set of natural numbers

We could use a little more formal definition of a countable set, than the one given above (as listed in LSIRA).

**Def** A set $A$ is countable if there exists an injective function $f: A \to \mathbb{N}$.

The function $f$ is the "indexing" or "numbering" function that assigns a separate natural number to each element of $A$.

Note that finite sets are always countable — we can always list the elements in a sequence. Things are more interesting for infinite sets.

**Def** If $f$ is also surjective, i.e., it is bijective, then $A$ is **countably infinite**, i.e., it is countable and is infinite.

e.g., $\mathbb{Z}$ is countable.

We can list all integers as

$$0, 1, -1, 2, -2, 3, -3, \ldots$$

This is just one way to list all integers. Other ways could be devised as well.

| index | 1 | 2 | 3 | 4 | 5 | 6 | 7 | ... |
|---|---|---|---|---|---|---|---|---|

Note how the indices are listed. The positive integers are the even entries in the list, and negative integers (& 0) are the odd entries in the list

Or, we can define $f: \mathbb{Z} \to \mathbb{N}$ as

$$f(z) = \begin{cases} 2z, & z > 0 \\ 1-2z, & z \leq 0 \end{cases}$$

We can specify $f^{-1}(\cdot)$ as follows:

$$f^{-1}(n) = \begin{cases} n/2, & n \text{ even} \\ \frac{-n+1}{2}, & n \text{ odd.} \end{cases}$$

$f$ is bijective, and hence $\mathbb{Z}$ in countably infinite.

**Proposition 1.6.1** If $A, B$ are countable, then so is $\underline{A \times B}$.
↳ Cartesian product

$A, B$ are countable $\Rightarrow \exists$ lists $\{a_n\}, \{b_n\}$ containing all elements of $A$ and $B$, respectively.

$$\Rightarrow \{\underbrace{(a_1,b_1)}_{1+1}, \underbrace{(a_1,b_2), (a_2,b_1)}_{\substack{1+2 \\ =3}\quad \substack{2+1 \\ =3}}, \underbrace{(a_1,b_3), (a_2,b_2), (a_3,b_1)}_{\substack{1+3 \\ =4}\quad \substack{2+2 \\ =4}\quad \substack{3+1 \\ =4}}, \dots\}$$

index

is a list containing all elements of $A \times B$.

Note the index trick: we list pairs of elements $(a_i, b_j)$ with $a_i \in \{a_n\}$ and $b_j \in \{b_n\}$ such that the sum of their indices increase as natural numbers. Thus, $i+j=2$, and then all options for $i+j=3$, followed by all options for $i+j=4$, and so on.

This index trick could be used to show other sets are countable, e.g., the cartesian product of $k$ countable sets is countable. ($A_1 \times A_2 \times \dots \times A_k$, where $A_i$ is countable for $1 \leq i \leq k$).

LSIRA 1.6 Prob 1 (pg 22) Show that the subset of a countable set is countable.

Let $B \subset A$, where $A$ is countable.

As $A$ is countable, there is a list $a_1, a_2, \dots, a_n, \dots$ such that every $a_i \in A$ is included in the list.

from the list

Let $n_1 \in \mathbb{N}$ be the smallest natural number such that $a_{n_1} \in B$.

And let $n_2 \in \mathbb{N}$, $n_2 > n_1$, be the smallest number such that $a_{n_2} \in B$, and let $n_3 > n_2$, $n_3 \in \mathbb{N}$, be the smallest number such that $a_{n_3} \in B$, and so on.

We form a new list with $b_i = a_{n_i}$, $i = 1, 2, 3, \dots$

$\Rightarrow$ $b_1, b_2, b_3, \dots$ is a listing of all elements in $B$, ensuring that $B$ is countable.

↳ indeed, we will miss no elements of $B$ in this process, and all of them are included in the new list. $\square$

Check Prop 1.6.2: $\bigcup_{n \in \mathbb{N}} A_n$ is countable when $A_n$ is countable $\forall n$.
(in LSIRA)

We can use a similar indexing trick as in Prop. 1.6.1.

Countability is one way to compare two infinite sets. We know $\mathbb{R} \supseteq \mathbb{Q}$, but both have infinitely many elements. Intuitively, we know $\mathbb{R}$ is bigger as it contains irrational numbers in addition to rationals.

We'll first show that $\mathbb{Q}$ is countable, but $\mathbb{R}$ is, in fact, uncountable. More in the next lecture...

# MATH 401: Lecture 6 (09/04/2025)

Today: * $\mathbb{Q}$ is countable, $\mathbb{R}$ is uncountable
* $\epsilon$-$\delta$ proofs, convergence

---

**Recall:** **Proposition 1.6.1** If $A, B$ are countable, then so is $A \times B$.

**Proposition 1.6.3** $\mathbb{Q}$ is countable.

↳ set of all rational numbers, $\frac{p}{q}$ for $p \in \mathbb{Z}$, $q \in \mathbb{N}$

This representation includes all negative rationals. Also, $q \in \mathbb{N}$ avoids $q = 0$.

We first observe that $\mathbb{Z} \times \mathbb{N}$ is countable, as we showed that $\mathbb{Z}$ and $\mathbb{N}$ are both countable, and then applying Proposition 1.6.1.

$\Rightarrow$ $\mathbb{Z} \times \mathbb{N}$ can be listed as, for instance,
$\{\{(a_1, b_i)\}_{i=1}^{\infty}, \{(a_2, b_i)\}_{i=1}^{\infty}, \ldots, \{(a_k, b_i)\}_{i=1}^{\infty}, \ldots\}$ where $\{a_n\}$ and $\{b_n\}$ are listings for $\mathbb{Z}$ and $\mathbb{N}$, respectively.

But $\left\{\left\{\frac{a_1}{b_i}\right\}_{i=1}^{\infty}, \left\{\frac{a_2}{b_i}\right\}_{i=1}^{\infty}, \ldots, \left\{\frac{a_k}{b_i}\right\}_{i=1}^{\infty}, \ldots\right\}$ is a listing of $\mathbb{Q}$. $\square$

Let's consider any rational number, e.g., $\frac{2}{5}$.
How many times does $\frac{2}{5}$ appear in this listing? Once, exactly as $\frac{2}{5}$.

But infinitely many times as a value, because $\frac{2}{5} = \frac{4}{10} = \frac{20}{50} = \cdots$

In fact, every rational number appears infinitely many times in this list. But that is not a problem for countability.

We now show that the set of all reals is uncountable.

**Theorem 1.64** $\mathbb{R}$ is uncountable.

Consider $[0,1] \subset \mathbb{R}$. We show that $[0,1]$ is uncountable.
To get a contradiction, assume that $[0,1]$ is countable.

As there are infinitely many real #s between 0 and 1.
$[0,1]$ is a countably infinite set (under assumption).

We can list all these real numbers as follows:

Note that each number has infinitely many decimal digits (they could be all zeros after some number of places)

$$r_1 = 0.a_{11}\, a_{12}\, a_{13} \cdots$$
$$r_2 = 0.a_{21}\, a_{22}\, a_{23} \cdots$$
$$r_3 = 0.a_{31}\, a_{32}\, a_{33} \cdots$$
$$\vdots \qquad \vdots$$

$a_{ij} =$ $j^{th}$ decimal digit in the $i^{th}$ real number (in the list).

$a_{ij} \in \{0,1,2,\ldots,9\}$.

We create a new real number in $[0,1]$ as follows.

$$s = 0.d_1 d_2 d_3 \cdots \quad \text{where}$$

$$d_i = \begin{cases} 1 & \text{if } a_{ii} \neq 1, \text{ and} \\ 2 & \text{if } a_{ii} = 1. \end{cases}$$

e.g.,
$$r_1 = 0.02534\cdots$$
$$r_2 = 0.8176\cdots$$
$$r_3 = 0.30914\cdots$$
$$r_4 = 0.00207\cdots$$
$$\vdots$$

Then $s = 0.1211\ldots$

Note that $s$ has infinitely many decimal digits.

So, $s$ is different from $r_i$ for each $i$.
This contradicts the assumption that $\{r_i\}$ contains **every** real number in $[0,1]$. Hence $[0,1]$ is uncountable.

Since $\mathbb{R} \supset [0,1]$, and $[0,1]$ is uncountable, $\mathbb{R}$ is also uncountable. $\square$

This is a standard trick we use to show a set is uncountable. We assume it is countable, and start with a listing. Then we identify an element that is distinct from every element in the listing, contradicting the assumption that the listing includes all such elements.

**Corollary.** The set of irrational numbers is uncountable.

We showed $\mathbb{Q}$ is countable, and $\mathbb{R}$ is uncountable.
The set of irrationals $= \mathbb{R}/\mathbb{Q}$ is hence uncountable.

# LSIRA Chapter 2 Foundations of Calculus

## 2.1. $\epsilon$-$\delta$ Definitions and Proofs

### Norms and Distances

Euclidean distance, by default

**Def** The **distance** between $\bar{x} = (x_1, \dots, x_m)$ $\left(\text{or } \bar{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{bmatrix}\right)$ and $\bar{y} = (y_1, \dots, y_m)$, two points in $\mathbb{R}^m$ is

$$\|\bar{x} - \bar{y}\| = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2 + \cdots + (x_m - y_m)^2}.$$

**My notation:** $\bar{x}, \bar{y}, \bar{\alpha}, \bar{\theta}$, etc. are vectors → lower case letters with a bar.

For $m = 1$, $\|x - y\| = \sqrt{(x-y)^2} = |x - y|$ → absolute value of $x - y$

think of it as just the distance between two points in $\mathbb{R}$.

### Triangle Inequality

$$\forall\, \bar{x}, \bar{y} \in \mathbb{R}^m, \quad \|\bar{x} + \bar{y}\| \leq \|\bar{x}\| + \|\bar{y}\|.$$

We could interpret the triangle inequality as saying length of diagonal $\leq$ sum of lengths of sides of the parallelogram.




With $\bar{x} = \bar{u} - \bar{w}$, $\bar{y} = \bar{w} - \bar{v}$, we get

$$\|\bar{u} - \bar{v}\| = \|\bar{u} - \bar{w} + \bar{w} - \bar{v}\| \leq \|\bar{u} - \bar{w}\| + \|\bar{w} - \bar{v}\|$$

for $\bar{u}, \bar{v}, \bar{w} \in \mathbb{R}^m$

Illustration of the above version in 2D: notice the parallelogram here as well!



# Convergence of Sequences

As a first use of distances, we consider convergence of sequences. How do we say a sequence $\{x_n\}$ converges to a real number $a$? We should be able to get arbitrarily close to $a$ by going far enough (large $n$) into the sequence.

**Def 2.1.1** A sequence $\{x_n\}$ of real numbers converges to $a \in \mathbb{R}$ if for every $\epsilon > 0$ (no matter how small), there exists an $N \in \mathbb{N}$ such that $|x_n - a| < \epsilon$ for all $n \geq N$. We write $\lim_{n\to\infty} x_n = a$.

Here is a pictorial representation of the convergence, with the "path" drawn separately below for clarity.




## LSIRA 2.1 Prob 1 (Pg 29)

Show that if $\{x_n\}$ converges to $a$, then the sequence $\{Mx_n\}$ converges to $Ma$. Use the definition of convergence to explain how you choose $N$.

Given $\{x_n\} \to a \implies \forall \epsilon > 0, \exists N \in \mathbb{N}$ such that $|x_n - a| < \epsilon \quad \forall n \geq N$.
$\left(\lim_{n\to\infty} x_n = a\right)$

We want to show $\{Mx_n\} \to Ma$. We want to show that $\forall \epsilon > 0, \exists N' \in \mathbb{N}$ s.t. $|Mx_n - Ma| < \epsilon \quad \forall n \geq N$.

Note that when $M = 0$, the result holds trivially, as $Mx_n = 0 \ \forall n$, and $Ma = 0$. Hence $|Mx_n - Ma| = 0 < \epsilon$ for any $\epsilon > 0$ for $n \geq 1$. $N'$

Also note that both $M > 0$ and $M < 0$ are possible.

Let's assume $M \neq 0$.

First, observe that $|Mx_n - Ma| = |M(x_n - a)| = |M||x_n - a|$.

Note that when $|x_n - a| < \epsilon' = \frac{\epsilon}{|M|}$, $|M||x_n - a| < \epsilon$.

But since $\{x_n\} \to a$, given $\epsilon' = \frac{\epsilon}{|M|} > 0$, $\exists\, N' \in \mathbb{N}$ s.t. $|x_n - a| < \epsilon'$ for all $n \geq N'$. We can choose $N = N'$, and get

$$|x_n - a| < \epsilon' = \frac{\epsilon}{|M|} \quad \forall n \geq N'$$

$$\Rightarrow |M||x_n - a| = \underbrace{|Mx_n - Ma| < \epsilon \quad \forall\, n \geq N'}$$

$\Rightarrow \{Mx_n\}$ converges to $Ma$. $\square$

# MATH 401: Lecture 7 (09/09/2025)

Today: * convergence in $\mathbb{R}^m$
* continuity of functions

We extend the notion of convergence in $\mathbb{R}$ to $\mathbb{R}^m$:




The definition naturally extends to $\mathbb{R}^m$ once we think of $|x_n - a|$ as the distance between $x_n$ and $a$.

**Def 2.1.2** A sequence $\{\bar{x}_n\}$ of points in $\mathbb{R}^m$ converges to $\bar{a} \in \mathbb{R}^m$ if $\forall \epsilon > 0$, $\exists$ an $N \in \mathbb{N}$ such that $\|\bar{x}_n - \bar{a}\| < \epsilon$ $\forall n \geq N$. We write $\lim_{n\to\infty} \bar{x}_n = \bar{a}$.

**LSIRA Prob 2.1.3** $\{\bar{x}_n\}, \{\bar{y}_n\}$ are two sequences in $\mathbb{R}^m$ where $\{x_n\} \xrightarrow{\text{converges}} \bar{a}$, and $\{\bar{y}_n\} \to \bar{b}$. Then show that $\{\bar{x}_n + \bar{y}_n\}$ converges to $\bar{a} + \bar{b}$.

We want to show: $\forall \epsilon > 0$ (any given $\epsilon$), $\exists N \in \mathbb{N}$ such that

$$\|(\bar{x}_n + \bar{y}_n) - (\bar{a}_n + \bar{b}_n)\| < \epsilon \quad \forall n \geq N.$$

same $\epsilon$ as our target

Hint, hint, hint!
$\|\bar{x} + \bar{y} + \bar{z}\| \leq \|\bar{x}\| + \|\bar{y}\| + \|\bar{z}\|$
by applying triangle inequality twice. We often choose $\epsilon/3$ (instead of $\frac{\epsilon}{2}$) with 3 terms!

We are given $\{\bar{x}_n\} \to \bar{a}$, $\{\bar{y}_n\} \to \bar{b}$, so

$\exists N_1 \in \mathbb{N}$ s.t. $\|\bar{x}_n - \bar{a}\| < \frac{\epsilon}{2}$ $\forall n \geq N_1$ and

$\exists N_2 \in \mathbb{N}$ s.t. $\|\bar{y}_n - \bar{b}\| < \frac{\epsilon}{2}$ $\forall n \geq N_2$.

$\Rightarrow$ for $N = \max\{N_1, N_2\}$, we get

$$\begin{aligned}
\|(\bar{x}_n + \bar{y}_n) - (\bar{a} + \bar{b})\| &= \|(\bar{x}_n - \bar{a}) + (\bar{y}_n - \bar{b})\| \\
&\leq \|\bar{x}_n - \bar{a}\| + \|\bar{y}_n - \bar{b}\| \quad \text{by triangle inequality} \\
&< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \quad \text{as } N \geq N_1,\ N \geq N_2.
\end{aligned}$$

$\Rightarrow \{\bar{x}_n + \bar{y}_n\} \to \bar{a} + \bar{b}$. $\square$

# Continuity

$f: \mathbb{R} \to \mathbb{R}$. When is $f$ continuous at $x=a$?

For sequences $\{x_n\} \to a$, we go "far enough out", i.e., $\forall n \geq N \in \mathbb{N}$. Instead of $N \in \mathbb{N}$, here we say $\exists\, \delta > 0$ such that if $|x-a| < \delta$ then $|f(x)-f(a)| < \epsilon$ (for any given $\epsilon > 0$). In other words, $f(x)$ gets close enough to $f(a)$ whenever $x$ is close enough to $a$!

**Def 2.1.4** The function $f: \mathbb{R} \to \mathbb{R}$ is **continuous** at $a \in \mathbb{R}$ if $\forall \epsilon > 0$ (no matter how small), $\exists$ a $\delta > 0$ such that $|f(x)-f(a)| < \epsilon$ whenever $|x-a| < \delta$.

Equivalently, if $|x-a| < \delta$ then $|f(x)-f(a)| < \epsilon$.

We naturally extend the definition to $\mathbb{R}^m$ using distances/norms.

LSIRA uses **F** (bold upper case F)

**Def 2.1.7** The function $\bar{f}: \mathbb{R}^n \to \mathbb{R}^m$ is **continuous** at $\bar{a} \in \mathbb{R}^n$ if $\forall \epsilon > 0$ (no matter how small), $\exists$ a $\delta > 0$ such that $\|\bar{f}(\bar{x}) - \bar{f}(\bar{a})\| < \epsilon$ whenever $\|\bar{x} - \bar{a}\| < \delta$.

By restricting our attention to a subset $A$ of $\mathbb{R}^n$, we naturally extend the above definition to subsets of interest.

**Def 2.1.8** Let $A \subset \mathbb{R}^n$, and $\bar{a} \in A$.

The function $\bar{f}: \mathbb{R}^n \to \mathbb{R}^m$ is **continuous** at $\bar{a} \in A$ if $\forall \epsilon > 0$ (no matter how small), $\exists$ a $\delta > 0$ such that $\|\bar{f}(\bar{x}) - \bar{f}(\bar{a})\| < \epsilon$ whenever $\|\bar{x} - \bar{a}\| < \delta$ and $\bar{x} \in A$.

LSIRA Section 2.1 Prob 4 (extension): If $f_i : \mathbb{R} \to \mathbb{R}$, $i = 1,2,3$ are all continuous at $a \in \mathbb{R}$, the show that so is $f_1 + f_2 - f_3$.
(i.e., show $f_1(x) + f_2(x) - f_3(x)$ is continuous at $x = a$).

Prob 4 considers $f+g$ for two functions $f, g$.

Let $g(x) = f_1(x) + f_2(x) - f_3(x)$. We want to show that

$\forall \epsilon > 0, \exists \delta > 0$ s.t. $|g(x) - g(a)| < \epsilon$ whenever $|x - a| < \delta$.

We know: since $f_i(x)$ are continuous at $x = a$,

$\exists \delta_i > 0$ s.t. $|f_i(x) - f_i(a)| < \frac{\epsilon}{3}$ whenever $|x - a| < \delta_i$, $i = 1,2,3$.

Let $\delta = \min_{i=1,2,3} \{\delta_i\}$. Then

We want $x$ to as close to $a$ as required in each case!

e.g., if $\delta_1 = 0.1$, $\delta_2 = 0.05$ and $\delta_3 = 0.08$, then $\delta \leq 0.05$ works!

$$|g(x) - g(a)| = |(f_1(x) + f_2(x) - f_3(x)) - (f_1(a) + f_2(a) - f_3(a))|$$

$$= |(f_1(x) - f_1(a)) + (f_2(x) - f_2(a)) + (f_3(a) - f_3(x))|$$

$$\leq |f_1(x) - f_1(a)| + |f_2(x) - f_2(a)| + \underbrace{|f_3(a) - f_3(x)|}_{= |f_3(x) - f_3(a)|}$$

by triangle inequality (applied twice)

$$< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} \quad \text{as } \delta \leq \delta_i \text{ for } i = 1,2,3$$

$$= \epsilon \quad \text{whenever } |x - a| < \delta.$$

$\square$

**LSIRA Proposition 2.1.9** Let $g: \mathbb{R} \to \mathbb{R}$ be continuous at $a \in \mathbb{R}$, and $g(a) \neq 0$. Show that $h(x) = \dfrac{1}{g(x)}$ is continuous at $x = a$.

Need to show: $\forall \epsilon > 0$, $\exists \delta > 0$ s.t. $|h(x) - h(a)| < \epsilon$ whenever $|x-a| < \delta$.

We want to show that

$$|h(x) - h(a)| = \left| \frac{1}{g(x)} - \frac{1}{g(a)} \right| < \epsilon.$$

$$\left| \frac{1}{g(x)} - \frac{1}{g(a)} \right| = \left| \frac{g(a) - g(x)}{g(x)\, g(a)} \right| = \frac{|g(x) - g(a)|}{|g(x)|\, \underbrace{|g(a)|}_{\neq 0}}$$

We want to show that $|g(x)|$ is not too small. Else, the fraction could be too large.

There must exist some $\delta' > 0$ such that $|g(x)| > \dfrac{|g(a)|}{2}$ whenever $|x-a| < \delta'$, as $g(a) \neq 0$.




In the picture here, notice that $g(x)$ lies "below" the $\frac{g(a)}{2}$ level, i.e., far enough away from zero, when $|x-a| < \delta'$.

Also, $g(x)$ is continuous at $x = a \Rightarrow$

$\exists\, \delta'' > 0$ s.t. $|g(x) - g(a)| < \epsilon'$ whenever $|x-a| < \delta''$.

Pick $\delta = \min\{\delta', \delta''\}$. Then we get that

$$\left|\frac{1}{g(x)} - \frac{1}{g(a)}\right| = \frac{|g(x)-g(a)|}{|g(x)|\,|g(a)|} < \frac{\epsilon'}{\frac{|g(a)|}{2}\,|g(a)|} = \frac{2\epsilon'}{|g(a)|^2}$$

whenever $|x-a| < \delta$.

If we choose $\epsilon' = \frac{|g(a)|^2}{2}\epsilon$, so that $\frac{2\epsilon'}{|g(a)|^2} = \epsilon$,

we get that $\left|\frac{1}{g(x)} - \frac{1}{g(a)}\right| < \epsilon$ whenever $|x-a| < \delta$.

Hence $\frac{1}{g(x)}$ is continuous at $x = a$

$\square$

In the next section, we consider the setting where the target or candidate limit $(a)$ is not given to us.
Can we still conclude that $\{\bar{x}_n\}$ converges? When?

# MATH 401: Lecture 8 (09/11/2025)

Today: * completeness
* sup, inf, lim sup, lim inf

## Completeness (LSIRA 2.3)

If we don't know the limit target $\bar{a}$, can we still say $\{\bar{a}_n\}$ converges?
If $\{\bar{a}_n\}$ "behaves nicely" and $\bar{a}_n$'s are in a "nice space", then yes!

Here is an intuition for what we mean by "nice space". Suppose $a_n \in A$ where $A$ is a "finite" interval (open or closed). Then we can be sure that the $a_n$'s cannot become arbitrarily large or arbitrarily small.




But in this example, the $a_n$'s belong to a bounded interval $[-1,1]$, but they are not "behaving nicely" as the values oscillate between 1 and $-1$.




We should ideally plot the discrete pts (x), instead of the continuous curve

But if the $a_n$'s are increasing and are bounded from above, or decreasing and bounded from below, we can conclude that $\{a_n\}$ converges!




OR




Finally, even if $a_n$'s are oscillating, and hence not increasing/decreasing, it could still be nice if the oscillations become smaller and smaller — as shown here. →



Intuitively, we want the upper and lower "envelopes" to get closer and closer.

We formalize these intuitive notions of "nice" space and "nice" behavior.

**Def** A nonempty set $A \subset \mathbb{R}$ is **bounded above** if there exists $b \in \mathbb{R}$ such that $a \leq b \ \forall a \in A$, and is **bounded below** if there exists $c \in \mathbb{R}$ such that $a \geq c \ \forall a \in A$. We refer to $b$ as an **upper bound**, and $c$ as a **lower bound**.

If $b$ is an upper bound, then any $b' > b$ is also an upper bound. Similarly, and $c' < c$ is also a lower bound.




We usually want to find a smallest upper bound, and a largest lower bound. This idea is ubiquitous in optimization, where finding the correct maximum value for a function $z = f(\bar{x})$ may be hard, but it may be easier to obtain lower/upper bounds. In order to get as best a handle on the actual max $z$ value, we try to find the smallest upper bound, and the biggest lower bound that work.




In the same way, we want to "estimate" $A$ as accurately as possible by finding the smallest upper bound and the largest lower bound for the set.

# The Completeness Principle

Every nonempty subset $A$ of $\mathbb{R}$ that is bounded above has a least upper bound. This bound is called the **supremum of A**, written $\sup A$.

Similarly every nonempty subset $A$ of $\mathbb{R}$ that is bounded below has a greatest lower bound, called the **infimum of A**, written $\inf A$.

LSIRA 2.2 Problem 1 Argue that $\sup [0,1) = 1$ and $\sup [0,1] = 1$.

Let $A = [0,1) = \{x \in \mathbb{R} \mid 0 \le x < 1\}$. So $x \in A$ can be arbitrarily close to 1, i.e., $x = 1-\epsilon$, $\epsilon > 0$, arbitrarily small. Hence any $1-\epsilon$ cannot be an upper bound for $A$, since $\forall \epsilon > 0$, $\exists\, 1-\epsilon' \in A$ s.t. $1-\epsilon' > 1-\epsilon$.

$\Rightarrow$ $b \geq 1$ satisfies $x \le b$ $\forall x \in A$, and hence $\sup A = 1$.

The same argument holds for $[0,1]$ too. Note that the sup is in $A$ in the latter case, but $\sup A \notin A$ for $A = [0,1)$.

So, what is the big deal about the completeness principle?

First, it does not hold over $\mathbb{Q}$ (rationals), as, e.g.,

$$A = \{x \in \mathbb{R} \mid x^2 < 3\} \text{ has } \sup A = \sqrt{3}. \text{ But}$$

$$B = \{x \in \mathbb{Q} \mid x^2 < 3\} \text{ has } \underline{\text{no supremum}} \text{ in } \mathbb{Q}!$$

$\sqrt{3}$ is irrational, and we can get arbitrarily close to $\sqrt{3}$ using rational numbers!

We say that $\mathbb{Q}$ does not satisfy completeness principle.

# Monotone Sequences, lim sup, lim inf

We now describe sequences that behave "nicely" like the bounded sets introduced earlier. We then consider how to handle sequences that are not as "nice".

"nondecreasing" if you want $a_{n+1} > a_n$ to mean "increasing"

**Def** A sequence $\{a_n\}$ in $\mathbb{R}$ is **increasing** if $a_{n+1} \geq a_n \ \forall n$.

A sequence $\{a_n\}$ in $\mathbb{R}$ is **decreasing** if $a_{n+1} \leq a_n \ \forall n$.

$\{a_n\}$ is **monotone** if it is either increasing or decreasing.

$\{a_n\}$ is **bounded** if $\exists M \in \mathbb{R}$ s.t. $|a_n| \leq M \ \forall n$.

LSIRA Theorem 2.2.2 Every monotone bounded sequence in $\mathbb{R}$ converges to a number in $\mathbb{R}$. → we do not specify which number!

Proof (for increasing sequences). We proceed in two steps.

1. $\{a_n\}$ is bounded $\Rightarrow$ $A = \{a_1, a_2, \dots, a_n, \dots\}$ is bounded. → set

$\Rightarrow \exists\, a \in \mathbb{R}$ such that $\sup A = a$. → using completeness of $\mathbb{R}$

2. $a$ is the least upper bound. → We show $\{a_n\} \to a$

$\Rightarrow a - \epsilon$ is not an upper bound for any $\epsilon > 0$.

$\{a_n\}$ is increasing $\Rightarrow$ $a - \epsilon < a_n \leq a \ \forall n \geq N$ for some $N$.

$\Rightarrow |a - a_n| < \epsilon \ \forall n \geq N$, i.e., $\{a_n\}$ converges.

→ $a_n - a > -\epsilon$ and $a - a_n < \epsilon$

$\square$

But what if $\{a_n\}$ is not monotone and/or not bounded?
Can we still say something about $\{a_n\}$ as $n \to \infty$?

Given a general sequence $\{a_n\}$, we define two related sequences that are monotone themselves.

**Def** Given $\{a_k\}$, $a_k \in \mathbb{R}$, we define two new sequences $\{M_n\}$ and $\{m_n\}$ as follows.

$$M_n = \sup\{a_k \mid k \geq n\} \quad \text{and}$$
$$m_n = \inf\{a_k \mid k \geq n\}.$$

$M_n = \infty$, $m_n = -\infty$ are allowed here.

$M_n$ "captures" how large $\{a_k\}$ can be "after" $n$, and $m_n$ captures how small $\{a_k\}$ can be "after" $n$.

Note that $\{M_n\}$ and $\{m_n\}$ are monotone!

$\{M_n\}$ is decreasing, as suprema are taken over smaller subsets.
and $\{m_n\}$ is increasing, as infima are taken over smaller subsets.

e.g., consider $A = \{1, 2, \ldots, 10\}$. The largest number in $A$ cannot be bigger than the largest number in $A' = \{1, 2, \ldots, 7\}$, or in any $A' \subset A$, in general.

$\Rightarrow \lim_{n\to\infty} M_n$ and $\lim_{n\to\infty} m_n$ exist!

**Def** The **limit superior** or lim sup of the original sequence $\{a_n\}$ is $\limsup_{n\to\infty} a_n = \lim_{n\to\infty} M_n$.

The **limit inferior** of $\{a_n\}$ is $\liminf_{n\to\infty} a_n = \lim_{n\to\infty} m_n$.

We ideally want to draw a sequence of points "....." in place of the continuous curve here




It appears while $\{x_n\}$ may be "oscillating" the upper bounds $M_n$ and lower bounds $m_n$ appear to be converging. Hence, $\{a_n\}$ also appears to converge! But we could have $\{a_n\}$ oscillate forever, even when $M_n$ and $m_n$ are finite $\forall n \geq N$.

<u>LSIRA 2.2 Problem 4</u>

Let $a_n = (-1)^n$. What is $\limsup_{n\to\infty} a_n$? $\liminf_{n\to\infty} a_n = ?$

$$\limsup_{n\to\infty} a_n = \lim_{n\to\infty} M_n = 1.$$

$$\liminf_{n\to\infty} a_n = \lim_{n\to\infty} m_n = -1.$$

Note than $a_n = 1 \ \forall\, n = 2k$, and $a_n = -1 \ \forall\, n = 2l+1$.

Hence $a_n \leq 1 \ \forall\, n$, and $a_n \geq -1 \ \forall\, n$.

In fact, $\{M_n\}$ and $\{m_n\}$ behave identical to $\{a_n\}$ here!

In the above problem, even though lim sup and lim inf are both finite, they are not equal, and we cannot say anything about $\{a_n\}$ converging to a limit. But when the lim sup and lim inf are equal, we get the picture drawn earlier, with $\{a_n\}$ converging to that value!

**LSIRA Proposition 2.2.3** Let $\{a_n\}$ be a sequence of real numbers.

Then $\lim_{n\to\infty} a_n = b$ if and only if

$\limsup_{n\to\infty} a_n = \liminf_{n\to\infty} a_n = b.$ b can $\pm\infty$ here!

($\Leftarrow$) Assume $\limsup_{n\to\infty} a_n = \liminf_{n\to\infty} a_n = b$

$\Rightarrow \lim_{n\to\infty} M_n = \lim_{n\to\infty} m_n = b$




Also, $m_n \le a_n \le M_n \quad \forall n$

$\Rightarrow \lim_{n\to\infty} a_n = b.$ (by "squeeze law" or "squeeze theorem"; LSIRA 2.2 Problem 2 — assigned in HW4!)

We'll finish the proof in the next lecture..

# MATH 401: Lecture 9 (09/16/2025)

Today: * Cauchy sequences
* Intermediate value theorem (IVT)

---

We first present the proof of Proposition 2.2.3...

**LSIRA Proposition 2.2.3** Let $\{a_n\}$ be a sequence of real numbers. Then $\lim_{n\to\infty} a_n = b$ iff $\limsup_{n\to\infty} a_n = \liminf_{n\to\infty} a_n = b$.

$b$ can be $\pm\infty$ here!

($\Leftarrow$) Assume $\limsup_{n\to\infty} a_n = \liminf_{n\to\infty} a_n = b$

$$\Rightarrow \lim_{n\to\infty} M_n = \lim_{n\to\infty} m_n = b$$




Also, $m_n \le a_n \le M_n \quad \forall n$

$$\Rightarrow \lim_{n\to\infty} a_n = b.$$

(by "squeeze law" or "squeeze theorem"; LSIRA 2.2 Problem 2 — assigned in HW4!)

($\Rightarrow$) Assume $\lim_{n\to\infty} a_n = b$, and $b \in \mathbb{R}$.

$\Rightarrow \forall \epsilon > 0, \exists N \in \mathbb{N}$ s.t. $|a_n - b| < \epsilon \quad \forall n \ge N$.

$\Rightarrow a_n - b < \epsilon$ and $b - a_n < \epsilon$

$|x| < 5 \Rightarrow -x < 5$ and $x < 5$

$$\Rightarrow b - \epsilon < a_n < b + \epsilon \quad \forall n \ge N$$

$$\Rightarrow b - \epsilon < m_n < b + \epsilon \quad \text{and}$$
$$b - \epsilon < M_n < b + \epsilon \quad \forall n \ge N$$

We will repeatedly use this trick of splitting $|x-y| < \epsilon$ into $x - y < \epsilon$ and $y - x < \epsilon$

Since the result holds for any $\epsilon > 0$,

$$\limsup_{n\to\infty} a_n = \liminf_{n\to\infty} a_n = b.$$

$\square$

# Cauchy Sequences

We extend the idea of completeness in $\mathbb{R}$ to $\mathbb{R}^m$. But there is no natural way to order points in $\mathbb{R}^m$ (as in $\mathbb{R}$). Instead, we say the points get closer and closer to each other.

**Def 2.2.4** A sequence $\{\bar{x}_n\}$ in $\mathbb{R}^m$ is called a **Cauchy sequence** if $\forall \epsilon > 0, \exists N \in \mathbb{N}$, s.t. $\|\bar{x}_n - \bar{x}_k\| < \epsilon \quad \forall n, k \geq N$.

$n, k$ are two indices, and represent any two points that are both far out enough into the sequence $(n, k \geq N)$




## Completeness Result in $\mathbb{R}^m$

**Theorem 2.2.5** The sequence $\{\bar{x}_n\}$ in $\mathbb{R}^m$ converges **iff** it is cauchy.

This is an iff result. We prove both directions, but one of them is easier than the other. We show the easy direction in $\mathbb{R}^m$, but the reverse direction in $\mathbb{R}$ (and can be extended to $\mathbb{R}^m$).

**Proposition 2.2.6** All convergent sequences in $\mathbb{R}^m$ are cauchy.

Proof Let $\{\bar{a}_n\}$ converge to $\bar{a}$ in $\mathbb{R}^m$.

We want to show $\|\bar{a}_n - \bar{a}_k\| < \epsilon \quad \forall n, k \geq N$ for some $N \in \mathbb{N}$.

$\Rightarrow \forall \epsilon > 0, \exists N \in \mathbb{N}$ s.t. $\|\bar{a}_n - \bar{a}\| < \frac{\epsilon}{2}, \forall n \geq N$.

Ideally, we use $\epsilon'$ here, and then choose $\epsilon' = \frac{\epsilon}{2}$.

$\Rightarrow$ If $n, k \geq N$, then

$$\|\bar{a}_n - \bar{a}_k\| = \|\bar{a}_n - \bar{a} + \bar{a} - \bar{a}_k\| \leq$$

triangle inequality

$$\|\bar{a}_n - \bar{a}\| + \|\bar{a} - \bar{a}_k\| < \underbrace{\frac{\epsilon}{2} + \frac{\epsilon}{2}}_{\text{now we see why we chose } \frac{\epsilon}{2}\,!} = \epsilon \quad \forall n \geq N.$$

$\Rightarrow \{\bar{a}_n\}$ is cauchy. $\square$

We present proof for the reverse direction in $\mathbb{R}$. We can repeat this argument for each dimension to prove the result in $\mathbb{R}^m$. We need a lemma first.

**Lemma 2.2.7** Every Cauchy sequence $\{a_n\}$ in $\mathbb{R}$ is bounded.

Want to show: $|a_n| \leq M$ for some $M \in \mathbb{R}$. note, $M \geq 0$

$\{a_n\}$ is cauchy $\Rightarrow |a_n - a_k| < \epsilon \ \forall n,k \geq N \in \mathbb{N}$ for <u>any</u> $\epsilon > 0$.

$\Rightarrow |a_n - a_N| < 1$ (for $\epsilon = 1$) the definition applies for any $\epsilon$, so we choose $\epsilon = 1$. After all, we just need to find a valid bound

$\Rightarrow a_n - a_N < 1$ and $a_N - a_n < 1$

$\Rightarrow a_n < a_N + 1$ and $a_n > a_N - 1 \ \forall n \geq N$.

$\Rightarrow M = \max\{a_1, a_2, \ldots, a_{N-1}, a_N + 1\}$ is an upper bound, and

$m = \min\{a_1, a_2, \ldots, a_{N-1}, a_N - 1\}$ is a lower bound. $\square$

Could also get $|a_n| - |a_N| \leq |a_n - a_N| < 1$

$\Rightarrow |a_n| \leq |a_N| + 1$.

We could have a larger number among $a_1, a_2, \ldots, a_{N-1}$, which are not considered earlier since the Cauchy definition stipulates $n, k \geq N$.



**Proposition 2.2.8** All Cauchy sequences in $\mathbb{R}$ converge.

**Proof** $\{a_n\}$ is cauchy $\Rightarrow$ $\{a_n\}$ is bounded (by Lemma 2.2.7).

$\Rightarrow M = \limsup_{n\to\infty} a_n$ and $m = \liminf_{n\to\infty} a_n$ are both finite.

We can use Proposition 2.2.3 now, if we can show $M = m$.

$\forall \epsilon > 0, \exists N \in \mathbb{N}$ s.t. $|a_n - a_k| < \epsilon \ \forall n, k \geq N$.

In particular, $|a_n - a_N| < \epsilon \ \forall n \geq N$. (taking $k = N$)

$\Rightarrow a_n - a_N < \epsilon$ and $a_N - a_n < \epsilon \ \forall n \geq N$

$\Rightarrow a_n < a_N + \epsilon$ and $a_n > a_N - \epsilon$

i.e., $a_N - \epsilon < a_n < a_N + \epsilon \ \forall n \geq N$ holds for any $\epsilon > 0$.

$\Rightarrow M_n = \sup\{a_k \mid k \geq n\} < a_N + \epsilon$ ADD

$-(m_n = \inf\{a_k \mid k \geq n\} > a_N - \epsilon) \Rightarrow -m_n < -a_N + \epsilon$

$\Rightarrow M_n - m_n < 2\epsilon \ \forall n \geq N$ and for any $\epsilon > 0$, arbitrary.

$\Rightarrow M = m$ (as $n \to \infty$).

$\square$

We now present four fundamental theorems, the proofs of which use many of the results we have presented. These theorems are quite fundamental in analysis, and also finds use in many applied domains as well.

# Intermediate Value Theorem

This is a rather straightforward result to understand — if a function goes from above the x-axis to below it, and is continuous, then it must cross the x-axis.




**Theorem 2.3.1** (Intermediate Value Theorem) Assume $f:[a,b]\to\mathbb{R}$ is continuous, and $f(a)$ and $f(b)$ have opposite signs. Then there exists $c\in(a,b)$ such that $f(c)=0$.

We will use a characterization of continuity using sequences in the proof (from LSIRA 2.1, actually!).

**Proposition 2.1.5** $f:\mathbb{R}\to\mathbb{R}$ is continuous at $x=a$ iff $\lim_{n\to\infty} f(x_n)=f(a)$ for all sequences $\{x_n\}$ that converge to $a$.

Proof

($\Rightarrow$) Assume $f$ is continuous at $x=a$.

Consider $\{x_n\}\to a$, i.e., $\lim_{n\to\infty} x_n = a$.

Need to show: $\forall \epsilon>0, \exists N\in\mathbb{N}$ s.t. $|f(x_n)-f(a)|<\epsilon$ $\forall n\geq N$.

same

$\Rightarrow$ $\exists \delta>0$ s.t. $|f(x)-f(a)|<\epsilon$ whenever $|x-a|<\delta$.

$\exists N'\in\mathbb{N}$ s.t. $|x_n-a|<\delta$ whenever $n\geq N'$. $\longrightarrow$ plays the "role of $\epsilon$", i.e. the convergence definition must hold for any $\epsilon>0$, and here we choose $\epsilon=\delta$.

$\Rightarrow$ If $n\geq N'$, then $|f(x_n)-f(a)|<\epsilon$, as $|x_n-a|<\delta$.

$\Rightarrow$ $\{f(x_n)\}\to f(a)$. Reverse direction in the next lecture...

# MATH 401: Lecture 10 (09/18/2025)

Today:
* Intermediate Value Theorem (IVT)
* Bolzano-Weierstrass (BW) theorem
* Extreme Value theorem (EVT)

---

Recall: **Proposition 2.1.5** $f: \mathbb{R} \to \mathbb{R}$ is continuous at $x=a$ iff $\lim_{n\to\infty} f(x_n) = f(a)$ for all sequences $\{x_n\}$ that converge to $a$.

($\Leftarrow$) Contrapositive argument

We assume $f$ not continuous at $x=a$, and show there must exist a sequence $\{x_n\}$ that converges to $a$, but $\{f(x_n)\}$ does not converge to $f(a)$.

If $f$ is not continuous at $x=a$, we take the converse of what is implied by it being continuous.

$\Rightarrow \exists\, \epsilon > 0$ s.t. no matter how small you choose $\delta > 0$,

$\exists\, x$ s.t. $|x-a| < \delta$ but $|f(x) - f(a)| \geq \epsilon$.

$x$ here depends on $\delta$

Also, note that $\delta$ can be chosen arbitrarily small.

Pick $\delta = \frac{1}{n}$. $\Rightarrow \exists\, x_n$ s.t. $|x_n - a| < \frac{1}{n}$ but

$|f(x_n) - f(a)| \geq \epsilon$. as $f$ is not continuous at $x=a$.

$\Rightarrow \{x_n\} \to a$ (as $n \to \infty$), but $\{f(x_n)\} \not\to f(a)$. $\square$

This notion of continuity defined in terms of sequences can be quite useful in many contexts, especially when we try to generalize results to higher dimensions.

We now get back to the proof of intermediate value theorem.

Recall:

**Theorem 2.3.1** (Intermediate Value Theorem) Assume $f:[a,b]\to\mathbb{R}$ is continuous, and $f(a)$ and $f(b)$ have opposite signs. Then there exists $c\in(a,b)$ such that $f(c)=0$.

**Proof** Consider $f(a)<0<f(b)$. → The other case of $f(a)>0>f(b)$ can be argued similarly

Let $A=\{x\in[a,b] \mid f(x)<0\}$ and $c=\sup A$. → $A$ is bounded (subset of $[a,b]$), hence $\sup A$ exists.

We show $f(c)=0$.

$f$ is continuous and $f(b)>0$. $\Rightarrow$ $c<b$.

$\Rightarrow$ The sequence $x_n=c+\frac{1}{n}\in[a,b]$ $\forall n\geq N$, for sufficiently large $N$.

$\Rightarrow$ $\{x_n\}\to c$ as $n\to\infty$. → $c<b$,

Also, $f(x_n)>0$ $\forall$ such $n$. → as $x_n\notin A$, since $x_n>c$.

By Proposition 2.1.5, as $f$ is continuous, $\lim_{n\to\infty} f(x_n)=f(c)$, and since $f(x_n)>0$ $\forall n$, $f(c)\geq 0$. → We can be sure that $f(c)\neq 0$.

On the other hand, by definition of $c$, consider $z_n=c-\frac{1}{n}$ for sufficiently large $n$. → When $n$ is large enough, $c-\frac{1}{n}\geq a$.

$\Rightarrow$ $z_n\leq c$ $\forall n$ large enough, and $\{z_n\}\to c$ (as $n\to\infty$).

Also, $z_n\in A\subset[a,b]$ for $n$ large enough. $\Rightarrow$ $f(z_n)<0$.

Again, by Proposition 2.1.5, $f(c)=\lim_{n\to\infty} f(z_n)$ and since $f(z_n)<0$ $\forall n$, we get $f(c)\leq 0$. Hence $f(c)\geq 0$ and $f(c)\leq 0$, i.e., $f(c)=0$. $\square$

Again, we can be sure that $f(c)\neq 0$.

The Intermediate Value Theorem does not hold in $\mathbb{Q}$!

Consider $f(x) = x^2 - 3 \Rightarrow f(0) = -3$ and $f(2) = 1$.

But $\nexists x \in [0,2] \cap \mathbb{Q}$ s.t. $f(x) = 0$, as $\sqrt{3} \notin \mathbb{Q}$.

## The **Bolzano-Weierstrass** (BW) Theorem

We saw that every Cauchy sequence converges. But what if a sequence is not Cauchy, and hence does not converge? Can we still say something nice about its structure? It turns out yes, when the sequence is bounded! We need the notion of a subsequence first.

**Def** (Subsequence) Given a sequence $\{\bar{x}_n\}$ in $\mathbb{R}^m$, we choose an infinite subset of its terms to form another sequence $\{\bar{y}_k\}$. (Of course, it is interesting only when we do not choose all terms of $\{\bar{x}_n\}$).

$$\bar{x}_1, \bar{x}_2, \ldots, \bar{x}_n, \ldots \quad \bar{y}_1 \quad \bar{y}_2 \quad \bar{y}_3 \quad \bar{y}_4 \quad \ldots$$

$\{\bar{y}_k\} \to$ subsequence of $\{\bar{x}_n\}$.

If $n_1 < n_2 < \cdots < n_k < \cdots$ are indices of terms picked to form a new sequence, then

$\{\bar{y}_k\} = \{\bar{x}_{n_k}\}$ is a **subsequence** of $\{\bar{x}_n\}$.

We state and prove BW theorem in $\mathbb{R}$.

**Proposition 2.3.2** Every bounded sequence in $\mathbb{R}$ has a convergent subsequence.

$\{x_n\}$ is bounded $\Rightarrow \exists\ a \le b$ s.t. $x_n \in [a,b] = I_0\ \forall n$.

We identify a Cauchy subseq.

We argue we can pick smaller and smaller subintervals of $I_0$, each of which has infinitely many terms of $\{x_n\}$.




Let $I_1 = [\frac{a+b}{2}, b]$ be such that it has infinitely many terms of $\{x_n\}$.

It could happen that $I_1' = [a, \frac{a+b}{2}]$ is the one with infinitely many terms, or both $I_1$ and $I_1'$ have infinitely many terms of $\{x_n\}$. But since $\{x_n\}$ has infinitely many terms, at least one of the two half intervals is guaranteed to have infinitely many terms. We always choose a half interval with infinitely many terms, and continue the process.

In general, $I_k$ is a half interval of $I_{k-1}$ that has infinitely many terms of $\{x_n\}$. Note that $I_k$ is a subinterval of $I_{k-1}$ for each $k$ ($k \geq 1$), and we get a sequence of nested subintervals that are shrinking in size by a factor of $(\frac{1}{2})$ in each step.

Since $|I_0| = |[a,b]| = b-a$ is finite, $|I_k| \to 0$ as $k \to \infty$.

We can now specify how to construct the convergent subsequence. Essentially, we pick one term of $\{x_n\}$ from each subinterval $I_k$ as follows.

Let $y_1$ be the first element of $\{x_n\}$ in $I_1$. And let $y_2$ be the first element of $\{x_n\}$ **after** $y_1$ that is in $I_2$.

$\vdots$

In general, let $y_k$ be the first element of $\{x_n\}$ **after** $y_{k-1}$ that is in $I_k$, for $k \geq 1$.

Note that the $y_k$'s are included in nested, shorter and shorter subintervals, and hence are getting closer and closer to each other.

we could make this argument more formal

$\Rightarrow$ $\{y_k\}$ is Cauchy!

$\Rightarrow$ $\{y_k\}$ converges by Proposition 2.2.8 $\square$

Consider a somewhat trivial example. Let $x_n = (-1)^n$, $n \in \mathbb{N}$.

One can immediately see that $a_n \leq 1$ $\forall n$ and $a_n \geq -1$ $\forall n$ hold, i.e., we can choose $[a,b] = [-1,1]$ in the proof above.




Then $y_k = (-1)^k$ for $k = 2n$, $n \in \mathbb{N}$ defines a subsequence $\{y_k\} \to 1$, and $z_k = (-1)^k$ for $k = 2n-1$, $n \in \mathbb{N}$ defines a subsequence $\{z_k\} \to -1$.

The BW theorem naturally extends to $\mathbb{R}^m$ — we essentially repeat the above argument one dimension at a time! See LSIRA for details.

We now present two theorems that use the results on sequences to specify properties of "good" (continuous or differentiable) functions defined on the sequences.

## The **Extreme Value Theorem** (EVT) in $\mathbb{R}$

**Theorem 2.3.4** Let $f:[a,b]\to\mathbb{R}$ be a continuous function on the closed bounded interval $[a,b]$. Then $\exists$ points $c, d \in [a,b]$ such that $f(d) \le f(x) \le f(c)$ $\forall\, x \in [a,b]$.

In words, $f$ has maximum and minimum points in $[a,b]$.

Proof (for maximum) → A similar argument can be made for minimum

Let $M = \sup\{f(x) \mid x \in [a,b]\}$. → We're not sure yet whether $M$ is finite

Choose sequence $\{x_n\}$ in $[a,b]$ such that $f(x) \to M$.

As $f$ is continuous, such a sequence exists. → irrespective of whether $M$ is finite or not

$[a,b]$ is bounded $\Rightarrow$ By BW Theorem, $\{x_n\}$ has a convergent subsequence $\{y_k\}$.

$[a,b]$ is closed $\Rightarrow$ $c = \lim_{k\to\infty} y_k \in [a,b]$.

$\Rightarrow f(y_k) \to M$ by construction. → We chose $\{x_n\}$ so that $f(x_n) \to M$ in the first place.

$f$ is continuous $\Rightarrow$ by Proposition 2.1.5, $f(y_k) \to f(c)$.

$\Rightarrow f(c) = M$, i.e., $M$ is the maximum, and $c \in [a,b]$ is the corresponding maximum point for $f$. $\square$

# MATH 401: Lecture 11 (09/23/2025)

Midterm exam: Oct 7

Take-home exam; sections 1.1–1.6, 2.1, 2.2

Today: * Mean value theorem
* Metric spaces

## Mean Value Theorem (MVT) on $\mathbb{R}$

For the final theorem ($4^{th}$ one, after IVT, BW, EVT), we assume the function is much "nicer", i.e., it's differentiable, to be able to present a stronger result on its structure. We recall the definition of derivative first.

Recall: **Derivative** of function $f$ at $x=a$ is

$$f'(a) = \lim_{x \to a} \frac{f(x)-f(a)}{x-a}$$

$f$ is **differentiable** at $x=a$ if the above limit exists.

The mean value theorem says for a differentiable and continuous function $f$ on $[a,b]$, there exists a point inside the interval where the instantaneous slope of the function is equal to the "mean" slope of $f$ over the interval. We need two results to be used as building blocks first.

**Lemma 2.3.5** Let $f:[a,b]\to\mathbb{R}$ have a maximum or minimum at an inner point $c\in(a,b)$ where the function is differentiable. Then $f'(c)=0$.

<u>Proof</u> We show $f'(c)>0$ or $f'(c)<0$ is not possible.

Assume $f'(c)>0$. → A similar argument works for $f'(c)<0$.

$$f'(c) = \lim_{x\to c} \frac{f(x)-f(c)}{x-c} \quad \text{by definition.}$$

$$\Rightarrow \frac{f(x)-f(c)}{x-c} > 0 \quad \text{for all } x \text{ sufficiently close to } c.$$

as $f'(c)>0$

$$\Rightarrow \quad x>c \Rightarrow f(x)>f(c), \text{ and}$$
$$x<c \Rightarrow f(x)<f(c)$$

If $x=c$ is a maximum, then $f(x)\le f(c)$ for $\forall x$.

$\Rightarrow x=c$ is neither a <u>maximum</u> nor minimum. The result follows by the contrapositive argument now. □

**Lemma 2.3.6** (Rolle's Theorem) Let $f:[a,b]\to\mathbb{R}$ be continuous at all $x\in[a,b]$ and is differentiable at all inner points $x\in(a,b)$. If $f(a)=f(b)$, then there exists a point $c\in(a,b)$ such that $f'(c)=0$.

<u>Proof</u> EVT (Theorem 2.3.4) $\Rightarrow$

$f$ has a maximum and minimum in $[a,b]$. Since $f(a)=f(b)$, at least one of these optima must be at an inner point $c$.

So Lemma 2.3.5 $\Rightarrow f'(c)=0$. □




Trivial case:
$f(x)=f(a)\ \forall x\in[a,b]$
↳ straight line!

**Theorem 2.3.7** (The Mean Value Theorem (MVT))

Let $f:[a,b]\to\mathbb{R}$ be continuous in all $[a,b]$ and differentiable at all inner points $x\in(a,b)$. Then there exists $c\in(a,b)$ s.t.

$$f'(c) = \frac{f(b)-f(a)}{b-a}$$

The mean (or average) slope of $f(x)$ over $[a,b]$ is $f(b)-f(a)/(b-a)$. This theorem says there is a point $c\in(a,b)$ where the instantaneous slope, i.e., slope of tangent, is equal to the mean slope!




Proof

Let $g(x) = f(x) - \left(\frac{f(b)-f(a)}{b-a}\right)(x-a)$.

How did we come up with this function?!! See next page...

$g(a) = f(a)$, and

$g(b) = f(b) - \left(\frac{f(b)-f(a)}{\cancel{b-a}}\right)\cancel{(b-a)} = f(a)$.

$b>a$ by assumption, and hence $b-a\neq 0$.

We can show that $g(x)$ is indeed continuous in $[a,b]$ and differentiable at all $x\in(a,b)$.

$g(x) = f(x) + m(x-a)$ for constant $m$; $f(x)$ is continuous and differentiable, and so is $(x-a)$; their sum is so as well.

So, Rolle's theorem (Lemma 2.3.6) $\Rightarrow \exists\, c\in(a,b)$ s.t. $g'(c)=0$.

$$\Rightarrow g'(x) = f'(x) - \left(\frac{f(b)-f(a)}{b-a}\right) = 0 \text{ at } x=c.$$

$$\Rightarrow f'(c) = \frac{f(b)-f(a)}{b-a}.$$

$\square$

Now, how did we come up with the $g(x)$ function?!

want to show!

$$f'(c) = \lim_{x \to c} \frac{f(x)-f(c)}{x-c} = \frac{f(b)-f(a)}{b-a}$$

$$f'(x)\Big|_{x=c} =$$

$$\Longrightarrow \quad \underbrace{f'(x) - \left(\frac{f(b)-f(a)}{b-a}\right)}\Bigg|_{x=c} = 0$$

Looks like $g'(x)=0$ for some function $g(x)$.

We want to find $g(x)$ such that $g(a)=g(b)=0$, and then we could use Rolle's theorem!

So we take antiderivative of $f'(x) - \dfrac{f(b)-f(a)}{b-a}$ to get

$$f(x) - \left(\frac{f(b)-f(a)}{b-a}\right)x + C. \longrightarrow \text{constant}$$

With $g(x) = f(x) - \left(\dfrac{f(b)-f(a)}{b-a}\right)x + C$, we choose $C$ such that $g(a)=g(b)$! Note that

$$\underbrace{f(b) - \left(\frac{f(b)-f(a)}{b-a}\right)b + \left(\frac{f(b)-f(a)}{b-a}\right)a}_{g(b)} = f(a) = \underbrace{f(a) - \left(\frac{f(b)-f(a)}{b-a}\right)a + \left(\frac{f(b)-f(a)}{b-a}\right)a}_{g(a)}.$$

# Chapter 3 Metric Spaces

We have showed several results on sequences and functions in $\mathbb{R}$ and $\mathbb{R}^m$. But many of these results could be shown for far more general spaces which have many of the nice properties of $\mathbb{R}$ (or $\mathbb{R}^m$). We define metric spaces with this goal in mind.

## 3.1 Definitions

nonempty

**Def** A **metric space** $(X, d)$ consists of a set $X \neq \emptyset$, and a function $d: X \times X \longrightarrow [0, \infty)$ such that

(i) (positivity) $d(x,y) \geq 0 \ \forall \ x, y \in X$, and
$d(x,y) = 0$ iff $x = y$;

(ii) (symmetry) $d(x,y) = d(y,x) \ \forall x, y \in X$; and

(iii) (triangle inequality) $d(x,y) \leq d(x,z) + d(z,y) \ \forall x, y, z \in X$.

hold. A function $d$ satisfying (i)–(iii) on $X$ is a **metric** on $X$.

We sometimes write just $X$, when the metric $d$ is evident. At the same time, note that a space $X$ could have multiple metrics defined on it. The first example we consider studies a metric on $\mathbb{R}^2$ that is different from the usual Euclidean metric.

## Examples

**LSIRA eg. 3** Manhattan or taxicab metric (in $\mathbb{R}^2$). $X$

For $\bar{x}, \bar{y} \in \mathbb{R}^2$, let $\bar{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$, $\bar{y} = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}$, $\bar{z} = \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}$

$$d(\bar{x}, \bar{y}) = |y_1 - x_1| + |y_2 - x_2|.$$

Check that this is a metric space.




The intuition is that if a taxi is to go from point $x$ to point $y$ in downtown Manhattan with perpendicular streets, it will have to go East/West (horizontal) and then North/South (vertical). We add these two straight line distances to get the taxicab distance between $x$ and $y$.

(i) $d(\bar{x}, \bar{y}) \geq 0$ holds, as $|x_1 - y_1| \geq 0$ and $|x_2 - y_2| \geq 0$.
The only way we get $d(\bar{x}, \bar{y}) = 0$ is when both absolute differences are zero, i.e., when $x_1 = y_1$ and $x_2 = y_2$, i.e., when $\bar{x} = \bar{y}$.

(ii) $d(\bar{x}, \bar{y}) = d(\bar{y}, \bar{x})$ follows from absolute differences being symmetric, i.e., $|x_i - y_i| = |y_i - x_i|$ for $i = 1, 2$.

(iii) Triangle inequality:

$$\begin{aligned} d(\bar{x}, \bar{y}) &= |y_1 - x_1| + |y_2 - x_2| \\ &= |y_1 - z_1 + z_1 - x_1| + |y_2 - z_2 + z_2 - x_2| \\ &\leq |y_1 - z_1| + |z_1 - x_1| + |y_2 - z_2| + |z_2 - x_2| \\ &= d(x, z) + d(z, y). \end{aligned}$$

standard triangle inequality in $\mathbb{R}$

Hence $d(\bar{x}, \bar{y}) \leq d(\bar{x}, \bar{z}) + d(\bar{z}, \bar{y}) \ \forall \bar{x}, \bar{y}, \bar{z} \in \mathbb{R}^2$. $\square$

# MATH 401: Lecture 12 (09/25/2025)

Today: * Examples of metric spaces
* isometry

---

**Recall** $(X,d)$: Metric space (i) positivity: $d(x,y) \geq 0 \ \forall x,y \in X$, and $d(x,y)=0 \iff x=y$.

(ii) symmetry: $d(x,y)=d(y,x) \ \forall x,y \in X$

(iii) triangle inequality: $d(x,y) \leq d(x,z)+d(z,y) \ \forall x,y,z \in X$

One more requirement: $d(x,y) < \infty \ \forall x,y \in X$. (finiteness)

The finiteness requirement is usually satisfied. But you should use your judgement to decide in which cases this property needs to be proved.

---

## LSIRA 3.1 Example 4 (Problem 1)

Let $X$ be the space of messages, where each message is a vector ($k$ is fixed)

$$\bar{x} = \begin{bmatrix} x_1 \\ \vdots \\ x_k \end{bmatrix}, \text{ with } x_i \in \underset{\text{alphabet}}{S} = \{\underbrace{s_1, \dots, s_N}_{\text{symbols in the alphabet}}\}.$$

Let $d(\bar{x},\bar{y}) = \#$ indices $i$ where $x_i \neq y_i$. Show $(X,d)$ is a metric space.

Note that shorter messages could be padded up with empty cells, or a "dummy" symbol (equivalent to 0 when using numbers). And larger messages can be chopped up into pieces of length $k$ each (with padding up if needed for the last piece).

An example: Let $S = \{R, G, B\}$ for colors red, green, blue. Consider vectorizing each image by assigning to each pixel its predominant color, and stacking these color symbols into a vector. For instance, a $12 \times 12$ pixel image is represented by a 144-vector of color values from $\{R, G, B\}$.

(i) $d(\bar{x},\bar{y}) \geq 0$, and $d(\bar{x},\bar{y})=0 \iff \bar{x}=\bar{y}$ as messages.

↳ $d(\bar{x},\bar{y})$ is the # places (or indices) where the messages differ, and hence is $\geq 0$.

$d(\bar{x},\bar{y})=0 \iff \bar{x}$ and $\bar{y}$ are identical in all entries, i.e., they do not differ at all. Hence $\bar{x}=\bar{y}$.

(ii) symmetry ✓

$$\begin{aligned} d(\bar{x},\bar{y}) &= \#\text{ indices where } x_i \neq y_i \\ &= \#\text{ indices where } y_i \neq x_i \\ &= d(\bar{y},\bar{x}) \end{aligned}$$

(iii) triangle inequality.

$d(\bar{x},\bar{y})$ counts # indices $i$ where $x_i \neq y_i$

$x_i \neq y_i \implies$ cannot have $x_i = z_i$ and $z_i = y_i$.

Combining with $z_i$, here are the possibilities:

1. $x_i \neq z_i$, $z_i = y_i$
2. $x_i = z_i$, $z_i \neq y_i$
3. $x_i \neq z_i$, $z_i \neq y_i$




$\implies d(x,y) \leq d(x,z) + d(z,y)$, as there are three possible cases for each index contributing 1 to the right-hand side sum corresponding to the one case possibly contributing 1 to the left-hand side distance.

□

# LSIRA Pg47, Problem 2 **Distance between Functions.**

Let $X$ = set of all continuous functions from $[a,b] \to \mathbb{R}$, and let

$$d_1(f,g) = \sup\{|f(x)-g(x)| : x \in [a,b]\}.$$

Show that $d_1$ is a metric over $X$.







Measuring distance between functions is a widely studied problem in analysis as well as applications. We illustrate use of $d_1$ here (top right). But $d_1$ may not be the best choice in all cases — in the second instance, $f$ and $g$ are quite close to each other except at one point $x=c$, where $g$ shoots up a lot. $d_1(f,g)$ will be quite large here, even though $f$ and $g$ are near equal. Measuring distance using the area between $f$ and $g$ may be better here.

<u>Proof</u>: We first show $d_1(f,g)$ is <u>finite</u> for any $f, g \in X$.

finitness is not obvious in this case!

$f, g$ are continuous over $[a,b]$

$\Rightarrow f-g$ is continuous over $[a,b]$.

By the Extreme Value Theorem (Theorem 2.3.4), $h = f-g$ has a maximum and minimum over $[a,b]$.

$\Rightarrow \sup\{|h(x)| : x \in [a,b]\}$ is finite.

$\Rightarrow d_1(f,g)$ is finite.

(i) (positivity) $d_1(f,g) = \sup\{|f(x)-g(x)| : x \in [a,b]\} \geq 0$, as the supremum of a set of $\geq 0$ values is $\geq 0$.

Need to also show $d_1(f,g) = 0 \iff f = g$ over $[a,b]$

$(\Rightarrow)$ $f(x) = g(x)\ \forall x \in [a,b]$

$\Rightarrow |f(x)-g(x)| = 0\ \forall x \in [a,b]$

$\Rightarrow \sup\{|f(x)-g(x)| : x \in [a,b]\} = 0 \Rightarrow d_1(f,g) = 0.$

$(\Leftarrow)$ $d_1(f,g) = 0 \Rightarrow \sup\{|f(x)-g(x)| : x \in [a,b]\} = 0$

The supremum of a set of $\geq 0$ is zero $\Rightarrow$ each element $= 0$!

$\Rightarrow f(x) = g(x)\ \forall x \in [a,b].$

(ii) (symmetry) $|f(x)-g(x)| = |g(x)-f(x)|$

$\Rightarrow d(f,g) = \sup\{|f(x)-g(x)| : x \in [a,b]\} =$
$\sup\{|g(x)-h(x)| : x \in [a,b]\} = d_1(g,f).$

(iii) <u>Triangle inequality</u>

$|f(x)-g(x)| = |f(x)-h(x)+h(x)-g(x)|$
$\leq |f(x)-h(x)| + |h(x)-g(x)|$ by standard triangle inequality over $\mathbb{R}$.

$$\begin{aligned} d_1(f,g) &= \sup\{|f(x)-g(x)| : x \in [a,b]\} \\ &\leq \sup\{|f(x)-h(x)| + |h(x)-g(x)| : x \in [a,b]\} \\ &\leq \sup\{|f(x)-h(x)| : x \in [a,b]\} + \\ &\qquad \sup\{|h(x)-g(x)| : x \in [a,b]\} \\ &= d_1(f,h) + d_1(h,g) \qquad \forall f,g,h \in X. \end{aligned}$$

as $\sup\{a+b\} \leq \sup\{a\} + \sup\{b\}$

$\square$

# LSIRA Pg48, Problem 7

Let $(X,d)$ be a metric space, and $x_i \in X$, $i=1,\dots,n$.

Show $d(x_1,x_n) \leq \sum_{i=1}^{n-1} d(x_i,x_{i+1})$.

$d(x_1,x_2)+d(x_2,x_3)+\dots+d(x_{n-1},x_n)$




## Can use induction

$n=2$ (base case)

$d(x_1,x_2) \leq d(x_1,x_2)$ holds, as both sides are the same.

$n=3$ case could be considered as the base case as well:

$d(x_1,x_3) \leq d(x_1,x_2)+d(x_2,x_3)$ follows from triangle inequality for $(X,d)$.

Assume result is true for $n=k$, i.e.,

$$d(x_1,x_k) \leq \sum_{i=1}^{k-1} d(x_i,x_{i+1}) \qquad (*)$$

For $n=k+1$,

$$d(x_1,x_{k+1}) \leq d(x_1,x_k)+d(x_k,x_{k+1}) \quad \text{by triangle inequality in } (X,d)$$

$$\leq \sum_{i=1}^{k-1} d(x_i,x_{i+1}) + d(x_k,x_{k+1}) \quad \text{by } (*)$$

$$= \sum_{i=1}^{k} d(x_i,x_{i+1}) \quad \checkmark$$

Hence the result holds for all $n$.

$\square$

We now talk about comparing two metric spaces, and functions between metric spaces. When are two metric spaces "the same"? As metric spaces are about pairwise distances between points, we want these distances to be preserved.

**Def 3.1.2** Let $(X, d_X)$ and $(Y, d_Y)$ are metric spaces.

An **isometry** between the spaces is a bijection $i: X \to Y$ such that $d_X(x,y) = d_Y(i(x), i(y))$ $\forall x, y \in X$.

The two spaces are isometric if an isometry exists between them.

Since $i$ is a bijection, its inverse exists, and $i^{-1}$ is an isometry from $(Y, d_Y)$ to $(X, d_X)$. Hence we can just say isometry between the spaces.

LSIRA Pg48, Problem 11 For $a \in \mathbb{R}$, let $f(x) = x + a$. Show $f$ is an isometry from $\mathbb{R}$ to $\mathbb{R}$.

$$d(x,y) = |x-y| \quad \forall x, y \in \mathbb{R}.$$

To show $f$ is an isometry, we need to show

1. $f$ is a bijection; and
2. $d(f(x), f(y)) = d(x,y)$ $\forall x, y \in \mathbb{R}$

1. $f(x) = x + a$ is a bijection as $\underbrace{x_1 \neq x_2 \Rightarrow f(x_1) = x_1 + a \neq x_2 + a = f(x_2)}_{\text{injection}}$; and $\underbrace{\forall y \in \mathbb{R}, \exists x = f^{-1}(y) = y - a}_{\text{surjection}}$.

2. $d(f(x), f(y)) = |f(x) - f(y)| = |x + a - (y + a)| = |x - y|$
$= d(x,y)$ $\forall x, y \in \mathbb{R}$.

$\square$

The requirement of $i$ being a bijection is too strict in some settings. There may be spaces that look otherwise quite similar, even if they are not isometric.

If $i: X \to Y$ in Definition 3.1.2 is only an injection (and not a bijection), we call $i$ an <u>embedding</u>.

You may have heard about embeddings in other geometric settings. For instance, think of a sphere (surface of a ball) in 3D space. We can work with the sphere as a metric space — the distance between any two points is the length of the shortest curve connecting the points that lies entirely on the surface (of the sphere). This is called the shortest geodesic distance. We can prove it is a metric.

$S^2$ (2-sphere) surface of a ball




in $\mathbb{R}^3$

There are many curves between $x$ and $y$ that lie on the surface of the sphere. Length of a shortest geodesic curve defines the distance between $x$ and $y$.

The embedding here is literally the positioning of the sphere in $\mathbb{R}^3$.