

MATH 230 - Lecture 3 (01/18/2011)

3-1

Echelon form and reduced echelon form

Row reduction is the procedure of using EROs to reduce **any** matrix into echelon form, and then to reduced echelon form.

Def:

pivot \rightarrow the leading entry in a nonzero row

pivot position \rightarrow position of a leading entry (same as pivot)

pivot column \rightarrow a column that contains a pivot.

Prob 4, pg 25 Row reduce the given matrix to echelon form, and then to reduced echelon form. Circle the pivots, list the pivot columns.

$$\begin{bmatrix} \textcircled{1} & 3 & 5 & 7 \\ 3 & 5 & 7 & 9 \\ 5 & 7 & 9 & 1 \end{bmatrix} \xrightarrow[R_3 - 5R_1]{R_2 - 3R_1} \begin{bmatrix} 1 & 3 & 5 & 7 \\ 0 & \textcircled{-4} & -8 & -12 \\ 0 & -8 & -16 & -34 \end{bmatrix} \xrightarrow{R_3 - 2R_2} \begin{bmatrix} \textcircled{1} & 3 & 5 & 7 \\ 0 & \textcircled{-4} & -8 & -12 \\ 0 & 0 & 0 & \textcircled{-10} \end{bmatrix} \begin{array}{l} \text{is in} \\ \text{echelon form} \end{array}$$

$$\xrightarrow[R_3 \times (-\frac{1}{10})]{R_2 \times (-\frac{1}{4})} \begin{bmatrix} 1 & 3 & 5 & 7 \\ 0 & \textcircled{1} & 2 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \xrightarrow{R_1 - 3R_2} \begin{bmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & \textcircled{1} \end{bmatrix} \xrightarrow[R_2 - 2R_3]{R_1 + 2R_3} \begin{bmatrix} \textcircled{1} & 0 & -1 & 0 \\ 0 & \textcircled{1} & 2 & 0 \\ 0 & 0 & 0 & \textcircled{1} \end{bmatrix}$$

pivot columns

Columns 1, 2, and 4 are pivot columns.

is in
reduced echelon form

Solution of linear systems: Use row reduction to convert the augmented matrix to echelon form, and if found consistent, reduce further to reduced echelon form to write the solutions down.

Prob 12, pg 25 The augmented matrix of a system of linear equations is given. Find all solutions.

pivots are circled

$$\begin{array}{c}
 x_1 \quad x_2 \quad x_3 \quad x_4 \quad \text{rhs} \\
 \left[\begin{array}{cccc|c}
 \textcircled{1} & -7 & 0 & 6 & 5 \\
 0 & 0 & 1 & -2 & -3 \\
 -1 & 7 & 4 & 2 & 7
 \end{array} \right] \xrightarrow{R_3 + R_1} \left[\begin{array}{cccc|c}
 1 & -7 & 0 & 6 & 5 \\
 0 & 0 & \textcircled{1} & -2 & -3 \\
 0 & 0 & -4 & 8 & 12
 \end{array} \right] \xrightarrow{R_3 + 4R_2} \left[\begin{array}{cccc|c}
 \textcircled{1} & -7 & 0 & 6 & 5 \\
 0 & 0 & \textcircled{1} & -2 & -3 \\
 0 & 0 & 0 & 0 & 0
 \end{array} \right]
 \end{array}$$

Pivot columns 1 & 3 correspond to variables x_1 and x_3 .

Def. Variables corresponding to pivot columns are called **basic** variables. Variables that are not basic are **free** variables (or non-basic variables).

Here, x_1, x_3 are basic and x_2, x_4 are free variables.

We can choose the values of free variables arbitrarily. The basic variables can be written in terms of the free variables.

A system of linear equations has infinitely many solutions if and only if it has free variables

Back to Prob 12, pg 25

The reduced echelon form is $\left[\begin{array}{cccc|c} 1 & -7 & 0 & 6 & 5 \\ 0 & 0 & 1 & -2 & -3 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$.

$$\left. \begin{array}{l} x_1 - 7x_2 + 6x_4 = 5 \\ x_3 - 2x_4 = -3 \end{array} \right\}$$

$$\begin{array}{l} x_1 = 5 + 7x_2 - 6x_4 \\ x_3 = -3 + 2x_4 \\ x_2, x_4 \text{ are free} \end{array}$$

↪ general solution to the original system.

Summary of linear systems

1. If the echelon form of the augmented matrix has a row $[0 \ 0 \ \dots \ 0 \ | \ *]$ for $*$ non-zero, the system is inconsistent.
Else
2. If there are no free variables, the system has a unique solution.
3. If there are free variables, the system has infinitely many solutions.

Prob 15(a) pg 25

$$\begin{bmatrix} \blacksquare & * & * & * \\ 0 & \blacksquare & * & * \\ 0 & 0 & \blacksquare & 0 \end{bmatrix}$$

is the augmented matrix in echelon form. Is the system consistent?

YES, as there is no row of the form $[0\ 0\ 0\ *]$ for $*$ non-zero.

Is the solution unique?

YES. As there are no free variables.

Prob 25, pg 26 The coefficient matrix has a pivot in each row. Why is the system consistent?

There is at least one non-zero entry in each row of the coefficient matrix. Hence we cannot have an inconsistent row $([0\ 0\ \dots\ 0\ |\ \blacksquare])$ in the augmented matrix.

Does this system have a unique solution?

Don't know

There could be free variables even when there is a pivot in each row, in which case, the solution is not unique.

Prob 19, pg 26

$$x_1 + hx_2 = 2$$

$$4x_1 + 8x_2 = k$$

Choose h, k such that system is

(a) inconsistent;

(b) has unique solution; and

(c) has many solutions.

Also, in cases (b) and (c), write all solutions down.

$$\left[\begin{array}{cc|c} 1 & h & 2 \\ 4 & 8 & k \end{array} \right] \xrightarrow{R_2 - 4R_1} \left[\begin{array}{cc|c} 1 & h & 2 \\ 0 & 8-4h & k-8 \end{array} \right]$$

(a) We need $8-4h=0$ and $k-8 \neq 0$, so that we get $[0 \ 0 \ | \ \square]$. So, $h=2$ and $k \neq 8$.

can also write as $k \in \mathbb{R}/\{8\}$

set of all reals
without 8.

(b) $8-4h \neq 0$ so that there are no free variables.

$h \neq 2$, k is any real number. $h \in \mathbb{R}/\{2\}$, $k \in \mathbb{R}$.

(c) Need $8-4h=0$ and $k-8=0$ so that x_2 is a free variable in a consistent system.

i.e., $h=2, k=8$.

Now, let's find the solution(s) in (b) and (c).

(3.6)

(b) We have $8-4h \neq 0$.

$$\left[\begin{array}{cc|c} 1 & h & 2 \\ 0 & 8-4h & k-8 \end{array} \right] \xrightarrow{R_2 \times \frac{1}{(8-4h)}} \left[\begin{array}{cc|c} 1 & h & 2 \\ 0 & 1 & \frac{k-8}{8-4h} \end{array} \right] \xrightarrow{R_1 - hR_2}$$

$$\left[\begin{array}{cc|c} 1 & 0 & 2 - h\left(\frac{k-8}{8-4h}\right) \\ 0 & 1 & \frac{k-8}{8-4h} \end{array} \right]$$

$$x_1 = 2 - h\left(\frac{k-8}{8-4h}\right) = \frac{2(8-4h) - hk + 8h}{4(2-h)} \\ = \frac{16 - hk}{4(2-h)}$$

$$x_2 = \frac{k-8}{8-4h} = \frac{k-8}{4(2-h)}$$

The unique solution is $x_1 = \frac{16-hk}{4(2-h)}$, $x_2 = \frac{k-8}{4(2-h)}$.

(c) $h=2, k=8$ gives

$$\left[\begin{array}{cc|c} 1 & 2 & 2 \\ 0 & 0 & 0 \end{array} \right]$$

x_2 is free, x_1 is basic

$$\left. \begin{array}{l} x_1 + 2x_2 = 2 \\ x_2 \text{ free} \end{array} \right\} \Rightarrow$$

$$\left\{ \begin{array}{l} x_1 = 2 - 2x_2 \\ x_2 \text{ free} \end{array} \right\}$$

general solution.


Vector Equations (Section 1.3)

3-7

$$\begin{aligned} 3x_1 + x_2 &= 7 \\ x_1 + 2x_2 &= 4 \end{aligned} \quad \left[\begin{array}{cc|c} 3 & 1 & 7 \\ 1 & 2 & 4 \end{array} \right] \xrightarrow{R_1 - 3R_2} \left[\begin{array}{cc|c} 0 & -5 & -5 \\ 1 & 2 & 4 \end{array} \right] \xrightarrow{\begin{array}{l} R_1 \times (-\frac{1}{5}) \\ R_1 \leftrightarrow R_2 \end{array}}$$

$$\left[\begin{array}{cc|c} 1 & 2 & 4 \\ 0 & 1 & 1 \end{array} \right] \xrightarrow{R_1 - 2R_2} \left[\begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & 1 \end{array} \right] \quad \left\{ \begin{array}{l} x_1 = 2 \\ x_2 = 1 \end{array} \right\} \text{ unique solution.}$$

This is the "row picture" of the system. We'll now look at the "column picture".



$$\left[\begin{array}{cc|c} 3 & 1 & 7 \\ 1 & 2 & 4 \end{array} \right] \text{ corresponds to}$$

$$\begin{bmatrix} 3 \\ 1 \end{bmatrix} x_1 + \begin{bmatrix} 1 \\ 2 \end{bmatrix} x_2 = \begin{bmatrix} 7 \\ 4 \end{bmatrix} \quad \left. \vphantom{\begin{bmatrix} 3 \\ 1 \end{bmatrix}} \right\} \text{ is a vector equation}$$

vectors.

A vector (or a column vector, by default) is a column of entries. An n -vector is a column with n entries.

"bar" indicates a vector $\bar{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$. \bar{x} here is also a $1 \times n$ matrix.