




# MATH 524 – Lecture 17 (10/17/2023)

Today: \* barycentric subdivision  
\* simplicial approximation

Recall: barycentric subdivision



Explicit Description of the simplices in  $Sd K$


Notation  $\sigma_1 > \sigma_2$  means  $\sigma_2$  is a proper face of  $\sigma_1$ , or equivalently,  $\sigma_1$  is a proper coface of  $\sigma_2$ .

Lemma 15.3 [M]  $Sd K$  is the collection of simplices of the form  $\hat{\sigma}_1 \hat{\sigma}_2 \dots \hat{\sigma}_p$  where  $\sigma_1 > \sigma_2 > \dots > \sigma_p$ .

Illustration

The edges in  $Sd K$  are of the form  $\hat{e}_j \hat{v}_i$  where  $e_j > v_i$ ; or of the form  $\hat{f}_0 \hat{e}_j$  where  $f_0 > e_j$ . Similarly, the triangles

in  $Sd K$  are of the form  $\hat{f}_0 \hat{e}_j \hat{v}_i$  where  $f_0 > e_j > v_i$ .




Proof (by induction)

True for  $K^{(0)}$  (as  $\hat{v} = v \ \forall v \in K^{(0)}$ ).

Now suppose each simplex of  $Sd K$  lying in  $|K^{(p)}|$  is of this form. Let  $\tau$  be a simplex of  $Sd K$  lying in  $|K^{(p+1)}|$ , but not in  $|K^{(p)}|$ . Then  $\tau$  belongs to one of the complexes  $\hat{\sigma} * L_{\sigma}$ , where  $\sigma$  is a  $(p+1)$ -simplex of  $K$ , and  $L_{\sigma}$  is the first barycentric subdivision of the complex made of the proper faces of  $\sigma$ . By induction, each simplex of  $L_{\sigma}$  is of the form  $\hat{\sigma}_0 \hat{\sigma}_1 \dots \hat{\sigma}_p$ . Then  $\tau$  must be of the form  $\hat{\sigma} \hat{\sigma}_0 \dots \hat{\sigma}_p$ . □

Notice that the simplices in  $Sd^2 K$  are much "smaller" than the simplices in  $Sd K$ . This observation is formalized in the following theorem.



**Theorem 15.4 [M]** Given a finite complex  $K$ , a metric for  $|K|$ , and  $\varepsilon > 0$ , there exists an  $r$  such that each simplex in  $Sd^r K$  has diameter less than  $\varepsilon$ .

**Def** For a subset  $S$  of a metric space  $(X, d)$ , its **diameter** is  $\text{diam}(S) = \sup \{ d(\bar{x}, \bar{y}) \mid \bar{x}, \bar{y} \in S \}$ .

See [M] for the proof. ↪ the metric of the metric space.

# Simplicial Approximation

We now talk about how to use subdivision to find a simplicial approximation of any continuous function  $h: |K| \rightarrow |L|$ .

Recall: A simplicial approximation of a continuous map  $h: |K| \rightarrow |L|$  by a simplicial map  $f: K \rightarrow L$  satisfies

$$h(\text{St } v) \subset \text{St}(f(v)) \quad \forall v \in K^{(0)}.$$

We had also seen that homomorphisms  $f_{\#}$  associated with simplicial maps  $f: K \rightarrow L$  induce isomorphisms at the homology level. Our ultimate goal is to argue that the homology groups are determined by the underlying spaces, rather than specific choices of the complex.


We now look at the next step toward that goal - to show that we can always approximate a continuous map by a simplicial map once we have a fine enough subdivision of the original complex.

The result for the case when  $K$  is finite is quite accessible compared to that when  $K$  is infinite. We will discuss only the finite case in detail here.

Theorem 16.1 [M] (The finite simplicial approximation theorem)

Let  $K$  and  $L$  be complexes, and let  $K$  be finite. Given a continuous map  $h: |K| \rightarrow |L|$ , there is an  $r$  such that  $h$  has a simplicial approximation  $f: Sd^r K \rightarrow L$ .

Here is an illustration we already saw in Lecture 15.



Of course,  $K'$  is not a barycentric subdivision of  $K$  here. But the example illustrates the result nonetheless. The key idea is that by subdividing  $K$  enough, we could approximate  $h$  by a simplicial map from  $K'$  to  $L$ .

Even in the illustration shown, one could argue that  $f$  misses the detail in  $h$  in some places, e.g., between  $i_L$  and  $j_L$  where  $h$  looks like " $\subseteq$ ", while  $f$  looks like " $\bullet$ ".

But one could consider a finer subdivision  $K''$  of  $K'$  where we have some more vertices between  $i$  and  $j$ . The image under  $f$  could be closer to  $h$  in that case.

In this context, the barycentric subdivision is just one kind of subdivision we could use. At the same time, its nice structure makes it convenient to devise proofs of results. On the other hand,  $Sd^r K$  might produce "bad" simplices, e.g., triangles that are too skinny. There are other classes of subdivision where the triangles produced are "round" (while still holding a small diameter).

Proof Cover  $|K|$  by open sets  $h^{-1}(St \bar{w})$ , as  $\bar{w}$  ranges over  $L^{(0)}$ . Let this covering be called  $\mathcal{A}$ . Then  $\mathcal{A}$  is an open covering of the compact metric space  $|K|$ . So, there exists a number  $\lambda$  such that any set of  $|K|$  with diameter less than  $\lambda$  lies in one of the elements of  $\mathcal{A}$ . This number is called a **Lebesgue number** for  $\mathcal{A}$ .

Here is the standard argument for why a Lebesgue number should exist in this case.

Suppose there does not exist a Lebesgue number for  $\mathcal{A}$ . Then we can choose a sequence  $C_n$  of sets such that  $\text{diam}(C_n) < \frac{1}{n}$ , but  $C_n$  does not lie in any element of  $\mathcal{A}$ . Choose  $\bar{x}_n \in C_n$ . By compactness, some subsequence  $\{\bar{x}_{n_i}\}$  converges, to say,  $\bar{x}$ . Now,  $\bar{x} \in A$  for some  $A \in \mathcal{A}$ . As  $A$  is open, it contains  $C_{n_i}$  for  $i$  sufficiently large — a contradiction.

Back to the main proof now...

Choose  $r$  s.t. each simplex  $\sigma$  in  $Sd^r K$  has  $\text{diam}(\sigma) < \frac{1}{2}$ .  
 Then each  $st \bar{\sigma}$  for  $\bar{\sigma} \in (Sd^r K)^{(0)}$  has diameter  $< \lambda$ . So,  
 it  $(st \bar{\sigma})$  lies in one of the sets  $h^{-1}(st \bar{\omega})$ . So,  
 $h: |K| \rightarrow |L|$  satisfies the star condition relative to  
 $Sd^r K$  and  $L$ , and hence a simplicial approximation exists.  $\square$


Extending the simplicial approximation theorem to the case  
 when  $K$  is not finite ( $h: |K| \rightarrow |L|$ ) is much more involved.

We introduce a key technique related to subdivision  
 used in this process. In particular, the default barycentric  
 subdivision will not work.


Subdividing  $K$  while keeping  $K_0$  (a subcomplex) fixed

**Def** Here is a sequence of subdivisions of skeletons of  $K$ .  
 Let  $J_0 = K^{(0)}$ . In general,  $J_p$  is a subdivision of  $K^{(p)}$ ,  
 and each simplex  $\tau$  of  $K_0$  with  $\dim(\tau) \leq p$  belongs  
 to  $J_p$ . Define  $J_{p+1}$  to be the union of  $J_p$ , all  
 $\tau \in K_0$  with  $\dim(\tau) = p+1$ , and the cones  $\hat{\sigma} * J_\tau$  as  
 $\sigma$  ranges over all  $(p+1)$ -simplices of  $K$  **not in**  $K_0$ . Here  
 $J_\tau$  is a subcomplex of  $J_p$  whose polytope is  $Bd \sigma$ .  
 The union of all complexes  $J_p$   $\forall p$  is a subdivision  
 of  $K$ , denoted  $Sd(K/K_0)$ , and is called the  
 first barycentric subdivision of  $K$ , holding  $K_0$  fixed.

We define  $Sd^r(K/K_0)$  similarly:  $Sd^2(K/K_0) = Sd(Sd(K/K_0)/K_0)$ , for instance.



$K_0: \{\text{edges } e_0, e_1, e_3, \text{ and all } v_j\}.$



$Sd(K/K_0)$  and  $Sd^2(K/K_0)$

We finish by listing the main result. See [M] for proof.

### Theorem 16.5 [M] (The general simplicial approximation theorem)

Let  $K$  and  $L$  be complexes, and let  $h: |K| \rightarrow |L|$  be a continuous map. Then there exists a subdivision  $K'$  of  $K$  such that  $h$  has a simplicial approximation  $f: K' \rightarrow L$ .

That's all we will cover in this subtopic. Next we move on to an important algebraic technique — exact sequences.