

MATH 273 - Lecture 27 (12/04/2014)

27.1

Line integrals, continued... (Section 15.1)


Q5) Evaluate $\int_C (x + \sqrt{y}) ds$ where C is given in the picture.

$$C_1: y = x^2 \text{ for } 0 \leq x \leq 1$$

$$\vec{r}(t) = \frac{t}{x} \hat{i} + \frac{t^2}{y} \hat{j}, \quad 0 \leq t \leq 1$$

$$C_2: y = x \text{ for } 1 \leq x \leq 0$$

$$\vec{r}(t) = \frac{(1-t)}{x} \hat{i} + \frac{(1-t)}{y} \hat{j}, \quad 0 \leq t \leq 1$$



$$C_1: \vec{v}(t) = \hat{i} + 2t \hat{j} \Rightarrow |\vec{v}(t)| = \sqrt{1^2 + (2t)^2} = \sqrt{1+4t^2}$$

$$f(x(t), y(t)) = x + \sqrt{y} = t + \sqrt{t^2} = 2t$$

$$\int_C f(x,y) ds = \int_0^1 f(t) |\vec{v}(t)| dt = \int_0^1 2t \sqrt{1+4t^2} dt = \frac{1}{4} \cdot \frac{2}{3} (1+4t^2)^{3/2} \Big|_0^1$$

$$= \frac{1}{6} (5\sqrt{5} - 1).$$

$$C_2: \vec{v}(t) = -\hat{i} - \hat{j} \Rightarrow |\vec{v}(t)| = \sqrt{(-1)^2 + (-1)^2} = \sqrt{2}$$

$$\int_{C_2} f ds = \int_0^1 \left(\frac{(1-t)}{x} + \sqrt{\frac{(1-t)}{y}} \right) \sqrt{2} dt = \int_1^0 (u + \sqrt{u}) \sqrt{2} du$$

swap limits to
cancel -ve
sign

$$u = 1 - t$$
$$du = -dt$$

$$= \int_0^1 (u + \sqrt{u}) \sqrt{2} du$$

$$= \sqrt{2} \left(\frac{1}{2} u^2 + \frac{2}{3} u^{3/2} \right) \Big|_0^1 = \sqrt{2} \left(\frac{1}{2} + \frac{2}{3} \right) = \frac{7\sqrt{2}}{6}.$$

$$\text{Hence } \int_C f ds = \int_{C_1} f ds + \int_{C_2} f ds = \frac{(5\sqrt{5}-1)}{6} + \frac{7\sqrt{2}}{6} = \frac{5\sqrt{5}+7\sqrt{2}-1}{6}.$$

Line Integrals over Vector Fields

We now extend the idea of line integrals to the settings of a vector field. A gravitational field, or electromagnetic field are typical examples — we will study how to compute the work done in moving an object along a curve in such a field.


Def A **vector field** is a function that assigns a vector to each point in its domain.

$$\vec{F}(x,y,z) = M(x,y,z)\hat{i} + N(x,y,z)\hat{j} + P(x,y,z)\hat{k}$$

\vec{F} is continuous (differentiable) if $M, N,$ and P are continuous (differentiable).

Examples

1. fluid flow through a bottleneck



2. Gradient field:

$$\vec{F} = \nabla f$$

$$\vec{F}(x,y,z) = \frac{\partial f}{\partial x}\hat{i} + \frac{\partial f}{\partial y}\hat{j} + \frac{\partial f}{\partial z}\hat{k}.$$

The vector assigned to each point is directed along the direction of largest rate of change of f , and its magnitude is the directional derivative along this direction.

Line Integral of a Vector Field

Curve C : $\vec{r}(t)$, $a \leq t \leq b$.

Unit tangent vector: $\hat{T} = \frac{\vec{v}(t)}{|\vec{v}(t)|} = \frac{d\vec{r}}{ds}$

book uses \mathbf{T}
(bold T)

Def

The line integral of \vec{F} along C is


$$\int_C \vec{F} \cdot \hat{T} ds = \int_C \left(\vec{F} \cdot \frac{d\vec{r}}{ds} \right) ds = \int_C \vec{F} \cdot d\vec{r}$$

To evaluate, we compute $\int_a^b \left(\vec{F}(\vec{r}(t)) \cdot \left(\frac{d\vec{r}}{dt} \right) \right) dt$.

7. $\vec{F} = 3y \hat{i} + 2x \hat{j} + 4z \hat{k}$

(b) $C_2: \vec{r}(t) = \underbrace{t}_x \hat{i} + \underbrace{t^2}_y \hat{j} + \underbrace{t^4}_z \hat{k}, 0 \leq t \leq 1$.

Find line integral of vector field \vec{F} along C_2 from $(0,0,0)$ to $(1,1,1)$.



$$\vec{F}(t) = 3 \underbrace{t^2}_y \hat{i} + 2 \underbrace{t}_x \hat{j} + 4 \underbrace{t^4}_z \hat{k}$$

$$\frac{d\vec{r}}{dt} = 1 \hat{i} + 2t \hat{j} + 4t^3 \hat{k}$$

$$\int_{C_2} \vec{F} \cdot \left(\frac{d\vec{r}}{dt} \right) dt = \int_0^1 (3t^2 + 2t \cdot 2t + 4t^4 \cdot 4t^3) dt = \int_0^1 (7t^2 + 16t^7) dt$$

$$= \left. \frac{7}{3} t^3 + \frac{16}{8} t^8 \right|_0^1 = \frac{7}{3} + 2 = \frac{13}{3}.$$

This line integral can be used to evaluate the work done in moving an object along curve C from $\underbrace{t=a}_{\text{pt. A}}$ to $\underbrace{t=b}_{\text{pt. B}}$ over a force field \vec{F} .

$$W = \int_C \vec{F} \cdot \hat{T} ds = \int_a^b (\vec{F}(\vec{r}(t)) \cdot \frac{d\vec{r}}{dt}) dt.$$

(19). $\vec{F} = xy\hat{i} + y\hat{j} - yz\hat{k}$, $\vec{r}(t) = \underbrace{t}_x\hat{i} + \underbrace{t^2}_y\hat{j} + \underbrace{t}_z\hat{k}$, $0 \leq t \leq 1$.
Find the work done in moving along C in the direction of increasing t (from $t=0$ to $t=1$).

$$\vec{F}(\vec{r}(t)) = t t^2 \hat{i} + t^2 \hat{j} - t^2 t \hat{k} = t^3 \hat{i} + t^2 \hat{j} - t^3 \hat{k}$$

$$\frac{d\vec{r}}{dt} = \hat{i} + 2t\hat{j} + \hat{k}. \quad \vec{F} \cdot \left(\frac{d\vec{r}}{dt}\right) = t^3 + 2t^3 - t^3 = 2t^3$$

$$W = \int_0^1 \vec{F} \cdot \left(\frac{d\vec{r}}{dt}\right) dt = \int_0^1 2t^3 dt = \left. \frac{2}{4} t^4 \right|_0^1 = \frac{1}{2}.$$