

STABLE COMPARISON OF TIME SERIES USING TOPOLOGY

Bala Krishnamoorthy Elizabeth Thompson
Washington State University

arXiv: 2501.02817

MADEPS

COMPARING TIME SERIES

→ stock price v/s volume of trading



(image:www)


COMPARING TIME SERIES

→ stock price v/s volume of trading



(image:www)

→ rainfall amount v/s river levels



(image:www)


COMPARING TIME SERIES

→ stock price v/s volume of trading



(image:www)


→ rainfall amount v/s river levels



(image:www)

? How to compare?


CROSS-RECURRENCE (%DET)



(Mallot & Leonardi, 2018)

Cross-recurrence
matrix \mathcal{C}


CROSS-RECURRENCE (%DET)



(Vallot & Leonardi, 2018)

Cross-recurrence
matrix \mathcal{C}


CROSS-RECURRENCE (%DET)



(Vallot & Leonardi, 2018)

Cross-recurrence
matrix \mathcal{C}

CROSS-RECURRENTENCE (%DET)




(Mallot & Leonardi, 2018)

Cross-recurrence
matrix \mathcal{C}

→ %Determinism (%DET)

— % of 1s in diagonal strips of \mathcal{C}

CROSS-RECURRENT (%DET)



(Vallot & Leonardi, 2018)


Cross-recurrence
matrix \mathcal{C}

→ % Determinism (%DET)

— % of 1s in diagonal strips of \mathcal{C}

→ four parameters: time lag τ , embed. dimension, dist. threshold (tol), # diagonal strips (minDI)



%DET: INSTABILITY






%DET = 95

→ minDL = 7

%DET = 83



%DET: INSTABILITY



%DET = 95

→ minDL = ~~7~~5

%DET = ~~88~~97.1

STABLE MEASURE?

- ? Define a measure of similarity that is probably stable?
- small change in input \Rightarrow small change in measure

STABLE MEASURE?

- ? Define a measure of similarity that is probably stable?
 - small change in input \Rightarrow small change in measure
- ? Minimize # parameters?

STABLE MEASURE?

- ? Define a measure of similarity that is probably stable?
 - small change in input \Rightarrow small change in measure
- ? Minimize # parameters?
- ? Effects of large dimensions?

STABLE MEASURE?

- ? Define a measure of similarity that is probably stable?
 - small change in input \Rightarrow small change in measure
- ? Minimize # parameters?
- ? Effects of large dimensions?
- ? Does it work well in practice?

RESULTS

- ✓ $\text{score}(f_1 | f_2)$: conditional periodicity of f_1 given f_2
 - persistent homology of a single point cloud

RESULTS

- ✓ $\text{Score}(f_1 | f_2)$: conditional periodicity of f_1 given f_2
 - persistent homology of a single point cloud
- ✓ theoretical stability under
 - small changes to periodicity
 - Gaussian noise

RESULTS

- ✓ $\text{score}(f_1|f_2)$: conditional periodicity of f_1 given f_2
 - persistent homology of a single point cloud
- ✓ theoretical stability under
 - small changes to periodicity
 - Gaussian noise
- ✓ single parameter: embedding dimension
 - min embedding dimension to control precision of $\text{score}(f_1|f_2)$


RESULTS

- ✓ $\text{score}(f_1|f_2)$: conditional periodicity of f_1 given f_2
 - persistent homology of a single point cloud
- ✓ theoretical stability under
 - small changes to periodicity
 - Gaussian noise
- ✓ single parameter: embedding dimension
 - min embedding dimension to control precision of $\text{score}(f_1|f_2)$
- ✓ stable under PCA truncation

RESULTS




- ✓ $\text{score}(f_1|f_2)$: conditional periodicity of f_1 given f_2
 - persistent homology of a single point cloud
- ✓ theoretical stability under
 - small changes to periodicity
 - Gaussian noise
- ✓ single parameter: embedding dimension
 - min embedding dimension to control precision of $\text{score}(f_1|f_2)$
- ✓ stable under PCA truncation
- ✓ computational evidence

SCORE($f_1|f_2$) vs %DEIT






VICTORIS-RIPS(VR) PERSISTENCE

Edelsbrunner et al., 2002




VICTORIS-RIPS(VR) PERSISTENCE

Edelsbrunner et al., 2002



Persistence diagram (PD):



PERSISTENCE STABILITY

Chazal, de Silva, Oudot (2014)

Chazal, de Silva, Glisse, Oudot (2016)

$$d_B(d_{gm}(VR(X)), d_{gm}(VR(Y))) \leq 2d_{GH}(X, Y) \leq 2d_H(X, Y)$$

\nwarrow \swarrow
Gromov-Hausdorff Hausdorff

PERSISTENCE STABILITY

Chazal, de Silva, Oudot (2014)

Chazal, de Silva, Glisse, Oudot (2016)

$$d_B(d_{gm}(VR(X)), d_{gm}(VR(Y))) \leq 2d_{GH}(X, Y) \leq 2d_H(X, Y)$$

\nwarrow
Gromov-Hausdorff

\nwarrow
Hausdorff

X, Y in

common

metric

space

\nwarrow
under all isometric
embeddings of X, Y
into common metric
space

PERSISTENCE STABILITY

Chazal, de Silva, Oudot (2014)

Chazal, de Silva, Glisse, Oudot (2016)

$$d_B(d_{gm}(VR(X)), d_{gm}(VR(Y))) \leq 2d_{GH}(X, Y) \leq 2d_H(X, Y)$$

\nwarrow
Gromov-Hausdorff

\nwarrow
Hausdorff

X, Y in

common

metric


space

\nwarrow
under all isometric
embeddings of X, Y
into common metric
space

PH: VR Persistent Homology


PH ON TIME SERIES

Perea & Harter (2015)
Perea et al. (2015)




PH ON TIME SERIES

Perea & Harter (2015)
Perea et al. (2015)





PH ON TIME SERIES

Perea & Harter (2015)
Perea et al. (2015)



PH ON TIME SERIES



Perea & Harer (2015)
Perea et al. (2015)



$mp(dgm(x))$
max 1-persistence

PH ON TIME SERIES




Perea & Harer (2015)
Perea et al. (2015)




$mp(dgm(x))$
max 1-persistence

$$\rightarrow \text{score}(f) = \frac{mp(dgm(x))}{\sqrt{3}}$$

PH ON TIME SERIES



PH ON TIME SERIES



→ stability: closeness of $SW(f)$ and $SW(S_N f)$

N -truncated Fourier series of f

CONDITIONAL PERIODICITY SCORE

Def $f_1, f_2 : [0, 2\pi] \rightarrow \mathbb{R}$ continuous, periodic time series
 f_2 more-periodic than f_1 $\left(\frac{2\pi}{\omega_2} \leq \frac{2\pi}{\omega_1} \right)$

CONDITIONAL PERIODICITY SCORE

Def $f_1, f_2 : [0, 2\pi] \rightarrow \mathbb{R}$ continuous, periodic time series
 f_2 more-periodic than f_1 $\left(\frac{2\pi}{\omega_2} \leq \frac{2\pi}{\omega_1} \right)$

Conditional SW Embedding of f_1 given f_2 :

$$SW_{M,\tau} f_{1|2}(t) = (f_1(t), \dots, f_1(t + M\tau))^T \text{ for } \tau = \frac{2\pi}{\omega_2(m+1)}.$$

CONDITIONAL PERIODICITY SCORE

Def $f_1, f_2 : [0, 2\pi] \rightarrow \mathbb{R}$ continuous, periodic time series f_2 more-periodic than f_1 $\left(\frac{2\pi}{\omega_2} \leq \frac{2\pi}{\omega_1} \right)$

Conditional SW Embedding of f_1 given f_2 :

$$SW_{M,\tau} f_{1|2}(t) = (f_1(t), \dots, f_1(t + M\tau))^T \text{ for } \tau = \frac{2\pi}{\omega_2(M+1)}.$$

$$\text{score}(f_1|f_2) = \frac{\text{mp}(\text{dgm}(SW f_{1|2}(\tau)))}{\sqrt{3}}, \tau \in [0, \frac{2\pi}{\omega_1}].$$


CONDITIONAL PERIODICITY SCORE

Def $f_1, f_2 : [0, 2\pi] \rightarrow \mathbb{R}$ continuous, periodic time series f_2 more-periodic than f_1 $\left(\frac{2\pi}{\omega_2} \leq \frac{2\pi}{\omega_1} \right)$

Conditional SW Embedding of f_1 given f_2 :

$$SW_{M,\tau} f_{1|2}(t) = (f_1(t), \dots, f_1(t + M\tau))^T \text{ for } \tau = \frac{2\pi}{\omega_2(M+1)}$$

$$\text{score}(f_1|f_2) = \frac{\text{mp}(\text{dgm}(SW f_{1|2}(\tau)))}{\sqrt{3}}, \tau \in [0, \frac{2\pi}{\omega_1}]$$



REDUCTION TO PERIODICITY SCORE



Proposition 1

$$\lim_{\frac{2\pi}{w_2} \rightarrow \frac{2\pi}{w_1}} \text{score}(f_1, f_2) = \text{score}(f_1)$$

REDUCTION TO PERIODICITY SCORE

Proposition 1


$$\lim_{\frac{2\pi}{w_2} \rightarrow \frac{2\pi}{w_1}} \text{score}(f_1, f_2) = \text{score}(f_1)$$



REDUCTION TO PERIODICITY SCORE

Proposition 1


$$\lim_{\frac{2\pi}{w_2} \rightarrow \frac{2\pi}{w_1}} \text{score}(f_1, f_2) = \text{score}(f_1)$$



REDUCTION TO PERIODICITY SCORE

Proposition 1

$$\lim_{\frac{2\pi}{\omega_2} \rightarrow \frac{2\pi}{\omega_1}} \text{score}(f_1, f_2) = \text{score}(f_1)$$



STABILITY RESULTS

✓ small change in periodicity of $f_2 \Rightarrow$
 $\text{score}(f, l_2)$ changes only a little

STABILITY RESULTS

✓ small change in periodicity of $f_2 \Rightarrow$
score(f_1, f_2) changes only a little

Theorem 2 $f_1, f_{21}, f_{22} : [0, 2\pi] \rightarrow \mathbb{R} : \frac{2\pi}{w_1} \geq \frac{2\pi}{w_{21}} \geq \frac{2\pi}{w_{22}}$
 $X_1 = SW_{M, \tau_1} f_{1|21}(T), X_2 = SW_{M, \tau_2} f_{1|22}(T)$

$$d_H(X_1, X_2) \leq \sqrt{M+1} \left| \frac{2\pi}{w_{21}} - \frac{2\pi}{w_{22}} \right| \sqrt{\sum_i |f'_i(c_i)|^2}$$

STABILITY RESULTS

✓ small change in periodicity of $f_2 \Rightarrow$
 $\text{score}(f_1, f_2)$ changes only a little

Theorem 2 $f_1, f_{21}, f_{22} : [0, 2\pi] \rightarrow \mathbb{R} : \frac{2\pi}{w_1} \geq \frac{2\pi}{w_{21}} \geq \frac{2\pi}{w_{22}}$
 $X_1 = SW_{M, \tau_1} f_{1|21}(T), X_2 = SW_{M, \tau_2} f_{1|22}(T)$

$$d_H(X_1, X_2) \leq \sqrt{M+1} \left| \frac{2\pi}{w_{21}} - \frac{2\pi}{w_{22}} \right| \sqrt{\sum_i |f'_i(c_i)|^2}$$

$$|\text{score}(f_1, f_{21}) - \text{score}(f_1, f_{22})| \leq 4 \sqrt{\frac{M+1}{3}} \left| \frac{2\pi}{w_{21}} - \frac{2\pi}{w_{22}} \right| \sqrt{\sum_i |f'_i(c_i)|^2}$$

STABILITY RESULTS

- ✓ Small Gaussian noise added to $f_1 \Rightarrow$
 $\text{score}(f_1, f_2)$ changes only a little w.h.p

STABILITY RESULTS

✓ Small Gaussian noise added to $f_1 \Rightarrow$

$\text{score}(f_1, f_2)$ changes only a little w.h.p

Lemma 3 $f_1^\sigma(t) = f_1(t) + \epsilon_t$ for $\epsilon_t \sim \mathcal{N}(0, \sigma^2)$

For $\delta \in (0, 1)$, it is at least $(1 - \delta) \cdot 100\%$ likely that

$$|\text{score}(f_1, f_2) - \text{score}_\sigma(f_1, f_2)| \leq 4\sigma \sqrt{\frac{M+1}{3\delta}}$$

STABILITY RESULTS

- ✓ Using top K principal components of PCA projection of $SW f_{1/2}(t)$ preserves score (f_1, f_2)

STABILITY RESULTS

✓ Using top K principal components of PCA projection of $SW f_{1|2}(t)$ preserves $\text{score}(f_1, f_2)$

Theorem 4 $\phi: \mathbb{R}^{M+1} \rightarrow \mathbb{R}^K$: PCA projection with eigenvectors/values $\{\zeta_k, \lambda_k\}_{k=1}^N$.

$$|\text{score}(f_1, f_2) - \text{score}_\phi(f_1, f_2)| \leq \sqrt{\frac{8}{3}} \sqrt{\sum_{i=k+1}^N \lambda_i^2}$$

"unused" \nwarrow $(N-K)$ eigenvalues

MIN. EMBEDDING DIMENSION

- ✓ Embedding dim. N above which $\text{Score}(f_1|f_2)$ does not change much with dimension

MIN. EMBEDDING DIMENSION

✓ Embedding dim. \mathcal{N} above which $\text{Score}(f_1|f_2)$ does not change much with dimension

Theorem 5 For $\epsilon > 0$, with $\mathcal{N} = \left\lceil \frac{2\pi}{\omega_2 \epsilon} \right\rceil$, for any $M_2 > M_1 \geq \mathcal{N}$


$$|\text{score}_{M_1}(f_1|f_2) - \text{score}_{M_2}(f_1|f_2)| \leq \epsilon \cdot g(M_1, f_1) + h(M_1, M_2, f_1)$$

constants w.r.t. ϵ

COMPUTATIONAL RESULTS

Score($f_1|f_2$)

$w_2: 15 \rightarrow 3$ ($w_1=3$)




\mathcal{U} for $\epsilon=0.1, 0.05, 0.03$


COMPUTATIONAL RESULTS

Score($f_1|f_2$) under Gaussian noise


5%



10%







15%



$w_2: 15 \rightarrow 3$ ($w_1=3$)

COMPUTATIONAL RESULTS

%DET under 10% Gaussian noise



$W_2: 15 \rightarrow 3$ ($w_i=3$)

OPEN QUESTIONS

? Testing on real data?

OPEN QUESTIONS

? Testing on real data?

? (Bound on) min # PCs needed to "preserve"
 $\text{score}_{\phi}(f_1|f_2)$?

OPEN QUESTIONS

? Testing on real data?

? (Bound on) min # PCs needed to "preserve"
 $\text{score}_{\phi}(f_1|f_2)$?

Thank you!