

Class 8: Breast Cancer Mini Project

Michael McClellan (PID: A169692395)

Table of contents

Background	1
Data Import	1
Clustering	3
Principle Component Analysis	4
The importance of data scaling	4
PCA of wisc.data	9
5. Combining methods	14
Clustering on PCA results	14
7. Prediction	17

Background

This source provides materials for a class mini-project focused on unsupervised learning analysis of human breast cancer cell data. Students will conduct principal component analysis (PCA) for dimensionality reduction and then apply hierarchical and k-means clustering techniques. The project involves exploratory data analysis, interpreting PCA results, evaluating clustering performance by comparing cluster assignments to actual diagnoses, and optionally combining PCA with clustering. The goal is to identify potential groupings within the cell data based on their characteristics without prior knowledge of malignancy, and the project concludes with an application of the PCA model to classify new patient samples.

Data Import

Our data comes from the University of Wisconsin Medical Center

```
wisc.df <- read.csv("WisconsinCancer.csv", row.names=1)
```

Q1. How many patients/samples are in this dataset?

```
nrow(wisc.df)
```

```
[1] 569
```

Q2. How many of the observations have a malignant diagnosis?

```
table(wisc.df$diagnosis)
```

```
  B   M  
357 212
```

```
sum(wisc.df$diagnosis == "M")
```

```
[1] 212
```

Q3. How many variables/features in the data are suffixed with mean?

```
colnames(wisc.df)
```

```
[1] "diagnosis"           "radius_mean"  
[3] "texture_mean"        "perimeter_mean"  
[5] "area_mean"           "smoothness_mean"  
[7] "compactness_mean"    "concavity_mean"  
[9] "concave.points_mean" "symmetry_mean"  
[11] "fractal_dimension_mean" "radius_se"  
[13] "texture_se"          "perimeter_se"  
[15] "area_se"             "smoothness_se"  
[17] "compactness_se"      "concavity_se"  
[19] "concave.points_se"   "symmetry_se"  
[21] "fractal_dimension_se" "radius_worst"  
[23] "texture_worst"       "perimeter_worst"  
[25] "area_worst"          "smoothness_worst"  
[27] "compactness_worst"   "concavity_worst"  
[29] "concave.points_worst" "symmetry_worst"  
[31] "fractal_dimension_worst"
```

```
length(grep("mean", colnames(wisc.df), value = T))
```

```
[1] 10
```

There is a diagnosis column that is the clinician consensus that I want to exclude from any further analysis. We will come back later and compare our results to this diagnosis.

```
diagnosis <- as.factor(wisc.df$diagnosis)
head(diagnosis)
```

```
[1] M M M M M M
Levels: B M
```


Now we can remove it from the wisc.df

```
wisc.data <- wisc.df[,-1]
```

Clustering

Let's try a hclust()

```
hc <- hclust(dist(wisc.data))
plot(hc)
```



We can extract clusters from this rather poor dendrogram/tree with `cutree()`

```
grps <- cutree(hc, k=2)
```

How many individuals in each cluster?

```
table(grps)
```

```
grps
  1  2
549 20
```

```
table(diagnosis)
```

```
diagnosis
  B  M
357 212
```

We can generate a cross-table that compares our cluster `grps` vector with our `diagnosis` vector values.

```
table(diagnosis, grps)
```

```
      grps
diagnosis  1  2
  B 357   0
  M 192  20
```

Principle Component Analysis

The importance of data scaling


The main function for PCA in base R is `prcomp()` it has a default input parameter of `scale=FALSE`.

```
# prcomp()
head(mtcars)
```

	mpg	cyl	disp	hp	drat	wt	qsec	vs	am	gear	carb
Mazda RX4	21.0	6	160	110	3.90	2.620	16.46	0	1	4	4
Mazda RX4 Wag	21.0	6	160	110	3.90	2.875	17.02	0	1	4	4
Datsun 710	22.8	4	108	93	3.85	2.320	18.61	1	1	4	1
Hornet 4 Drive	21.4	6	258	110	3.08	3.215	19.44	1	0	3	1
Hornet Sportabout	18.7	8	360	175	3.15	3.440	17.02	0	0	3	2
Valiant	18.1	6	225	105	2.76	3.460	20.22	1	0	3	1

We could do a PCA of this data as is and it could be mis-leading...

```
pc <- prcomp(mtcars)
biplot(pc)
```



Let's look at the mean values of each column and their standard deviation.

```
colMeans(mtcars)
```

mpg	cyl	disp	hp	drat	wt	qsec
20.090625	6.187500	230.721875	146.687500	3.596563	3.217250	17.848750
vs	am	gear	carb			
0.437500	0.406250	3.687500	2.812500			

```
apply(mtcars, 2, sd)
```

mpg	cyl	disp	hp	drat	wt
6.0269481	1.7859216	123.9386938	68.5628685	0.5346787	0.9784574
qsec	vs	am	gear	carb	
1.7869432	0.5040161	0.4989909	0.7378041	1.6152000	

We can “scale” this data before PCA to get a much better representation and analysis of all the columns.

```
mtscale <- scale(mtcars)
```

```
colMeans(mtscale)
```

mpg	cyl	disp	hp	drat
6.678685e-16	-6.938894e-18	-2.949030e-16	-2.428613e-17	-1.113692e-15
wt	qsec	vs	am	gear
5.221518e-16	-1.465841e-15	1.387779e-17	8.326673e-17	-5.030698e-17
carb				
1.387779e-17				

```
apply(mtscale, 2, sd)
```

mpg	cyl	disp	hp	drat	wt	qsec	vs	am	gear	carb
1	1	1	1	1	1	1	1	1	1	1


```
pc.scale <- prcomp(mtscale)
```

We can look at the two main analysis results figures from PCA - the “PC plot” (aka score plot, ordination plot, or PC1 vs. PC2 plot). The “loadings plot” how the original variables contribute to the new PCs.

A loadings plot of the unscaled PCA results


```
library(ggplot2)
```

```
ggplot(pc$rotation) +  
  aes(PC1, rownames(pc$rotation)) +  
  geom_col()
```



Loadings plot of the scaled data.

```
ggplot(pc.scale$rotation) +  
  aes(PC1, rownames(pc$rotation)) +  
  geom_col()
```




PC plot of scaled PCA results

```
library(ggrepel)

ggplot(pc.scale$x) +
  aes(PC1, PC2, label=rownames(pc.scale$x)) +
  geom_point() +
  geom_text_repel()
```

Warning: ggrepel: 9 unlabeled data points (too many overlaps). Consider increasing max.overlaps



Key point: In general we will set `scale=TRUE` when we do PCA. This is not the default but probably should be...

We can check the SD and mean of the different columns in `wisc.data` to see if we need to scale - hint: we do!

PCA of wisc.data

```
wisc.pr <- prcomp(wisc.data, scale=TRUE)
```

To see how well PCA is doing here in terms capturing the variance(or spread) in the data we can use the `summary()` function.

```
summary(wisc.pr)
```

Importance of components:


	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	3.6444	2.3857	1.67867	1.40735	1.28403	1.09880	0.82172
Proportion of Variance	0.4427	0.1897	0.09393	0.06602	0.05496	0.04025	0.02251
Cumulative Proportion	0.4427	0.6324	0.72636	0.79239	0.84734	0.88759	0.91010

	PC8	PC9	PC10	PC11	PC12	PC13	PC14
Standard deviation	0.7811	0.6811	0.5811	0.4811	0.3811	0.2811	0.1811
Proportion of Variance	0.0111	0.0111	0.0111	0.0111	0.0111	0.0111	0.0111
Cumulative Proportion	0.9212	0.9323	0.9434	0.9545	0.9656	0.9767	0.9878

Standard deviation	0.69037	0.6457	0.59219	0.5421	0.51104	0.49128	0.39624
Proportion of Variance	0.01589	0.0139	0.01169	0.0098	0.00871	0.00805	0.00523
Cumulative Proportion	0.92598	0.9399	0.95157	0.9614	0.97007	0.97812	0.98335
	PC15	PC16	PC17	PC18	PC19	PC20	PC21
Standard deviation	0.30681	0.28260	0.24372	0.22939	0.22244	0.17652	0.1731
Proportion of Variance	0.00314	0.00266	0.00198	0.00175	0.00165	0.00104	0.0010
Cumulative Proportion	0.98649	0.98915	0.99113	0.99288	0.99453	0.99557	0.9966
	PC22	PC23	PC24	PC25	PC26	PC27	PC28
Standard deviation	0.16565	0.15602	0.1344	0.12442	0.09043	0.08307	0.03987
Proportion of Variance	0.00091	0.00081	0.0006	0.00052	0.00027	0.00023	0.00005
Cumulative Proportion	0.99749	0.99830	0.9989	0.99942	0.99969	0.99992	0.99997
	PC29	PC30					
Standard deviation	0.02736	0.01153					
Proportion of Variance	0.00002	0.00000					
Cumulative Proportion	1.00000	1.00000					

Let's make the main PC1 vs. PC2 figure.

```
ggplot(wisc.pr$x) +
  aes(PC1, PC2, col=diagnosis) +
  geom_point() +
  xlab("PC1 (44.3%)")
```



```
ylab("PC2 (19%)")
```

```
$y
[1] "PC2 (19%)"
```

```
attr("class")
[1] "labels"
```

Q4. From your results, what proportion of the original variance is captured by the first principal components (PC1)?

```
summary(wisc.pr)
```

Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	3.6444	2.3857	1.67867	1.40735	1.28403	1.09880	0.82172
Proportion of Variance	0.4427	0.1897	0.09393	0.06602	0.05496	0.04025	0.02251
Cumulative Proportion	0.4427	0.6324	0.72636	0.79239	0.84734	0.88759	0.91010
	PC8	PC9	PC10	PC11	PC12	PC13	PC14
Standard deviation	0.69037	0.6457	0.59219	0.5421	0.51104	0.49128	0.39624
Proportion of Variance	0.01589	0.0139	0.01169	0.0098	0.00871	0.00805	0.00523
Cumulative Proportion	0.92598	0.9399	0.95157	0.9614	0.97007	0.97812	0.98335
	PC15	PC16	PC17	PC18	PC19	PC20	PC21
Standard deviation	0.30681	0.28260	0.24372	0.22939	0.22244	0.17652	0.1731
Proportion of Variance	0.00314	0.00266	0.00198	0.00175	0.00165	0.00104	0.0010
Cumulative Proportion	0.98649	0.98915	0.99113	0.99288	0.99453	0.99557	0.9966
	PC22	PC23	PC24	PC25	PC26	PC27	PC28
Standard deviation	0.16565	0.15602	0.1344	0.12442	0.09043	0.08307	0.03987
Proportion of Variance	0.00091	0.00081	0.0006	0.00052	0.00027	0.00023	0.00005
Cumulative Proportion	0.99749	0.99830	0.9989	0.99942	0.99969	0.99992	0.99997
	PC29	PC30					
Standard deviation	0.02736	0.01153					
Proportion of Variance	0.00002	0.00000					
Cumulative Proportion	1.00000	1.00000					

0.4427

Q5. How many principal components (PCs) are required to describe at least 70% of the original variance in the data?


PC3 Cumulative Proportion = 0.72636, so PC1 + PC2 + PC3 are required

Q6. How many principal components (PCs) are required to describe at least 90% of the original variance in the data?

PC7 Cumulative Proportion = 0.91010, so PC1 + PC2 + PC3 + PC4 + PC5 + PC6 + PC7 are required

Q7. What stands out to you about this plot? Is it easy or difficult to understand? Why?


```
biplot(wisc.pr)
```



It is very difficult to read and understand because the data points are all overlapping with one another and cannot be distinguished from each other.

Q8. Generate a similar plot for principal components 1 and 3. What do you notice about these plots?

```
ggplot(wisc.pr$x) +  
  aes(PC1, PC3, col=diagnosis) +  
  geom_point() +  
  xlab("PC1 (44.3%)")
```



```
ylab("PC3 (9.4%)")
```

```
$y
[1] "PC3 (9.4%)"

attr("class")
[1] "labels"
```

There is a noticeable separation between the benign and malignant (looks like they are separated into 2 groups).

Q9. For the first principal component, what is the component of the loading vector (i.e. `wisc.pr$rotation[,1]`) for the feature `concave.points_mean`?

```
wisc.pr$rotation["concave.points_mean", 1]
```

```
[1] -0.2608538
```

Q10. What is the minimum number of principal components required to explain 80% of the variance of the data?

```
summary(wisc.pr)
```

Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	3.6444	2.3857	1.67867	1.40735	1.28403	1.09880	0.82172
Proportion of Variance	0.4427	0.1897	0.09393	0.06602	0.05496	0.04025	0.02251
Cumulative Proportion	0.4427	0.6324	0.72636	0.79239	0.84734	0.88759	0.91010

	PC8	PC9	PC10	PC11	PC12	PC13	PC14
Standard deviation	0.69037	0.6457	0.59219	0.5421	0.51104	0.49128	0.39624
Proportion of Variance	0.01589	0.0139	0.01169	0.0098	0.00871	0.00805	0.00523
Cumulative Proportion	0.92598	0.9399	0.95157	0.9614	0.97007	0.97812	0.98335

	PC15	PC16	PC17	PC18	PC19	PC20	PC21
Standard deviation	0.30681	0.28260	0.24372	0.22939	0.22244	0.17652	0.1731
Proportion of Variance	0.00314	0.00266	0.00198	0.00175	0.00165	0.00104	0.0010
Cumulative Proportion	0.98649	0.98915	0.99113	0.99288	0.99453	0.99557	0.9966

	PC22	PC23	PC24	PC25	PC26	PC27	PC28
Standard deviation	0.16565	0.15602	0.1344	0.12442	0.09043	0.08307	0.03987
Proportion of Variance	0.00091	0.00081	0.0006	0.00052	0.00027	0.00023	0.00005
Cumulative Proportion	0.99749	0.99830	0.9989	0.99942	0.99969	0.99992	0.99997

	PC29	PC30
Standard deviation	0.02736	0.01153
Proportion of Variance	0.00002	0.00000
Cumulative Proportion	1.00000	1.00000

The cumulative score of PC5 is 0.84734, which is the combined scores of PC1 + PC2 + PC3 + PC4 + PC5. So, you would need 5 PCs.


5. Combining methods

We can take our PCA results and use them as a basis set

Clustering on PCA results

```
wisc.pr.hclust <- hclust(dist(wisc.pr$x[, 1:2]), method="ward.D2")
plot(wisc.pr.hclust)
```

Cluster Dendrogram



```
dist(wisc.pr$x[, 1:2])
hclust (*, "ward.D2")
```

We can “cut” this tree to yield our clusters (groups):

```
pc.grps <- cutree(wisc.pr.hclust, k=2)
table(pc.grps)
```

```
pc.grps
 1    2
195 374
```

How do my cluster grps compare to the expert diagnosis

```
table(diagnosis, pc.grps)
```

```
      pc.grps
diagnosis  1    2
  B    18  339
  M   177   35
```

```
table(diagnosis)
```

```
diagnosis
  B   M
357 212
```

```
wisc.pr.hclust.clusters <- cutree(wisc.pr.hclust, k=4)
table(wisc.pr.hclust.clusters, diagnosis)
```

```
              diagnosis
wisc.pr.hclust.clusters  B   M
1         0 112
2        18  65
3       232  18
4       107  17
```

Q13. Which method gives your favorite results for the same data `data.dist` dataset? Explain your reasoning.

The ward.D2 method gives good results for the `data.dist` dataset because it minimizes the variance within the clusters, making them more neat and compact in comparison to the other methods listed (single, complete, average).

Q15. How well does the newly created model with four clusters separate out the two diagnoses?

The newly created model with four clusters makes it more difficult to separate out the two diagnoses because there are now 3 groups (out of 4) that have both benign and malignant cases. The distinction (compared to when there were 2 groups) is not as clear now.

Q16. How well do the k-means and hierarchical clustering models you created in previous sections (i.e. before PCA) do in terms of separating the diagnoses? Again, use the `table()` function to compare the output of each model (`wisc.km$cluster` and `wisc.hclust.clusters`) with the vector containing the actual diagnoses.

They did really badly. We do much better after PCA - the new PCA variables (what we call a basis set) give us much better separation of M and B.

```
table(wisc.pr.hclust.clusters, diagnosis)
```

```
              diagnosis
wisc.pr.hclust.clusters  B   M
1         0 112
2        18  65
3       232  18
4       107  17
```



```
wisc.km <- kmeans(wisc.data, centers= 2, nstart= 20)
table(wisc.km$cluster, diagnosis)
```

```
diagnosis
  B   M
1  1 130
2 356  82
```

7. Prediction

We can use our PCA model for the analysis of new “unseen” data. In this case from U. Mich.

```
url <- "https://tinyurl.com/new-samples-CSV"
new <- read.csv(url)
npc <- predict(wisc.pr, newdata=new)
npc
```

```
      PC1      PC2      PC3      PC4      PC5      PC6      PC7
[1,]  2.576616 -3.135913  1.3990492 -0.7631950  2.781648 -0.8150185 -0.3959098
[2,] -4.754928 -3.009033 -0.1660946 -0.6052952 -1.140698 -1.2189945  0.8193031
      PC8      PC9      PC10      PC11      PC12      PC13      PC14
[1,] -0.2307350 0.1029569 -0.9272861 0.3411457  0.375921 0.1610764 1.187882
[2,] -0.3307423 0.5281896 -0.4855301 0.7173233 -1.185917 0.5893856 0.303029
      PC15      PC16      PC17      PC18      PC19      PC20
[1,] 0.3216974 -0.1743616 -0.07875393 -0.11207028 -0.08802955 -0.2495216
[2,] 0.1299153  0.1448061 -0.40509706  0.06565549  0.25591230 -0.4289500
      PC21      PC22      PC23      PC24      PC25      PC26
[1,] 0.1228233 0.09358453 0.08347651 0.1223396 0.02124121 0.078884581
[2,] -0.1224776 0.01732146 0.06316631 -0.2338618 -0.20755948 -0.009833238
      PC27      PC28      PC29      PC30
[1,] 0.220199544 -0.02946023 -0.015620933 0.005269029
[2,] -0.001134152 0.09638361 0.002795349 -0.019015820
```

Q18. Which of these new patients should we prioritize for follow up based on your results?

You should prioritize PC1 patients because that group has the highest malignant to benign ratio (130 to 1), where almost all of the patients in that group (130) are malignant cases.