

A Discussion on Symmetry and Awesome Sets

An Introduction to the Isometries of \mathbb{R}^3

R. Baldwin

Math/Stat Department
Swarthmore College

Swarthmore College

Outline

1 Motivation

- A Reminder on the Definition
- Motivating Example

2 Discussion

- D_n and Isometries of \mathbb{R}^3
- Ideas Moving Forward

Outline

1 Motivation

- A Reminder on the Definition
- Motivating Example

2 Discussion

- D_n and Isometries of \mathbb{R}^3
- Ideas Moving Forward

What are *aba*-sets?

Also known as "awesome"-sets, A-sets

Definition

Let G be a group.

A set $S \subseteq G$ is an *aba-set* if for all $a, b \in S$, $aba \in S$.

■ Previous results:

- If $x \in S$ is an element of odd order, then $\langle x \rangle \in S$.

What are *aba*-sets?

Also known as "awesome"-sets, A-sets

Definition

Let G be a group.

A set $S \subseteq G$ is an *aba-set* if for all $a, b \in S$, $aba \in S$.

■ Previous results:

- If $x \in S$ is an element of odd order, then $\langle x \rangle \in S$.

How much can these sets generate?

Questions to consider:

- Can an *aba*-set with only generators and the identity recover the whole group?
- Is there a more geometric quality that *aba*-sets induce?

Let's look at an example to further explore this!

How much can these sets generate?

Questions to consider:

- Can an *aba*-set with only generators and the identity recover the whole group?
- Is there a more geometric quality that *aba*-sets induce?

Let's look at an example to further explore this!

How much can these sets generate?

Questions to consider:

- Can an *aba*-set with only generators and the identity recover the whole group?
- Is there a more geometric quality that *aba*-sets induce?

Let's look at an example to further explore this!

How much can these sets generate?

Questions to consider:

- Can an *aba*-set with only generators and the identity recover the whole group?
- Is there a more geometric quality that *aba*-sets induce?

Let's look at an example to further explore this!

Outline

1 Motivation

- A Reminder on the Definition
- **Motivating Example**

2 Discussion

- D_n and Isometries of \mathbb{R}^3
- Ideas Moving Forward

The Dihedral Group

$$\forall a, b \in S, aba \in S$$

Example

Let $S = \{1, \sigma, \zeta\} \subset D_n$ be a set with property *aba*-closure. What must follow is that S must contain ('generate') all powers of the flips and rotations.

Proof.

- Flips have order 2, meaning $\sigma = \sigma^{-1} \in S$ a priori.
- We can **build-up** all powers of *zeta* through different recursive combinations with the identity.
For example, $\zeta 1 \zeta = \zeta^2$ and $\zeta \zeta \zeta = \zeta^3$ and so forth.
- Thus, we can generate all powers of the flip- σ and rotation- ζ elements.



The Dihedral Group

$$\forall a, b \in S, aba \in S$$

Example

Let $S = \{1, \sigma, \zeta\} \subset D_n$ be a set with property aba -closure. What must follow is that S must contain ('generate') all powers of the flips and rotations.

Proof.

- Flips have order 2, meaning $\sigma = \sigma^{-1} \in S$ a priori.
- We can **build-up** all powers of zeta through different recursive combinations with the identity.
For example, $\zeta 1 \zeta = \zeta^2$ and $\zeta \zeta \zeta = \zeta^3$ and so forth.
- Thus, we can generate all powers of the flip- σ and rotation- ζ elements.



The Dihedral Group

$$\forall a, b \in S, aba \in S$$

Example

Let $S = \{1, \sigma, \zeta\} \subset D_n$ be a set with property *aba*-closure. What must follow is that S must contain ('generate') all powers of the flips and rotations.

Proof.

- Flips have order 2, meaning $\sigma = \sigma^{-1} \in S$ a priori.
- We can **build-up** all powers of *zeta* through different recursive combinations with the identity.
For example, $\zeta 1 \zeta = \zeta^2$ and $\zeta \zeta \zeta = \zeta^3$ and so forth.
- Thus, we can generate all powers of the flip- σ and rotation- ζ elements.



The Dihedral Group

$$\forall a, b \in S, aba \in S$$

Example

Let $S = \{1, \sigma, \zeta\} \subset D_n$ be a set with property aba -closure. What must follow is that S must contain ('generate') all powers of the flips and rotations.

Proof.

- Flips have order 2, meaning $\sigma = \sigma^{-1} \in S$ a priori.
- We can **build-up** all powers of ζ through different recursive combinations with the identity.
For example, $\zeta 1 \zeta = \zeta^2$ and $\zeta \zeta \zeta = \zeta^3$ and so forth.
- Thus, we can generate all powers of the flip- σ and rotation- ζ elements.



Outline

1 Motivation

- A Reminder on the Definition
- Motivating Example

2 Discussion

- D_n and Isometries of \mathbb{R}^3
- Ideas Moving Forward

Subgroups of \mathbb{R}^3

$$\forall a, b \in S, aba \in S$$

To find groups similar to the dihedral group, let's impose firm positions on the vertices of D_n on a plane.

In other words, let us look at what **space** D_n lives in!

Subgroups of \mathbb{R}^3

$$\forall a, b \in S, aba \in S$$

To find groups similar to the dihedral group, let's impose firm positions on the vertices of D_n on a plane.

In other words, let us look at what **space** D_n lives in!

Subgroups of \mathbb{R}^3

$$\forall a, b \in S, aba \in S$$

- The dihedral groups are all *subgroups* of the group of rotations in \mathbb{R}^3 ($SO(3)$).
- Other examples of finite subgroups of $SO(3)$ are "cyclic, tetrahedral, octohedral group."

Subgroups of \mathbb{R}^3

$$\forall a, b \in S, aba \in S$$

- The dihedral groups are all *subgroups* of the group of rotations in \mathbb{R}^3 ($SO(3)$).
- Other examples of finite subgroups of $SO(3)$ are "cyclic, tetrahedral, octohedral group."

Subgroups of \mathbb{R}^3

$$\forall a, b \in S, aba \in S$$

- The dihedral groups are all *subgroups* of the group of rotations in \mathbb{R}^3 ($SO(3)$).
- Other examples of finite subgroups of $SO(3)$ are "cyclic, tetrahedral, octohedral group."

Another Interesting Example

$$\forall a, b \in S, aba \in S$$

Remark

Recall that if $|x|$ is odd and in S , $\langle x \rangle \in S$. We can extend this to *any* cyclic subgroup, so long as we start with the identity! That is to say:

Let $x \in G$ have finite order. If $\{1, x\} \in S$, then $\langle x \rangle \in S$.

Furthermore, if G is a cyclic group, then the entire group is an *aba-group*!

Another Interesting Example

$$\forall a, b \in S, aba \in S$$

Remark

Recall that if $|x|$ is odd and in S , $\langle x \rangle \in S$. We can extend this to *any* cyclic subgroup, so long as we start with the identity! That is to say:

Let $x \in G$ have finite order. If $\{1, x\} \in S$, then $\langle x \rangle \in S$.

Furthermore, if G is a cyclic group, then the entire group is an *aba-group*!

Make Titles Informative.

Make Titles Informative.

Make Titles Informative.

Outline

1 Motivation

- A Reminder on the Definition
- Motivating Example

2 Discussion

- D_n and Isometries of \mathbb{R}^3
- Ideas Moving Forward

Groups to Inspect

- Focus on investigating the *aba*-sets of subgroups and semi-direct products of primarily \mathbb{R}^3 (a topic for next week).
- Observe if there is any correlation between said *aba* sets (specifically with sets only containing the generators and identity of these groups).

Groups to Inspect

- Focus on investigating the *aba*-sets of subgroups and semi-direct products of primarily \mathbb{R}^3 (a topic for next week).
- Observe if there is any correlation between said *aba* sets (specifically with sets only containing the generators and identity of these groups).

Programming

- Working on code that will speed up the process of finding *aba*-sets.
- *Potentially attempt to create a 3D program that may enumerate any *geometric* qualities of *aba*-sets.

Summary

- I am now not only investigating *aba*-sets from an algebraic viewpoint, but seeing if the **space** where a group lives in induces a constraint in algebraic results.
- Observing how *aba*-sets act on symmetries may lead into questions like how much of an original group is guaranteed to be **encoded** in an *aba*-set.
- While larger conjectures in \mathbb{R}^n would be awesome, I will likely continue to work with groups in \mathbb{R}^2 and \mathbb{R}^3 due to some computational concerns (things get *big* and confusing).

Summary

- I am now not only investigating *aba*-sets from an algebraic viewpoint, but seeing if the **space** where a group lives in induces a constraint in algebraic results.
- Observing how *aba*-sets act on symmetries may lead into questions like how much of an original group is guaranteed to be **encoded** in an *aba*-set.
- While larger conjectures in \mathbb{R}^n would be awesome, I will likely continue to work with groups in \mathbb{R}^2 and \mathbb{R}^3 due to some computational concerns (things get *big* and confusing).

Summary

- I am now not only investigating *aba*-sets from an algebraic viewpoint, but seeing if the **space** where a group lives in induces a constraint in algebraic results.
- Observing how *aba*-sets act on symmetries may lead into questions like how much of an original group is guaranteed to be **encoded** in an *aba*-set.
- While larger conjectures in \mathbb{R}^n would be awesome, I will likely continue to work with groups in \mathbb{R}^2 and \mathbb{R}^3 due to some computational concerns (things get *big* and confusing).

Thank you!