

Statistics, fisheries management and spatial modelling.

Towards spatio-temporal stock assessment methods.


Based on a collaboration between: Maxime Olmos, James Thorson,
André Punt, Cole Monnahan, Baptiste Alglaive, Cody Szuwalski.

December 2023




Some graphics as introduction

FIGURE 1 WORLD CAPTURE FISHERIES AND AQUACULTURE PRODUCTION




NOTES: Excluding aquatic mammals, crocodiles, alligators, caimans and algae. Data expressed in live weight equivalent.
SOURCE: FAO.

FIGURE 23 GLOBAL TRENDS IN THE STATE OF THE WORLD'S MARINE FISHERY STOCKS, 1974–2019



SOURCE: FAO.


Collapse of Atlantic cod stocks (East Coast of Newfoundland), 1992.





Source : Millennium Ecosystems Assessment

Landed value of fish in Newfoundland and Labrador, 1989 - 2009.

1989



2009




Mather, C. (2013). From cod to shellfish and back again? The new resource geography and Newfoundland's fish economy. *Applied Geography*, 45, 402-409.

How to assess and manage fish resources ?

➡ Stock assessment models

How to assess and manage fish resources ?

➡ Stock assessment models



How to build a stock assessment model?



Plan

- 1 Introduction
- 2 Basics of stock assessment models
- 3 The hierarchical formalism: state-space modelling
- 4 Towards spatio-temporal modelling of marine resources
- 5 Conclusion

Plan

- 1 Introduction
- 2 Basics of stock assessment models
- 3 The hierarchical formalism: state-space modelling
- 4 Towards spatio-temporal modelling of marine resources
- 5 Conclusion

Surplus production model

Logistic model

(Continuous formulation)

$$\frac{dN(t)}{dt} = r \cdot N(t) \left(1 - \frac{N}{K}\right)$$

(Discrete formulation)


$$N(t+1) = N(t) + r \cdot N(t) \left(1 - \frac{N(t)}{K}\right)$$

Schaefer model

$$N(t+1) = N(t) + r \cdot N(t) \left(1 - \frac{N(t)}{K}\right) - C(t)$$

with $C(t) = q \cdot e \cdot N(t)$

N : abundance, C : catch, K : carrying capacity, r : intrinsic rate of population increase, q : catchability, e : effort
 $B, K, r, t, q, f \in \mathbb{R}^+$



The middle term is known as the surplus production (denoted SP_t).

If $SP_t > C_t$,
population size increases

If $SP_t = C_t$,
population size is steady

If $SP_t < C_t$,
population size decreases

Surplus production model

Logistic model

(Continuous formulation)

$$\frac{dN(t)}{dt} = r \cdot N(t) \left(1 - \frac{N}{K}\right)$$

(Discrete formulation)


$$N(t+1) = N(t) + r \cdot N(t) \left(1 - \frac{N(t)}{K}\right)$$

Schaefer model

$$N(t+1) = N(t) + r \cdot N(t) \left(1 - \frac{N(t)}{K}\right) - C(t)$$

with $C(t) = q \cdot e \cdot N(t)$

N : abundance, C : catch, K : carrying capacity, r : intrinsic rate of population increase, q : catchability, e : effort
 $B, K, r, t, q, f \in \mathbb{R}^+$



The middle term is known as the surplus production (denoted SP_t).

If $SP(t) > C(t)$,
population size increases

If $SP(t) = C(t)$,
population size is steady

If $SP(t) < C(t)$,
population size decreases

Virtual population analysis

Catch is disaggregated into age-groups.

Cohort is the key concept.

A cohort comprises all the individuals (fish in this case) that were born in the same year.

For a single cohort:

$$N(t+1) = N(t) - C(t) - D(t)$$

with $D(t) = N(t) \cdot (1 - S)$


$$\Rightarrow N(t) = (N(t+1) + C(t))/S$$

$N(t)$: the abundance at the beginning of time t

$C(t)$: the catch during time t

$D(t)$: the amount of removals due to natural death during time t

S : the survival rate




Let's assume there are **no more fish left of the oldest age** (all caught or died)

⇒ calculate the numbers last year with only catch and mortality as input values

⇒ start from the oldest ages and move backwards to the youngest

In both cases

- criticism of the model
- criticism of the fitting/estimation procedure



Plan

- 1 Introduction
- 2 Basics of stock assessment models
- 3 The hierarchical formalism: state-space modelling
- 4 Towards spatio-temporal modelling of marine resources
- 5 Conclusion

Modelling hidden processes of population dynamics


Population dynamics

$$N(a+1, t+1) = N(a, t) \cdot \exp(-F(a, t) - M(a, t))$$

Exploitation dynamics (Baranov equation)

$$C(a, t) = \frac{F(a, t)}{F(a, t) + M(a, t)} \cdot N(a, t) \cdot [1 - \exp(-(F(a, t) + M(a, t)))]$$

And there is also Recruitment, but that's a whole different issue...



Modelling hidden processes of population dynamics


Population dynamics

$$N(a+1, t+1) = N(a, t) \cdot \exp(-F(a, t) - M(a, t))$$

Exploitation dynamics (Baranov equation)

$$C(a, t) = \frac{F(a, t)}{F(a, t) + M(a, t)} \cdot N(a, t) \cdot [1 - \exp(-(F(a, t) + M(a, t)))]$$

And there is also **Recruitment**, but that's a whole different issue...



The available data

Scientific survey data ($N_{obs}(a, t)$)



Commercial catch data ($C_{obs}(a, t)$)



Hierarchical modelling

Parameter
layer




Latent
layer




Observation
layer


Hierarchical modelling




Hierarchical modelling




Hierarchical modelling



Hierarchical modelling



Hierarchical modelling



Model inference

Let's denote \mathcal{Y} the observations and θ the parameters.

Frequentist approach

Search for the parameter values $\hat{\theta}$ that maximize the likelihood $P(\mathcal{Y}|\theta)$:

- ➡ Search for $\operatorname{argmax}_{\theta} (P(\mathcal{Y}|\theta))$

Bayesian approach

Compute the posterior distribution of the parameters θ knowing the data \mathcal{Y} through the Bayes rule:

$$P(\theta|\mathcal{Y}) = \frac{P(\mathcal{Y}|\theta)P(\theta)}{P(\mathcal{Y})}$$

In the Bayesian approach, **Markov Chain Monte Carlo** (MCMC) methods were very popular in the past decades.

- ➡ Provide a generic modelling and inference framework for fitting any kind of model

Model inference

Let's denote \mathcal{Y} the observations and θ the parameters.

Frequentist approach

Search for the parameter values $\hat{\theta}$ that maximize the likelihood $P(\mathcal{Y}|\theta)$:

- ➡ Search for $\operatorname{argmax}_{\theta} (P(\mathcal{Y}|\theta))$

Bayesian approach

Compute the posterior distribution of the parameters θ knowing the data \mathcal{Y} through the Bayes rule:

$$P(\theta|\mathcal{Y}) = \frac{P(\mathcal{Y}|\theta)P(\theta)}{P(\mathcal{Y})}$$

In the Bayesian approach, **Markov Chain Monte Carlo** (MCMC) methods were very popular in the past decades.

- ➡ Provide a generic modelling and inference framework for fitting any kind of model

Model inference

Let's denote \mathcal{Y} the observations and θ the parameters.

Frequentist approach

Search for the parameter values $\hat{\theta}$ that maximize the likelihood $P(\mathcal{Y}|\theta)$:

- ➡ Search for $\operatorname{argmax}_{\theta} (P(\mathcal{Y}|\theta))$

Bayesian approach


Compute the posterior distribution of the parameters θ knowing the data \mathcal{Y} through the Bayes rule:

$$P(\theta|\mathcal{Y}) = \frac{P(\mathcal{Y}|\theta)P(\theta)}{P(\mathcal{Y})}$$


In the Bayesian approach, **Markov Chain Monte Carlo** (MCMC) methods were very popular in the past decades.

- ➡ Provide a generic modelling and inference framework for fitting any kind of model

Complexifying the hidden process




Complexifying the hidden process




Complexifying the hidden process

Von Bertalanffy growth curve




Complexifying the hidden process



Complexifying the hidden process

Modelling population demography in space




Cao, J., Thorson, J. T., Punt, A. E., & Szuwalski, C. (2020). A novel spatiotemporal stock assessment framework to better address fine-scale species distributions: development and simulation testing. *Fish and Fisheries*, 21(2), 350-367.

Olmos, M., Cao, J., Thorson, J. T., Punt, A. E., Monnahan, C. C., Alglave, B., & Szuwalski, C. (2023). A step towards the integration of spatial dynamics in population dynamics models: Eastern Bering Sea snow crab as a case study. *Ecological Modelling*, 485, 110484.

Complexifying the hidden process

Modelling population demography in space




Cao, J., Thorson, J. T., Punt, A. E., & Szuwalski, C. (2020). A novel spatiotemporal stock assessment framework to better address fine-scale species distributions: development and simulation testing. *Fish and Fisheries*, 21(2), 350-367.

Olmos, M., Cao, J., Thorson, J. T., Punt, A. E., Monnahan, C. C., Alglave, B., & Szuwalski, C. (2023). A step towards the integration of spatial dynamics in population dynamics models: Eastern Bering Sea snow crab as a case study. *Ecological Modelling*, 485, 110484.


Plan

- 1 Introduction
- 2 Basics of stock assessment models
- 3 The hierarchical formalism: state-space modelling
- 4 Towards spatio-temporal modelling of marine resources
- 5 Conclusion


**Including the spatial component
add extra complexity !**




**Including the spatial component
add extra complexity !**



**Including the spatial component
add extra complexity !**



**Including the spatial component
add extra complexity !**



Limits of the standard methods of inference

Standard methods are no more efficient.

The keystone is the likelihood:

$$L_M(\theta) = P(Y|\theta) = \int_{\mathbb{R}^q} P(Y, \delta|\theta) d\delta$$

Y are the observations, θ are the parameters, δ are the latent variables.

In a spatio-temporal context q can be very high

- require efficient numerical methods (1) to bypass the integration step,
(2) to reduce the dimensionality of δ and (3) to derive efficiently the likelihood.

Three methods to enhance computing:

- Laplace approximation (= approximation of the likelihood)
- SPDE approach (= approximation of the Gaussian random effect)
- Automatic differentiation (= efficient derivation technics)

R package: R-INLA for bayesian inference (Rue et al., 2017) and TMB for maximum likelihood inference (Kristensen et al., 2015).

Limits of the standard methods of inference

Standard methods are no more efficient.

The keystone is the likelihood:

$$L_M(\theta) = P(Y|\theta) = \int_{\mathbb{R}^q} P(Y, \delta|\theta) d\delta$$

Y are the observations, θ are the parameters, δ are the latent variables.

In a spatio-temporal context q can be very high

- require efficient numerical methods (1) to bypass the integration step,
(2) to reduce the dimensionality of δ and (3) to derive efficiently the likelihood.

Three methods to enhance computing:

- Laplace approximation (= approximation of the likelihood)
- SPDE approach (= approximation of the Gaussian random effect)
- Automatic differentiation (= efficient derivation technics)

R package: R-INLA for bayesian inference (Rue et al., 2017) and TMB for maximum likelihood inference (Kristensen et al., 2015).

Limits of the standard methods of inference

Standard methods are no more efficient.

The keystone is the likelihood:

$$L_M(\theta) = P(Y|\theta) = \int_{\mathbb{R}^q} P(Y, \delta|\theta) d\delta$$

Y are the observations, θ are the parameters, δ are the latent variables.

In a spatio-temporal context q can be very high

- ➡ require efficient numerical methods (1) to bypass the integration step,
(2) to reduce the dimensionality of δ and (3) to derive efficiently the likelihood.

Three methods to enhance computing:

- Laplace approximation (= approximation of the likelihood)
- SPDE approach (= approximation of the Gaussian random effect)
- Automatic differentiation (= efficient derivation technics)

R package: R-INLA for bayesian inference (Rue et al., 2017) and TMB for maximum likelihood inference (Kristensen et al., 2015).

Limits of the standard methods of inference

Standard methods are no more efficient.

The keystone is the likelihood:

$$L_M(\theta) = P(Y|\theta) = \int_{\mathbb{R}^q} P(Y, \delta|\theta) d\delta$$

Y are the observations, θ are the parameters, δ are the latent variables.

In a spatio-temporal context q can be very high

- ➡ require efficient numerical methods (1) to bypass the integration step,
(2) to reduce the dimensionality of δ and (3) to derive efficiently the likelihood.



Three methods to enhance computing:

- Laplace approximation (= approximation of the likelihood)
- SPDE approach (= approximation of the Gaussian random effect)
- Automatic differentiation (= efficient derivation technics)


R package: R-INLA for bayesian inference (Rue et al., 2017) and TMB for maximum likelihood inference (Kristensen et al., 2015).

Some examples


Estimated density of big skate in the US waters of the California Current



Estimates of spatial variation in recruitment rate in the Gulf of Alaska




Spatiotemporal variation in exploitation rate for snow crab of Alaska



Exploitation Rate



Estimated spatial distribution and demography of a cohort of individuals over 3 years for cod



Plan

- 1 Introduction
- 2 Basics of stock assessment models
- 3 The hierarchical formalism: state-space modelling
- 4 Towards spatio-temporal modelling of marine resources
- 5 Conclusion

Take home message

A short history of stock assessment:

- **1950 - 1980:** development of the basis of stock assessment methods

Surplus production model, virtual population analysis.

- **1980 - 2010:** bringing these methods into statistics


State space modelling, maximum likelihood, bayesian inference, integrated approach.

- **2010 - today:** towards spatio-temporal modelling of population dynamics (among other things)

Spatio-temporal modelling, Laplace approximation, SPDE approach, automatic differentiation.

Does this mean that old methods are outdated ?

⇒ Of course not, all these methods are complementary.



Source : Ifremer, Diagnostic 2022 sur les ressources halieutiques débarquées par la pêche française (métropolitaine)



Take home message

A short history of stock assessment:

- **1950 - 1980:** development of the basis of stock assessment methods








Surplus production model, virtual population analysis.

- **1980 - 2010:** bringing these methods into statistics

State space modelling, maximum likelihood, bayesian inference, integrated approach.

- **2010 - today:** towards spatio-temporal modelling of population dynamics (among other things)

Spatio-temporal modelling, Laplace approximation, SPDE approach, automatic differentiation.



Does this mean that old methods are outdated ?

⇒ Of course not, all these methods are complementary.

Take home message

A short history of stock assessment:

- **1950 - 1980:** development of the basis of stock assessment methods







Surplus production model, virtual population analysis.

- **1980 - 2010:** bringing these methods into statistics

State space modelling, maximum likelihood, bayesian inference, integrated approach.

- **2010 - today:** towards spatio-temporal modelling of population dynamics (among other things)

Spatio-temporal modelling, Laplace approximation, SPDE approach, automatic differentiation.



Source : Ifremer, Diagnostic 2022 sur les ressources halieutiques débarquées par la pêche française (métropolitaine)



Does this mean that old methods are outdated ?

→ Of course not, all these methods are complementary.

Bibliography

- Bonfil, R. (2005). 10. Fishery stock assessment models and their application to sharks. Management techniques for elasmobranch fisheries, 154.
- Cao, J., Thorson, J. T., Punt, A. E., & Szuwalski, C. (2020). A novel spatiotemporal stock assessment framework to better address fine-scale species distributions: development and simulation testing. *Fish and Fisheries*, 21(2), 350-367.
- Krainski, E., Gómez-Rubio, V., Bakka, H., Lenzi, A., Castro-Camilo, D., Simpson, D., ... & Rue, H. (2018). Advanced spatial modeling with stochastic partial differential equations using R and INLA. Chapman and Hall/CRC.
- Kristensen, K., Thygesen, U. H., Andersen, K. H., & Beyer, J. E. (2014). Estimating spatio-temporal dynamics of size-structured populations. *Canadian Journal of Fisheries and Aquatic Sciences*, 71(2), 326-336.
- Kristensen, K., Nielsen, A., Berg, C. W., Skaug, H., & Bell, B. (2015). TMB: automatic differentiation and Laplace approximation. arXiv preprint arXiv:1509.00660.
- Mather, C. (2013). From cod to shellfish and back again? The new resource geography and Newfoundland's fish economy. *Applied Geography*, 45, 402-409.
- Olmos, M., Cao, J., Thorson, J. T., Punt, A. E., Monnahan, C. C., Alglave, B., & Szuwalski, C. (2023). A step towards the integration of spatial dynamics in population dynamics models: Eastern Bering Sea snow crab as a case study. *Ecological Modelling*, 485, 110484.
- Rue, H., Riebler, A., Sørbye, S. H., Illian, J. B., Simpson, D. P., & Lindgren, F. K. (2017). Bayesian computing with INLA: a review. *Annual Review of Statistics and Its Application*, 4, 395-421.
- Thorson, J. T., Jannot, J., & Somers, K. (2017). Using spatio-temporal models of population growth and movement to monitor overlap between human impacts and fish populations. *Journal of Applied Ecology*, 54(2), 577-587.
- Thorson, J. T., Ianelli, J. N., Munch, S. B., Ono, K., & Spencer, P. D. (2015). Spatial delay-difference models for estimating spatiotemporal variation in juvenile production and population abundance. *Canadian journal of fisheries and aquatic sciences*, 72(12), 1897-1915.
- Xiao, Y. (2005). Catch equations: restoring the missing terms in the nominally generalized Baranov catch equation. *Ecological modelling*, 181(4), 535-556.

Modelling hidden processes of population dynamics

Age-dependent only (*Continuous formulation*)

$$\frac{dN(a)}{da} = -(F + M)N(a)$$

$$\Rightarrow N(a) = N(a_0) \exp(-(F + M)(a - a_0)), a_0 \leq a$$

$$C(a, \Delta a) = \int_a^{a+\Delta a} F \cdot N(a) da \quad (\text{Baranov equation})$$
$$= \frac{F}{F + M} N(a) [1 - \exp(-(F + M)\Delta a)]$$


Usually, age is discretized and the catch is modelled between age a and $a + 1$ ($\Delta a = 1$). Then, it gives:

$$C(a) = \frac{F}{F + M} N(a) [1 - \exp(-(F + M))]$$

Age- and time-dependent (*Discrete formulation*)

$$N(a + 1, t + 1) = N(a, t) \exp(-F(a, t) - M(a, t))$$

$$C(a, t) = \frac{F(a, t)}{F(a, t) + M(a, t)} N(a, t) [1 - \exp(-(F(a, t) + M(a, t)))]$$



And there is also **Recruitment**, but that's a whole different issue...

Laplace approximation

Aim: approximate the integral where $f(x)$ is the log-likelihood of a Gaussian variable x :

$$I_n = \int_x \exp(nf(x))dx \quad \text{as } n \rightarrow \infty$$

Let's denote x_0 , the point in which $f(x)$ has its maximum, then

$$\begin{aligned} I_n &\approx \int_x \exp[n(f(x_0) + \frac{1}{2}(x - x_0)^2 f''(x_0))]dx \\ &= \exp(nf(x_0)) \sqrt{\frac{2\pi}{-nf''(x_0)}} = \tilde{I}_n. \end{aligned}$$

Application to likelihood:

If $f(\theta, \delta)$ is the joint log-likelihood, then $L_M(\theta)$ can be written as:

$$\begin{aligned} L_M(\theta) &\approx L_M^*(\theta) \\ &= (2\pi)^{q/2} |H(\theta)|^{-1/2} \exp[f(\theta, \hat{\delta}_\theta)] \end{aligned}$$


$H(\theta)$: the Hessian of the negative joint log-likelihood — f

q : the size of the latent effect

$$\hat{\delta}_\theta = \underset{\delta}{\operatorname{argmax}}(f(\theta, \delta))$$

Conditions required:

- normality of the random effect δ
- some regularity conditions on f (e.g. only one maximum)




The true marginal (blue), the Laplace approximation (red) and the Gaussian approximation (orange).

SPDE approach

$$\delta | \theta \sim GF(\mu, \Sigma) \quad \text{Latent Gaussian field}$$


$$Y | \delta, \theta \sim \mathcal{L}(\delta | \theta) \quad \text{Observation process}$$

- Y : observations.
- θ : parameters.
- δ : a spatial random effect (= Gaussian field)
- $\Sigma = Q^{-1}$: variance-covariance matrix of a spatial random effect.
 Q is the precision matrix.




First set of results

A Gaussian field with Matérn Covariance can be represented as a Gauss-Markov field \Rightarrow sparse representation of the spatial effect



Second set of results




SPDE approach

$$\delta|\theta \sim GF(\mu, \Sigma) \quad \text{Latent Gaussian field}$$


$$Y|\delta, \theta \sim \mathcal{L}(\delta|\theta) \quad \text{Observation process}$$

- Y : observations.
- θ : parameters.
- δ : a spatial random effect (= Gaussian field)
- $\Sigma = Q^{-1}$: variance-covariance matrix of a spatial random effect.
 Q is the precision matrix.



First set of results

A Gaussian field with Matérn Covariance can be represented as a Gauss-Markov field \Rightarrow sparse representation of the spatial effect



Second set of results

