



Survey Header

Username: Rohit, Age: 21.0

Percentage of Correct Answers by Domain



Percentage of Correct Answers by Subdomain



Survey Report

Anxiety

4 = The subject reports severe anxiety. The subject has no control over their anxiety when it occurs and cannot turn their attention to other things, even when a distinct and sustained effort is made. At least one marked and persistent sign of motor tension or autonomic hyperactivity should accompany the anxiety. The anxiety has been present in this form on the majority of days in the last month.

Muscular Tension

Signs of motor tension include: physical restlessness, trembling, involuntarily tensed muscles, tension pains affecting neck, back or legs and tension headaches. Signs of autonomic hyperactivity include: gastro-intestinal: dry mouth, difficulty swallowing, epigastric discomfort, frequent loose motions; respiratory: feeling of constriction in the chest, difficulty inhaling, hyperventilation; cardiovascular: discomfort over the heart, palpitations, missed heartbeats, throbbing in the neck; genitourinary: frequency and urgency of micturition, failure of erection, lack of libido, increased menstrual discomfort; nervous system: tinnitus, blurring of vision, dizziness, prickling sensations, sweating, blushing.

Situational Anxiety

Some of the subject's utilise avoidance strategies as a means of coping with their anxiety.

They may report experiencing little or no anxiety in the previous month because they have avoided those situations which would have provoked anxiety. For example, a person who experiences severe anxiety in public situations may have avoided this by staying at home all the time, relying on a relative or other carer to carry out essential tasks like shopping or going to work. In these circumstances it is recommended that the score for anxiety should be based on the level of reported anxiety experienced by the subject, but the presence of avoidance strategies, the frequency with which they are employed and the disruption they cause to the person's social functioning should also be noted.

Depression

the sample recommendations