# Males and Females (Machine) Learning to Express Emotions More Granularly

Bradley Allen

School of Computing and Information Sciences, Willamette University

DATA 515: Advanced Applied Machine Learning

August 14, 2026

## Abstract

As Americans report heightened levels of depression, anxiety, and stress, the need to identify underlying predictors of mental health is increasingly important. Identifying one's emotions with granular detail is associated with down regulation of negative emotions. However, the frequency and intensity one experiences these mental health indicators differ by gender, and the individual capacity to identify emotions is highly variable. Existing research has explored gender differences in diagnoses and behavioral affect of females and males with moderate to severe depression, anxiety, and stress, but the relative importance of features predicting these mental health indicators by gender is under explored. This study leveraged machine learning algorithms to identify gender differences in important features predictive of mental health in a sample of 38,835 participants. Results revealed the relative importance of anxiety and stress features were similar for females and males. By contrast, female depression was better predicted by anxiety and stress features, while male depression was best predicted by features associated with depression. Similarly, personality was the strongest predictor of female depression, anxiety, and stress, while male stress was better predicted by features such as family size and sexual orientation. Finally, emotional stability was more important for males across all three mental health indicators than for females. These results highlight the need to equip females and males with more granular emotional language to parse their experiences with, and sources of, depression, anxiety, and stress.

"When we can talk about our feelings, they become less overwhelming, less upsetting, and less scary." – Fred Rogers (1994, p. 115)

Whether the advice comes from a beloved neighbor like Mr. Rogers or emotional intelligence researchers Peter Salovey and John D. Mayer (1990), the ability to identify and regulate emotions is essential for developing a healthy personality, cultivating friendships, and enacting social change.

As Americans report worsening mental health (Jones, 2025), elevated levels of depression (Witters, 2026), heightened anxiety (American Psychiatric Association, 2026), and the highest levels of stress in decades (Fioroni & Foy, 2024), the need to identify physiological and behavioral symptoms of mental health indicators such as depression, anxiety, and stress is of increasing importance. After all, as Mr. Rogers put it, "Anything that's human is mentionable, and anything that is mentionable can be more manageable" (1994, p. 115).

Indeed, labeling emotions is associated with regulation of distress and other negative affect emotions (Kashdan et al., 2015; Kircanski et al., 2012; Levy-Gigi & Shamay-Tsoory, 2022; Yue et al., 2025). Problematically, research by authors Bradberry and Greaves (2009) revealed that only 36% of people could identify their emotions when they experienced them (p. 19). Nook et al. (2020) found that while emotional vocabulary increased during childhood, it plateaued between 11 and 18 years of age.

Given the importance of identifying emotions in developing healthy emotional regulation (Mayer & Salovey, 1995; Salovey & Mayer, 1990), the gap between emotional experience and the granularity with which a person can verbalize that experience has become a field of study in psychology.

**Emotional Granularity**

The term emotional granularity was coined by researcher Lisa Barrett (2004) to describe the level of verbal precision with which a person identified emotions they experienced. An individual who rates low in emotional granularity is said to report their experiences in general terms as positively or negatively valenced. An individual who rates high in emotional granularity is said to report to their emotional experiences in more descriptive terms (e.g., happy, sad, angry) such that a distinct feeling is conveyed.

The ability to distinctly describe one's emotions is associated healthier coping mechanisms and better mental health (Kashdan et al., 2015; Smidt & Suvak, 2015). For example, individuals who rate high on emotional granularity are less prone to maladaptive behavior like hitting or binge eating (Dixon-Gordon et al., 2014).

Furthermore, individuals with high emotional granularity self-report more frequent emotional regulation (Barrett et al., 2001) and increased resilience (Tugade et al., 2004), while

individuals low on emotional granularity tend to be less successful in downregulating negative emotions (Kalokerinos et al., 2019).

Taken together, the connection between experiencing and explaining is beneficial to wellbeing such that providing people with the vocabulary clinicians use to describe the physiological and behavioral predictors of depression, anxiety, and stress is a necessary undertaking.

**Gender Differences**

The extent to which individuals experience indicators of depression, anxiety, and stress can depend on gender. For instance, females experience a higher lifetime prevalence and diagnosis of anxiety disorders (Kessler et al., 2005; McLean et al., 2011) and score higher on scales measuring stress (Graves et al., 2021; Matud, 2004) compared to males.

Additionally, females rate lower on personality trait emotional stability (Lippa, 2010; Weisberg et al., 2011), and although personality tends to be a stronger predictor of depression, anxiety, and stress for males, emotional stability is the best personality predictor for depression, anxiety and stress for both males and females (Alizadeh et al., 2018; Uliaszek et al., 2010).

Evidence also suggests that females rely on more emotion-focused coping styles for stress compared to males (Matud, 2004), and in response to depression, females are more prone to internalizing behavior (Levkovich et al., 2025; Li et al., 2023), and rumination (Johnson & Whisman, 2013).

To date, much of the research on gender differences in experiencing depression, anxiety, and stress has focused either on the magnitude of the differences (Salk et al., 2017), the behavioral expressions of individuals experiencing high levels of one or the other (Kuehner, 2017; Sloan & Sandt, 2006), or associated factors such as biology (Donner & Lowry, 2013; Farhane-Medina et al., 2022).

For example, a widely cited study by Martin et al. (2013) found that although the proportion of females and males who experience depression is about the same (e.g., 33.3% vs. 30.6%, respectively), males reported higher rates of anger, aggression, substance abuse, and risk taking. Conversely, females tend to report depressed mood, appetite and sleep disturbance with higher frequency and intensity compared to males (Cavanagh et al., 2017).

While there is a broad swath of research on gender differences in symptomatic expression, a topic that remains under explored is whether females and males differ in how depression, anxiety, and stress are experienced in self-reported measures (e.g., does dysphoria better predict depression for females while hopelessness better predicts depression for males?). Given gender differences in diagnoses and symptomatic expression, it is plausible the underlying subscales of measures for depression, anxiety, and stress would differently predict symptoms for females and males.

## DASS Root Vocabulary

The 21-item Depression Anxiety Stress Scales (DASS-21) is a widely used and rigorously validated psychometric tool that operationally defines depression, stress, and anxiety in terms of clinically aligned physiological and behavioral subscales (Adu et al., 2025; Antony et al., 1998; Dwight et al., 2024; Henry & Crawford, 2005; P. F. Lovibond & Lovibond, 1995; S. H. Lovibond & Lovibond, 1995).

Researchers have demonstrated population validity in samples of youths (Dwight et al., 2024), across cultures (Adu et al., 2025; Henry & Crawford, 2005; P. F. Lovibond & Lovibond, 1995), and between the sexes (Gomez et al., 2014). Therefore, the DASS-21 subscales are an ideal source of clinically aligned, emotionally granular vocabulary.

The subscales of DASS-21 describe physiological and behavioral predictors. For example, depression is characterized by hopelessness (i.e., nothing to look forward to), while anxiety is characterized by autonomic arousal such as dryness of mouth or difficulty breathing (Antony et al., 1998; P. F. Lovibond & Lovibond, 1995). Table 1 shows the DASS-21 scales with the associated subscales, items, and root vocabulary. Parsing these predictors and properly labelling them should, in theory, add to an individual's capacity for emotional granularity, yielding all the associated benefits.

## Present Study

This study proposes applying machine learning models to the Answers to the Depression Anxiety Stress Scales data set available from the Open Psychometrics Project (2019). In a developing study by Bradley Allen (2026), confirmatory factor analysis was conducted on this dataset confirming the DASS-21 measure functioned the same for both females and males, allowing for comparisons between genders.

The data further replicated findings of gender differences in personality, personality as a predictor of depression, anxiety, and stress, as well as findings that females tended to experience higher levels of all three mental health indicators compared to males. Therefore, the Answers to the Depression Anxiety Stress Scales data set is a strong candidate for generalizable findings of predictors of depression, anxiety, and stress.

Existing machine learning research has analyzed this data set to predict depression, anxiety, and stress using demographic variables such as family size, gender, and race (Kumar et al., 2020; Yesudas, 2023). Other applied machine learning research has analyzed physiological predictors of depression, anxiety, and stress such as skin conductance and heart rate (Al Abdi et al., 2026; Liu et al., 2025). A metanalysis of models predicting depression, anxiety, and stress revealed the most common models were kNN, SVM, random forest, and ensemble (Taskynbayeva & Gutoreva, 2025).

Despite the extensive research on this topic, whether subscales of the DASS-21 are better predictors for depression, anxiety, and stress for females and males remains under explored. This project aims to extend research on this topic in two ways. First, by applying newer models such as XGBoost, LightGBM, and CatBoost. Second, by using these models in a novel way to predict moderate to severe levels of depression, anxiety, and stress for males and females. Toward that end, this study aims to answer the following:

RQ1: Do these models support the discriminant validity of the DASS-21 constructs via important feature extraction?

RQ2: Do important features vary between males and females with respect to DASS-21 subscales and demographic features as predictors of depression, anxiety, and stress?

RQ3: If important features vary by gender, what might this suggest about the value of learning clinically aligned vocabulary?

These findings will help males and females in multiple ways. First, if gender differences in physiological and behavioral predictors exist, these findings will help males and females identify symptomatic experiences of depression, anxiety, or stress separately. Properly identifying symptoms will help males and females communicate the feelings they experience to one another, as well as enhance interactions with mental health care professionals by using more precise language.

## Method

### Participants

The raw data contained 39,775 responses. Participants were required to pass a series of checks for their responses to be included. First, the survey included a 16-word list (e.g., boat, paucity, lucid), 3 of which were fake (e.g., cuivocal, florted, verdid). Participants were instructed to check a box next to words whose definitions they knew for certain. Those who indicated they knew the definitions of all 3 fake words were excluded from the analysis, resulting in a reduction of 326 participants.

Second, participants were required identified as male or female. As a result, 540 participants who selected “other” and 67 participants who did not provide a response were excluded. Finally, 7 participants who listed their age as greater than 100 were removed. The remaining sample size was 38,835.

### Measures

#### *Depression, Anxiety, and Stress*

Depression, anxiety, and stress scores were calculated using the 21-item Depression Anxiety Stress Scale (DASS-21; P. F. Lovibond & Lovibond, 1995). Depression was assessed using a 7-item scale that included items such as “I felt sad and depressed” and “I just couldn’t

seem to get going." Anxiety was assessed using a 7-item scale that included items such as "I had difficulty swallowing" and "I felt terrified." Stress was assessed using a 7-item scale that included items such as "I found it hard to wind down" and "I tended to over-react to situations."

Items for all three scales were measured using a 4-point Likert scale (0 = *Did not apply to me at all*, 3 = *Applied to me very much, or most of the time*). Scores were then doubled for the DASS-21 subset (S. H. Lovibond & Lovibond, 1995). Descriptive statistics are shown in Table 2.

### *Personality*

Personality scores were calculated using the Ten Item Personality Inventory (TIPI; Gosling et al., 2003). Participants responded to the prompt, "I see myself as" and rated items such as "Extraverted, enthusiastic" and "Calm, emotionally stable." Extraversion, agreeableness, conscientiousness, emotional stability, and openness to experience were all assessed using a 2-item scale, on a 7-point Likert scale (1 = *Disagree strongly*, 7 = *Agree strongly*).

Openness to experience measures intellectual curiosity, creativity, and the preference of novelty. Conscientiousness measures self-discipline, orderliness, and persistence toward goals. Extraversion measures traits such as energy, assertiveness, sociability, and the tendency to seek stimulation in the company of others. Agreeableness measures a person's tendency to be compassionate, cooperative, and seek social harmony. Emotional stability is the tendency to regulate negative emotions and exhibit low reactivity to stress, anxiety, and environmental pressure.

## Procedure

The Open Psychometrics (2019) data set Answers to the Depression Anxiety Stress Scales was explored. Observations included in the data were gathered through an online survey from 2017 to 2019 and recorded 39,775 responses. The raw data frame contained 172 variables, including 42 items for the Depression Anxiety Stress Scales (DASS; P. F. Lovibond & Lovibond, 1995), 10 items from the Ten Item Personality Inventory (TIPI; Gosling et al., 2003), 16 items for the attention check (i.e., one variable for each word), and a series of time related questions (e.g., how much time spent on each question, total survey completion time) and demographic variables (e.g., education level, college major, gender, religion, orientation).

## Data Preprocessing and Analytic Strategy

### *Software*

To analyze the data, 7 models were trained and tested using Python (Python Software Foundation, 2025) with Scikit-learn (Pedregosa et al., n.d.) in Visual Studio Code (Microsoft, 2026). Models included Random Forest (Breiman, 2001), AdaBoost (Freund & Schapire, 1997), Gradient Boosting (Friedman, 2001), Histogram Gradient Boosting (Friedman, 2001), XGBoost (Chen & Guestrin, 2016), LightGBM (Ke et al., 2017), and CatBoost (Prokhorenkova et al.,

2018). Additional software packages used in this project were matplotlib (Hunter, 2007), numpy (Harris et al., 2020), pandas (The Pandas Development Team, 2020), and Seaborn (Waskom, 2021).

DASS-21 and demographic plots contrasting feature importance by gender were created in R Studio (Posit team, 2026) using R (R Core Team, 2026) libraries patchwork (Pedersen, 2025), tidytext (Robinson et al., 2026), and tidyverse (Wickham et al., 2019).

### *Model Design*

The data was filtered into two data frames, one for males and one for females. The data frames were then split into one containing DASS-21 variables, and another containing demographic variables. These data frames were used to identify the top DASS-21 subscales and demographic predictors of mental health (i.e., depression, anxiety, and stress) for males and females, respectively.

In observance of cost-sensitive machine learning practices for applied psychology, Randomized Search CV was applied to optimize model parameter tuning (Feng et al., 2026; Sterner et al., 2025). Parameters included 20 n-iterations, 5-fold cross validation, and accuracy as the scoring metric. Models included "honesty" provisions such as minimum sample splits, variable tree depth, and weighting described by Athey et al. (2019). Given the large sample size, minimum sample splits were set to 30 to ensure that each node generalized to the population under the Central Limit Theorem.

Finally, the best model parameters were extracted, then applied to an Ensemble model. Models were trained on 80% of the data and tested on 20%. In all cases, random state was set to 42 for replicability.

The Ensemble model was used to find the best predictors of depression, anxiety, and stress for males and females, respectively. Soft voting was applied for its ability to capture the nuances of complex, multi-dimensional mental health data when using Ensemble with models such as decision trees, XGBoost, CatBoost, and LightGBM (Gupta & Pandit, 2026; Jain et al., 2025; Pavani et al., 2024). Data preprocessing was required before models were trained.

### *DASS-21 Clinical Cut-off Scores*

To facilitate classification, binary labels were created from established clinical cut-off scores by S.H. Lovibond and Lovibond (1995). Using the DASS-21 measure, scores within the normal to mild range were assigned to 0, while scores within the moderate to extremely severe range were assigned to 1. Score ranges are shown in Table 3.

Before the scores were calculated, respondent values needed to be adjusted downward by 1, as the original scale range was 0-3, but the survey calculated them from 1-4. This brought the values in line with the DASS-21 clinical cut-off ranges (P. F. Lovibond & Lovibond, 1995; S. H. Lovibond & Lovibond, 1995; Soria-Reyes et al., 2024).

### *Personality Reverse Scoring*

The TIPI items were assessed using a 7-point Likert scale, which included 5 reverse-scored items (i.e., questions 2, 4, 6, 8, and 10). Reversed scored items were calculated by subtracting each response from 8 (e.g., 8 - 1, 8 - 2, etc.) before taking the mean of each set and assigning them to their respective trait (Gosling et al., 2003).

### *Grouped Categories*

To address model bias in decision trees, religion and age groups were assigned numerical categories. These variables contained multiple categories or high ranges, which decision trees favor as important features to split on (Nembrini et al., 2018; Strobl et al., 2007). For instance, age ranged from 13 to 89 (Table 4), and religion contained 12 categories. Meaningful age groups (Geifman et al., 2013) were assigned as follows: 1 = *Adolescent*, 2 = *Young adult*, 3 = *Adult*, 4 = *Middle aged*, 5 = *Aged*, 6 = *80 and over* (Table 2).

Religion was converted to a binary variable (0 = *Not religious*, 1 = *Religious*). Participants who selected agnostic and atheist were assigned to the not religious group, while participants who selected Buddhist, Christian, Hindu, Jewish, Muslim, Sikh, and other were assign to the religious group (Table 2).

### *Casting Categorical Variables to Data Type Categorical*

Categorical variables in machine learning have been known to contribute to model bias in machine learning through spurious mathematical correlation (Kopp et al., 2025; Zhu et al., 2024). To eliminate a potential source of bias, ordered categorical variables were cast to type category using Pandas astype (Pandas via NumFOCUS, Inc., 2023).

Education was ordered by level of attainment (e.g., 1 = *Less than high school*, 4 = *Graduate degree*). Age group was ordered by epoch (1 = *Adolescent*, 6 = *80 and over*). The remainder of categorical variables were unordered (i.e., urban, gender, orientation, race, voted, married).

DASS-21 and TIPI items were either sums or averages, respectively, and thus treated as continuous numerical variables (Gosling et al., 2003; P. F. Lovibond & Lovibond, 1995).

### *Class Imbalance*

After applying the binary classifier to participant DASS scores, notable class imbalances were observed. Differences were more common for females, with 30.65%, 29.39%, and 38.56% experiencing low to mild symptoms of depression, anxiety, and stress, compared to 69.35%, 70.61%, and 61.44% experiencing moderate to severe symptoms, respectively (Table 2).

The proportion of males with anxiety and stress was approximately equal at 41.89% and 52.62% experiencing low to mild symptoms, compared to 58.11% and 47.38% experiencing moderate to severe symptoms. The largest difference was on depression, with 34.97% of males

experiencing low to mild symptoms, compared to 65.03 experiencing moderate to severe symptoms (Table 2).

To address class imbalance, class weight was set to “balanced” for Random Forest, LightGBM, and CatBoost. A custom weight was applied to XGBoost by multiplying by the proportion of negative to positive cases (i.e., low to mild divided by moderate to high). Weights were not applied to AdaBoost, Gradient Boosting, or HistGradientBoost.

### *Missing Data*

No patterns of missing data were found for DASS-21, TIPI, gender, or other demographic variables.

### *Evaluation*

Model performance was evaluated using accuracy, precision, recall, and F1 score. Accuracy measures the overall percentage of correct predictions, calculated as the ratio of true results (both positive and negative) to all total outcomes. Precision evaluates the reliability of the model's positive predictions—such as identifying cases of moderate-to-severe depression—by calculating the ratio of true positives to all predicted positives. Recall assesses the model’s capacity to capture all actual positive cases by calculating the ratio of true positives to all true positive and false negative outcomes. F1 score is the ratio of the product of precision and recall over the sum of precision and recall, and is used as a balanced summary metric to indicate whether the model was over optimized for one metric over the other (Sujon et al., 2025; Yesudas, 2023)

## Results

### DASS-21 Subscales as Predictors of Depression, Anxiety, and Stress

### *Model Performance*

As expected, the baseline models performed with nearly 100% accuracy. This is the case because the subscales should accurately predict the scale (e.g., the subscales aggregated for the depression score should accurately predict depression). The Ensemble model performed equally well for males and females, with universal accuracy at or above 99% (Table 5). Optimized parameter values and model performance contrasts by gender are included in the Supplement (Tables S1, S2, S3; Figs. S1, S2, S3).

Allowing model parameters to freely vary between females and males resulted in average differences in model parameters. For example, to predict anxiety, AdaBoost models were optimized for females vs. males with 1000 vs 500 n-estimators, .10 vs .50 learning rate, and entropy vs. Gini criterion (Table S2). This suggests that Random Grid Search CV was useful in tuning cost-sensitive models (Sterner et al., 2025).

### *Feature Importance*

Feature importance was calculated and extracted from the Ensemble model. The values were standardized to allow between group comparisons by dividing each feature score by the highest feature score. The result is a relative feature score where each is a fraction of the highest feature score (Table 6).

The feature scores for anxiety and stress performed as expected for both females and males, with important features strictly grouping under the associated scale (Table 1). Anxiety, for example, had a top feature of autonomic arousal, followed by subjective experience of anxious affect, situational anxiety, and skeletal musculature effects, all of which ranked 74% or higher for females and 80% or higher for males (Fig. 3). All other subscales (i.e., stress and depression) ranked 2% or less for females, and 5% or less for males. The order of importance was approximately the same for both females and males. Stress subscales ranged from 68% to 90% of the top stress features for females, and 70% to 92% for males (Table 6). The order of importance was approximately the same (Fig. 1). All other subscales (i.e., anxiety and depression) ranked 11% or lower for females, and 27% or lower for males. Thus, indicating strong discriminant validity for both anxiety and stress.

By contrast, depression trended higher across all features, with a minimum relative feature importance of 47% for females, and 35% for males (Table 6). Additionally, there were no shared features between females and males in the top 5 important features (Table 6). The top 5 features for females included a mixture of anxiety and stress subscales (e.g., easily upset, impatient, irritable / over-reactive) but none for depression. Conversely, the top 5 features for males were strictly subscales associated with depression (e.g., inertia, hopelessness). See Fig. 2 for chart of top features in ascending order by gender. These results indicate DASS-21 depression subscales perform differently as predictors depression for females experiencing moderate to severe symptoms when compared to males.

## Demographic Information as Predictors of Depression, Anxiety, and Stress

### *Model Performance*

Baseline models demonstrated variable performance for females and males, with accuracy fluctuating between 70% to 76% for depression (Table S4; Fig. S4), anxiety (Table S5; Fig. S5), and stress (Table S6; Fig. S6). On balance, the Esemble models performed higher than the baseline for females and males with depression at 76% for both, anxiety at 74% and 72%, and stress at 75% and 74% accuracy, respectively (Table 7). Given the extent to which demographic variables may only be tangentially related to depression, anxiety, and stress, these rates should be expected to trend lower than the subscales of DASS-21. Confusion matrices revealed that demographic variables were between 20 and 30% less accurate predicting negative cases of depression, anxiety, and stress compared to positive cases (Figs. S7, 8, 9). This relation held for all models except male stress, where accuracy was 73% in both cases.

### *Important Features*

Personality tended to be the best predictor of depression, and anxiety for both females and males. Across depression, anxiety and stress, the top three predictors for females were extraversion, conscientiousness, and agreeableness, all of which ranked 80% or higher compared to the top feature (Table 8). For anxiety and stress, the next top predictors were family size and openness to experience, while emotional stability, then family size were the next best predictors for depression (Table 8; Fig. 3).

Similarly, personality was also the top predictor of depression and anxiety for males. Depression was best predicted by extraversion, conscientiousness, emotional stability, openness to experience, then family size. Anxiety was best predicted by extraversion, emotional stability, conscientiousness, agreeableness, and family size (Table 8; Fig. 3). Notably, compared to females, the relative importance of each subsequent feature from the main feature for males was 3% to 20% lower. This suggests that a broader range of features elicits depression, anxiety, and stress in females compared to males.

Conversely, the top features indicative of moderate to severe stress in males differed significantly. Compared to anxiety and depression, the top feature for males was emotional stability, followed by family size, sexual orientation, agreeableness, and education level (Fig. 4). In total, personality was not as predictive of stress for males.

## Discussion

Results revealed strong discriminant validity for stress and anxiety subscales as important features in predicting stress and anxiety, respectively. However, ensemble machine learning revealed distinct patterns in relative feature importance for females and males who reported moderate to severe levels of depression. For instance, male depression was best predicted by depression features, compared to females where depression was best predicted by a mixture of anxiety and stress features. This suggests that the discriminant validity of DASS-21 depression subscales as predictors of clinically high levels of depression, vary by gender (i.e., strong for males, and potentially weak for females due to interrelatedness with anxiety and stress).

Interrelated important features sheds some light into why females might experience higher lifetime diagnoses of anxiety disorders and score higher on scales measuring stress compared to males (Graves et al., 2021; Kessler et al., 2005; Matud, 2004; McLean et al., 2011). It further suggests that what it means to experience depression for females may differ from males in clinically important ways. For instance, it may be the case the females are better treated for underlying anxiety and stress disorders in tandem with depression, while males may be treated strictly for depression.

With respect to important demographic features, results supported findings that personality tends to be a strong predictor of depression, anxiety, and stress. However, this relationship is slightly different for males compared to females with respect to stress. Feature

importance revealed that emotional stability, followed by family size, sexual orientation, agreeableness, and education level, all of which rated 50% or higher in importance compared to emotional stability, mattered more than openness to experience, conscientiousness, and extraversion.

While emotional stability has been found to exert the most influence over depression, anxiety, and stress for both females and males compared to other personality traits (Alizadeh et al., 2018; Uliaszek et al., 2010), the pathway to emotional stability as an important predictor looks different for females compared to males. These results suggest that for females, the pathway to emotional stability runs through extraversion and conscientiousness, with agreeableness playing a significant role along the way. By contrast, for males, emotional stability was the primary feature for stress, second for anxiety, and third for depression.

This is significant because females tend to rate higher in trait agreeableness and lower in trait emotional stability (Lippa, 2010; Weisberg et al., 2011), findings which have been replicated in this dataset (Allen, 2026). Thus, higher relative feature importance of these traits for females who tend rate higher in agreeableness and lower in emotional stability further sheds light into lifetime difference in clinical diagnoses of depression, anxiety, and stress. For males, a higher overall relative feature importance of emotional stability suggests emotional stability drives symptoms of mental health for males who suffer from moderate to severe depression, anxiety, and stress.

Thus, in answer to RQ1, machine models revealed potential gender differences in discriminant validity of DASS-21 depression subscales. In answer to RQ2, stress and anxiety subscales were more important in predicting clinically high levels of depression for females, compared to males where depression subscales were more important. Additionally, models revealed differences in feature importance of personality traits and other person level features between males and females.

While these findings do not depart from convention, they offer a sense of ordinality to the experiences and traits that most affect mental health for females and males separately. These results are supported by rigorously designed machine models including cost sensitivity (Feng et al., 2026; Sterner et al., 2025), population-level generalizable minimum sample splits (i.e., no less than 30 participants per node), and model honesty provisions (Athey et al., 2019).

Regarding RQ3, ordinal feature importance suggests learning the language clinicians use to diagnose depression, anxiety, and stress would help males and females in different ways. For females, placing a strong emphasis on language describing anxiety and stress, and how to parse them from depression would improve the granularity with which symptoms are described and managed. For males, similar attention should be paid to descriptors of anxiety and stress, but also on depression. Pairing this language with understanding how personality differently affects experiencing depression, anxiety, and stress would equip females and males alike with more tools to identify sources associated with declines in mental health.

## Limitations and Future Directions

The data was collected via convenience sampling. Convenience sampling is known to introduce bias such as social desirability and halo effect, especially in survey on desirable traits (Speklé & Widener, 2018). While attempts were made to ensure response integrity, such as removing participants who failed the attention check, and participants who listed their age as 100 years or more, the need for a large sample size for the project prohibited more stringent methodological constraints such as down sampling to mimic a random sample from the responses. It is also likely that the sample underrepresents low socio-economic individuals with limited access to devices capable of completing the survey.

Furthermore, the results are in terms of relative importance to the most important feature. No distinction between degree of importance was made with respect to effect size for females or males. Thus, no conclusion about how each feature meaningfully contributes to clinical levels of depression, anxiety, or stress may be drawn.

With respect to the models, unweighted models (i.e., AdaBoost, Gradient Boosting, or HistGradientBoost) tended to perform between 2-to-6% better than unweighted models for the demographic predictors of depression (Table S4; Fig. S4), stress (Table S5; Fig. S5), and anxiety (Table S6; Fig. S6). Although modestly, these models artificially increased ensemble accuracy, thus highlighting the importance of weighting in classification models. This is clear in the confusion matrices for demographic predictors, where positive cases were predicted with much higher accuracy than negative cases (Figs. S7, 8, 9).

Future research should consider both effect size and important feature splits. The impact of each feature, and how those features split for various individuals (e.g., heterosexual vs. homosexual) would yield more inclusive and granular results, thus serving a wider population. Given the age groups, future research would further parse results by age range to understand at which point emphasizing language learning about depression, anxiety, or stress independently would be more advantageous.

## Conclusion

This study extracted important features from machine model predictions of depression, anxiety, and stress for females and males to better understand which features best described these mental health indicators. Gender differences suggest that the more granularly females can distinguish between anxiety and stress, the more likely they will be able to parse these from depression. For females, a granular understanding of personality, especially on agreeableness and emotional stability, might enhance their ability to describe and understand the sources of depression, anxiety, and stress.

Males, by contrast, would benefit from a breadth of descriptors. The results demonstrated that for males, DASS-21 subscales aligned with each mental health indicator. Thus, a greater emphasis on vocabulary to describe each indicator would be beneficial for males.

In terms of demographic variables, males would benefit from understanding the interaction between emotional stability, family, and sexual orientation.

A clinically aligned vocabulary would provide deeper personal insight, and enhance communication with friends, family, significant others, and mental health experts. As one beloved neighbor said, anything that is human is mentionable, and anything mentionable is more manageable. At a time when Americans are reporting higher levels of depression, anxiety, and stress, there is great utility in having the language to mention it, simply that one can manage it.

**Table 1**

*DASS-21 Scales, Subscales, Associated Questions, and Root Vocabulary*

| Item no. | **SCALE**<br>*Subscale*<br>Item | Root Vocabulary |
|---|---|---|
| | **DEPRESSION** | |
| | *Anhedonia* | Anhedonia |
| 3 | I couldn't seem to experience any positive feeling at all | Positive feeling |
| | *Devaluation of life* | Devalue |
| 38 | I felt that life was meaningless | Meaningless |
| | *Dysphoria* | Dysphoria |
| 26 | I felt down-hearted and blue | Down-hearted, blue |
| | *Hopelessness* | Hopeless |
| 10 | I felt that I had nothing to look forward to | Look forward to |
| | *Inertia* | Inertia |
| 42 | I found it difficult to work up the initiative to do things | Initiative |
| | *Lack of interest / involvement* | Interest, involvement |
| 31 | I was unable to become enthusiastic about anything | Enthusiasm |
| | *Self-deprecation* | Self-deprecation |
| 17 | I felt that I wasn't worth much as a person | Worth |
| | **ANXIETY** | Anxiety |
| | *Autonomic arousal* | |
| 2 | I was aware of dryness of my mouth | Dry mouth |
| 4 | I experienced breathing difficulty (e.g., excessively rapid breathing, breathlessness in the absence of physical exertion) | Difficulty breathing |
| 25 | I was aware of the action of my heart in the absence of physical exertion (e.g., sense of heart rate increase, heart missing a beat) | Heartbeat |
| | *Situational anxiety* | |
| 40 | I was worried about situations in which I might panic and make a fool of myself | Panic, fool |
| | *Skeletal musculature effects* | |
| 41 | I experienced trembling (e.g., in the hands) | Trembling |
| | *Subjective experience of anxious affect* | |
| 20 | I felt scared without any good reason | Scared |
| 28 | I felt I was close to panic | Panic |
| | **STRESS** | Stress |
| | *Difficulty relaxing* | Relaxing |
| 8 | I found it difficult to relax | |
| 22 | I found it hard to wind down | Wind-down |
| | *Easily upset / agitated* | Upset, agitated |
| 11 | I found myself getting agitated | |
| | *Impatient* | Impatient |
| 35 | I was intolerant of anything that kept me from getting on with what I was doing | Intolerant |
| | *Irritable / over-reactive* | Irritable, over-react |
| 6 | I tended to over-react to situations | |
| 18 | I felt I was rather touchy | Touchy |
| | *Nervous arousal* | |
| 12 | I felt that I was using a lot of nervous energy | Nervous |

*Note:* Item numbers are based on the DASS-42 survey items included in the Answers to the Depression Anxiety Stress Scales data set. The items in this table are those related to the DASS-21 only.

**Table 2**

*Descriptive Statistics for Age Group, Religion, Depression, Anxiety, and Stress by Gender*

| | Female | | Male | |
|---|---|---|---|---|
| **Age Group (years)** | n | % | n | % |
| Adolescent (13-18) | 7815 | 25.92 | 2044 | 23.55 |
| Young adult (19-24) | 14355 | 47.60 | 3645 | 42 |
| Adult (25-44) | 6992 | 23.19 | 2367 | 27.27 |
| Middle aged (45-65) | 943 | 3.13 | 566 | 6.52 |
| Aged (66-79) | 48 | .16 | 53 | .61 |
| 80 and over | 3 | .01 | 4 | .05 |
| **Religion** | | | | |
| Religious | 4243 | 14.07 | 2745 | 31.63 |
| Not Religious | 25913 | 85.93 | 5934 | 68.37 |
| **Depression** | | | | |
| Low to mild | 9243 | 30.65 | 3035 | 34.97 |
| Moderate to severe | 20913 | 69.35 | 5644 | 65.03 |
| **Anxiety** | | | | |
| Low to mild | 8862 | 29.39 | 3636 | 41.89 |
| Moderate to severe | 21294 | 70.61 | 5043 | 58.11 |
| **Stress** | | | | |
| Low to mild | 11629 | 38.56 | 4567 | 52.62 |
| Moderate to severe | 18527 | 61.44 | 4112 | 47.38 |

**Table 3**

*DASS-21 Clinical Cutoff Scores for Emotional States*

| Severity Level | Depression | Anxiety | Stress |
|---|---|---|---|
| Normal | 0-9 | 0-7 | 0-14 |
| Mild | 10-13 | 8-9 | 15-18 |
| Moderate | 14-20 | 10-14 | 19-25 |
| Severe | 21-27 | 15-19 | 26-33 |
| Extremely Severe | ≥ 28 | ≥ 20 | ≥ 34 |

**Table 4**

*Descriptive Statistics for Age by Gender*

| Age (years) | n | *M* | *SD* | Min | Max |
|---|---|---|---|---|---|
| **Gender** | | | | | |
| Female | 30156 | 22.99 | 7.89 | 13 | 85 |
| Male | 8679 | 25.05 | 10.37 | 13 | 89 |

*Note:* $M$ = mean; $SD$ = standard deviation; Min = minimum; Max = maximum

**Table 5**

*Ensemble Model Results for DASS-21 Subscales as Predictors of Depression, Anxiety, and Stress*

| | Accuracy | Precision | Recall | F1 Score |
|---|---|---|---|---|
| **Depression** | | | | |
| Female | 1 | 1 | 1 | 1 |
| Male | .99 | .99 | .99 | .99 |
| **Anxiety** | | | | |
| Female | 1 | 1 | 1 | 1 |
| Male | 1 | 1 | 1 | 1 |
| **Stress** | | | | |
| Female | 1 | 1 | 1 | 1 |
| Male | 1 | 1 | 1 | 1 |

*Note:* All reported scores are based on weighted averages

## Table 6


*Ensemble Model Feature Important for DASS-21 Subscales*

| SCALE | Female | | | SCALE | Male | | |
|---|---|---|---|---|---|---|---|
| Subscale | Split Value | Importance Score | Subscale Importance | Subscale | Split Value | Importance Score | Relative Importance |
| **DEPRESSION** | | | | **DEPRESSION** | | | |
| Easily Upset | 7 | 476.68 | 1.00 | Inertia | 3 | 126.15 | 1.00 |
| Impatient | 1 | 456.00 | .96 | Hopelessness | 1 | 122.21 | .97 |
| Irritable / Over-reactive | 5 | 409.68 | .86 | Self-Deprecation | 3 | 112.87 | .89 |
| Autonomic Arousal | 1 | 401.67 | .84 | Lack of Interest / Involvement | 1 | 98.29 | .78 |
| Subjective Experience of Anxious Affect | 3 | 350.52 | .74 | Dysphoria | 3 | 92.04 | .73 |
| Dysphoria | 3 | 310.74 | .65 | Devaluation of Life | 1 | 85.42 | .68 |
| Nervous Arousal | 3 | 306.67 | .64 | Anhedonia | 1 | 83.79 | .66 |
| Situational Anxiety | 3 | 284.34 | .60 | Autonomic Arousal | 1 | 80.53 | .64 |
| Inertia | 1 | 281.19 | .59 | Irritable / Over-reactive | 5 | 77.21 | .61 |
| Lack of Interest / Involvement | 1 | 265.28 | .56 | Easily Upset | 5 | 73.89 | .59 |
| Anhedonia | 1 | 261.59 | .55 | Situational Anxiety | 3 | 60.18 | .48 |
| Difficulty Relaxing | 1 | 260.33 | .55 | Impatient | 3 | 60.17 | .48 |
| Self-Deprecation | 1 | 257.59 | .54 | Subjective Experience of Anxious Affect | 3 | 54.19 | .43 |
| Devaluation of Life | 1 | 237.61 | .50 | Difficulty Relaxing | 3 | 49.19 | .39 |
| Skeletal Musculature Effects | 1 | 235.67 | .49 | Nervous Arousal | 1 | 47.67 | .38 |
| Hopelessness | 1 | 221.77 | .47 | Skeletal Musculature Effects | 1 | 43.54 | .35 |
| **ANXIETY** | | | | **ANXIETY** | | | |
| Autonomic Arousal | 1 | 139.90 | 1.00 | Autonomic Arousal | 3 | 132.03 | 1.00 |
| Situational Anxiety | 1 | 133.21 | .95 | Subjective Experience of Anxious Affect | 1 | 119.69 | .91 |
| Subjective Experience of Anxious Affect | 1 | 125.64 | .90 | Situational Anxiety | 1 | 115.67 | .88 |
| Skeletal Musculature Effects | 1 | 104.03 | .74 | Skeletal Musculature Effects | 1 | 108.14 | .82 |
| Irritable / Over-reactive | 7 | 3.00 | .02 | Easily Upset | 3 | 6.87 | .05 |
| Easily Upset | 5 | 2.53 | .02 | Hopelessness | 1 | 5.67 | .04 |
| Nervous Arousal | 1 | 2.01 | .01 | Irritable / Over-reactive | 3 | 5.51 | .04 |
| Self-Deprecation | 5 | 1.67 | .01 | Nervous Arousal | 1 | 4.36 | .03 |
| Impatient | 1 | 1.50 | .01 | Inertia | 3 | 3.67 | .03 |
| Inertia | 1 | 1.50 | .01 | Devaluation of Life | 1 | 3.51 | .03 |
| Dysphoria | 5 | 1.34 | .01 | Impatient | 1 | 3.35 | .03 |
| Devaluation of Life | 1 | .33 | .00 | Difficulty Relaxing | 3 | 2.68 | .02 |
| Hopelessness | 1 | .33 | .00 | Lack of Interest / Involvement | 1 | 1.84 | .01 |
| Lack of Interest / Involvement | 1 | .33 | .00 | Self-Deprecation | 1 | 1.84 | .01 |
| Difficulty Relaxing | 5 | .00 | .00 | Anhedonia | 1 | 1.34 | .01 |
| Anhedonia | 1 | .00 | .00 | Dysphoria | 1 | 1.18 | .01 |
| **STRESS** | | | | **STRESS** | | | |
| Easily Upset | 5 | 1025.23 | 1.00 | Easily Upset | 7 | 578.75 | 1.00 |
| Irritable / Over-reactive | 3 | 923.04 | .90 | Irritable / Over-reactive | 5 | 530.19 | .92 |
| Impatient | 1 | 701.89 | .68 | Nervous Arousal | 3 | 455.32 | .79 |
| Nervous Arousal | 1 | 694.67 | .68 | Impatient | 3 | 444.86 | .77 |
| Difficulty Relaxing | 1 | 692.96 | .68 | Difficulty Relaxing | 3 | 406.43 | .70 |
| Autonomic Arousal | 5 | 111.68 | .11 | Autonomic Arousal | 1 | 155.84 | .27 |
| Subjective Experience of Anxious Affect | 3 | 83.00 | .08 | Subjective Experience of Anxious Affect | 1 | 90.69 | .16 |
| Situational Anxiety | 3 | 75.34 | .07 | Situational Anxiety | 3 | 88.17 | .15 |
| Hopelessness | 1 | 75.17 | .07 | Inertia | 3 | 82.19 | .14 |
| Inertia | 1 | 70.84 | .07 | Lack of Interest / Involvement | 1 | 77.84 | .13 |
| Skeletal Musculature Effects | 1 | 64.17 | .06 | Self-Deprecation | 1 | 77.02 | .13 |
| Lack of Interest / Involvement | 1 | 60.33 | .06 | Hopelessness | 1 | 64.67 | .11 |
| Anhedonia | 1 | 51.67 | .05 | Dysphoria | 1 | 61.52 | .11 |
| Devaluation of Life | 1 | 48.67 | .05 | Skeletal Musculature Effects | 5 | 58.84 | .10 |
| Dysphoria | 5 | 45.17 | .04 | Devaluation of Life | 5 | 55.67 | .10 |
| Self-Deprecation | 1 | 45.00 | .04 | Anhedonia | 3 | 55.17 | .10 |

*Note:* Relative importance was calculated by dividing each importance score by the highest score

**Figure 1**


*Top DASS-21 Subscale Predictors of Anxiety and Stress*




This graphic shows relative feature importance for females and males with respect to anxiety (left) and stress (right). Subscales for anxiety and stress predictably align with their respective scales, suggesting the measures have strong discriminant validity. Feature importance for anxiety and stress are similar for both males and females.

**Figure 2**

*Top DASS-21 Subscale Predictors of Depression*




The chart on the left shows relative feature importance scores in descending order for females, whereas the chart on the right shows scores in descending order for males. This highlights how important features for females and males differ. For example, features more predictive of anxiety and stress drive depression scores for females, while the associated subscales for depression best predict depression for males.

**Table 7**

*Ensemble Model Results for Demographic Variables as Predictors of Depression, Anxiety, and Stress*

| | Accuracy | Precision | Recall | F1 Score |
|---|---|---|---|---|
| **Depression** | | | | |
| Female | .76 | .76 | .76 | .76 |
| Male | .76 | .76 | .76 | .76 |
| **Anxiety** | | | | |
| Female | .74 | .73 | .74 | .74 |
| Male | .72 | .72 | .72 | .72 |
| **Stress** | | | | |
| Female | .75 | .75 | .75 | .75 |
| Male | .74 | .74 | .74 | .74 |

*Note:* All reported scores are based on weighted averages

## Table 8


*Ensemble Model Feature Important for Demographic Variables*

| SCALE | Female | | | SCALE | Male | | |
|---|---|---|---|---|---|---|---|
| Demographic Variable | Split Value | Importance Score | Relative Importance | Demographic Variable | Split Value | Importance Score | Relative Importance |
| **DEPRESSION** | | | | **DEPRESSION** | | | |
| Extraversion | 3.83 | 314.29 | 1.00 | Extraversion | 3.17 | 167.70 | 1.00 |
| Conscientiousness | 4.75 | 285.75 | .91 | Conscientiousness | 4.25 | 148.34 | .88 |
| Agreeableness | 4.25 | 273.47 | .87 | Emotional Stability | 3.25 | 14.42 | .84 |
| Emotional Stability | 3.75 | 264.92 | .84 | Openness to Experience | 4.25 | 124.87 | .74 |
| Family Size | 3.50 | 261.03 | .83 | Family Size | 1.50 | 119.41 | .71 |
| Openness to Experience | 4.75 | 243.14 | .77 | Agreeableness | 4.25 | 117.24 | .70 |
| Education Level | 2.50 | 167.32 | .53 | Education Level | 2.50 | 77.85 | .46 |
| Sexual Orientation | 1.50 | 142.99 | .45 | Age Group | 1.50 | 69.30 | .41 |
| Urban vs. Rural Childhood | 2.50 | 134.41 | .43 | Urban vs. Rural Childhood | 1.50 | 66.76 | .40 |
| Age Group | 1.50 | 133.51 | .42 | Race | 5.50 | 64.19 | .38 |
| Race | 5.50 | 118.56 | .38 | Sexual Orientation | .50 | 63.31 | .38 |
| Voted | 1.50 | 72.87 | .23 | Voted | 1.50 | 42.12 | .25 |
| Married | 1.50 | 60.77 | .19 | Religious | .50 | 36.72 | .22 |
| Religious | .50 | 44.32 | .14 | Married | 1.50 | 29.09 | .17 |
| **ANXIETY** | | | | **ANXIETY** | | | |
| Extraversion | 3.50 | 635.39 | 1.00 | Extraversion | 2.83 | 75.10 | 1.00 |
| Conscientiousness | 4.25 | 580.54 | .91 | Emotional Stability | 2.75 | 64.18 | .85 |
| Agreeableness | 4.25 | 541.78 | .85 | Conscientiousness | 4.25 | 59.10 | .79 |
| Openness to Experience | 5.25 | 528.29 | .83 | Agreeableness | 4.75 | 48.92 | .65 |
| Family Size | 2.50 | 51.73 | .80 | Family Size | 2.50 | 48.76 | .65 |
| Emotional Stability | 2.25 | 509.79 | .80 | Openness to Experience | 4.25 | 47.72 | .64 |
| Education Level | 2.50 | 318.72 | .50 | Age Group | .50 | 31.42 | .42 |
| Sexual Orientation | .50 | 285.20 | .45 | Race | 5.50 | 30.12 | .40 |
| Urban vs. Rural Childhood | 2.50 | 273.23 | .43 | Education Level | 2.50 | 27.89 | .37 |
| Age Group | .50 | 244.44 | .38 | Urban vs. Rural Childhood | 2.50 | 22.85 | .30 |
| Race | 5.50 | 223.45 | .35 | Sexual Orientation | 1.50 | 22.30 | .30 |
| Voted | 1.50 | 158.85 | .25 | Religious | .50 | 15.15 | .20 |
| Married | 1.50 | 106.95 | .17 | Voted | 1.50 | 13.86 | .18 |
| Religious | .50 | 99.98 | .16 | Married | 1.50 | 9.97 | .13 |
| **STRESS** | | | | **STRESS** | | | |
| Extraversion | 3.50 | 635.39 | 1.00 | Emotional Stability | 2.75 | 97.89 | 1.00 |
| Conscientiousness | 4.25 | 580.54 | .91 | Family Size | 2.50 | 79.91 | .82 |
| Agreeableness | 4.25 | 541.78 | .85 | Sexual Orientation | .50 | 56.32 | .58 |
| Openness to Experience | 5.25 | 528.29 | .83 | Agreeableness | 4.25 | 53.83 | .55 |
| Family Size | 2.50 | 510.73 | .80 | Education Level | 2.50 | 48.02 | .49 |
| Emotional Stability | 2.25 | 509.79 | .80 | Age Group | .50 | 38.69 | .40 |
| Education Level | 2.50 | 318.72 | .50 | Openness to Experience | 5.75 | 32.02 | .33 |
| Sexual Orientation | .50 | 285.20 | .45 | Conscientiousness | 4.25 | 30.17 | .31 |
| Urban vs. Rural Childhood | 2.50 | 273.23 | .43 | Extraversion | 3.50 | 19.10 | .20 |
| Age Group | .50 | 244.44 | .38 | Voted | 1.50 | 15.02 | .15 |
| Race | 5.50 | 223.45 | .35 | Religious | .50 | 14.25 | .15 |
| Voted | 1.50 | 158.85 | .25 | Urban vs. Rural Childhood | 2.50 | 14.04 | .14 |
| Married | 1.50 | 106.95 | .17 | Race | 5.50 | 11.37 | .12 |
| Religious | .50 | 99.98 | .16 | Married | 1.50 | 6.70 | .07 |

*Note:* Relative importance was calculated by dividing each importance score by the highest score

**Figure 3**


*Top Demographic Predictors of Depression and Anxiety*




This graphic shows relative feature importance for females and males with respect to depression (left) and anxiety (right). Features are arranged in descending order by importance for females. While demographic feature importance is relatively similar for females and males, emotional stability is more important for males than females for predicting anxiety.

**Figure 4**

*Top Demographic Predictors of Stress*




The chart on the left shows relative feature importance scores in descending order for females, whereas the chart on the right shows importance scores in descending order for males. This highlights how important features for females and males differ. For example, personality features drive stress predictors for females, while non-personality features like family size, sexual orientation, and education level are of greater importance for males.

References

Adu, P., Popoola, T., Iqbal, N., Medvedev, O. N., & Simpson, C. R. (2025). Validating the depression anxiety stress scales (DASS-21) across Germany, Ghana, India, and New Zealand using Rasch methodology. *Journal of Affective Disorders*, *383*, 363–373. https://doi.org/10.1016/j.jad.2025.04.099

Al Abdi, R., AlKaabi, S., Elsifi, S., & Yousaf, J. (2026). Mental stress detection using physiological sensors and artificial intelligence: A review. *Sensors*, *26*(5), 1616. https://doi.org/10.3390/s26051616

Alizadeh, Z., Feizi, A., Rejali, M., Afshar, H., Keshteli, A. H., & Adibi, P. (2018). The Predictive Value of Personality Traits for Psychological Problems (Stress, Anxiety and Depression): Results from a Large Population Based Study. *Journal of Epidemiology and Global Health*, *8*(3), 124–133. https://doi.org/10.2991/j.jegh.2017.11.003

Allen, B. (2026). *Gender differences or schmifferences: Testing replication with open source data* [Unpublished manuscript]. School of Information and Computing Science, Willamette University.

American Psychiatric Association. (2026, May 12). *Annual mental health poll finds Americans anxious about current events, personal finances and emerging technology* [Press release]. Psychiatry.Org. https://www.psychiatry.org/News-room/News-Releases/2026-Annual-Mental-Health-Poll

Antony, M. M., Bieling, P. J., Cox, B. J., Enns, M. W., & Swinson, R. P. (1998). Psychometric properties of the 42-item and 21-item versions of the Depression Anxiety Stress Scales in clinical groups and a community sample. *Psychological Assessment*, *10*(2), 176–181. https://doi.org/10.1037/1040-3590.10.2.176

Athey, S., Tibshirani, J., & Wager, S. (2019). Generalized random forests. *The Annals of Statistics*, *47*(2), 1148–1178.

Barrett, L. F. (2004). Feelings or words? Understanding the content in self-report ratings of experienced emotion. *Journal of Personality and Social Psychology*, *87*(2), 266. https://doi.org/10.1037/0022-3514.87.2.266

Barrett, L. F., Gross, J., Christensen, T. C., & Benvenuto, M. (2001). Knowing what you're feeling and knowing what to do about it: Mapping the relation between emotion differentiation and emotion regulation. *Cognition & Emotion*, *15*(6), 713–724. https://doi.org/10.1080/02699930143000239

Bradberry, T., & Greaves, J. (2009). *Emotional intelligence 2.0*. TalentSmart.

Breiman, L. (2001). Random Forests. *Machine Learning*, *45*(1), 5–32. https://doi.org/10.1023/A:1010933404324

Cavanagh, A., Wilson, C. J., Kavanagh, D. J., & Caputi, P. (2017). Differences in the expression of symptoms in men versus women with depression: A systematic review and meta-analysis. *Harvard Review of Psychiatry*, *25*(1), 29. https://doi.org/10.1097/HRP.0000000000000128

Chen, T., & Guestrin, C. (2016). XGBoost: A scalable tree boosting system. *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, KDD '16*, 785–794. https://doi.org/10.1145/2939672.2939785

Dixon-Gordon, K. L., Chapman, A. L., Weiss, N. H., & Rosenthal, M. Z. (2014). A preliminary examination of the role of emotion differentiation in the relationship between borderline personality and urges for maladaptive behaviors. *Journal of Psychopathology and Behavioral Assessment*, *36*(4), 616–625. https://doi.org/10.1007/s10862-014-9423-4

Donner, N. C., & Lowry, C. A. (2013). Sex differences in anxiety and emotional behavior. *Pflügers Archiv - European Journal of Physiology*, *465*(5), 601–626. https://doi.org/10.1007/s00424-013-1271-7

Dwight, A. R., Briesch, A. M., Hoffman, J. A., & Rutt, C. (2024). Dependability and usability of the DASS-21 as a progress-monitoring tool for youth. *Journal of Psychoeducational Assessment*, *42*(4), 453–466. https://doi.org/10.1177/07342829241227758

Farhane-Medina, N. Z., Luque, B., Tabernero, C., & Castillo-Mayén, R. (2022). Factors associated with gender and sex differences in anxiety prevalence and comorbidity: A systematic review. *Science Progress*, *105*(4), 00368504221135469. https://doi.org/10.1177/00368504221135469

Feng, Y., Du, H., Song, J., Sun, Y., Wang, Y., & Joel, A. (2026). Advancing Psychological Research With Random Forests: A Review of Methods, Tools, and Applications. *Advances in Methods and Practices in Psychological Science*, *9*(2), 25152459251404358. https://doi.org/10.1177/25152459251404358

Fioroni, S., & Foy, D. (2024, April 15). *Americans sleeping less, more stressed*. Gallup.Com. https://news.gallup.com/poll/642704/americans-sleeping-less-stressed.aspx

Freund, Y., & Schapire, R. E. (1997). A decision-theoretic generalization of on-line learning and an application to boosting. *Journal of Computer and System Sciences*, *55*(1), 119–139. https://doi.org/10.1006/jcss.1997.1504

Friedman, J. H. (2001). Greedy function approximation: A gradient boosting machine. *The Annals of Statistics*, *29*(5), 1189–1232. https://doi.org/10.1214/aos/1013203451

Geifman, N., Cohen, R., & Rubin, E. (2013). Redefining meaningful age groups in the context of disease. *Age*, *35*(6), 2357–2366. https://doi.org/10.1007/s11357-013-9510-6

Gomez, R., Summers, M., Summers, A., Wolf, A., & Summers, J. (2014). Depression Anxiety Stress Scales-21: Measurement and structural invariance across ratings of men and women. *Assessment*, *21*(4), 418–426. https://doi.org/10.1177/1073191113514106

Gosling, S. D., Rentfrow, P. J., & Swann, W. B. (2003). A very brief measure of the Big-Five personality domains. *Journal of Research in Personality*, *37*(6), 504–528. https://doi.org/10.1016/S0092-6566(03)00046-1

Graves, B. S., Hall, M. E., Dias-Karch, C., Haischer, M. H., & Apter, C. (2021). Gender differences in perceived stress and coping among college students. *PLOS ONE*, *16*(8), e0255634. https://doi.org/10.1371/journal.pone.0255634

Gupta, P., & Pandit, A. (2026). Leveraging voting techniques to predict mental stress: Exploring the efficacy of hard and soft voting models. *National Research Journal of Information Technology & Information Science*, (13), 622–628.

Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van Kerkwijk, M. H., Brett, M., Haldane, A., del Río, J. F., Wiebe, M., Peterson, P., … Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, *585*(7825), 357–362. https://doi.org/10.1038/s41586-020-2649-2

Henry, J. D., & Crawford, J. R. (2005). The short-form version of the Depression Anxiety Stress Scales (DASS-21): Construct validity and normative data in a large non-clinical sample. *British Journal of Clinical Psychology*, *44*(2), 227–239. https://doi.org/10.1348/014466505X29657

Hunter, J. D. (2007). Matplotlib: A 2D Graphics Environment. *Computing in Science & Engineering*, *9*(3), 90–95. https://doi.org/10.1109/MCSE.2007.55

Jain, B., Chitaliya, H., Gopani, M., Mehta, N., Gawde, A., & Dhekane, S. (2025). Enhancing mental health disorder detection: A hybrid classifier system using soft voting and ensemble methods. *IJFMR - International Journal For Multidisciplinary Research*, *7*(1). https://doi.org/10.36948/ijfmr.2025.v07i01.35963

Johnson, D. P., & Whisman, M. A. (2013). Gender differences in rumination: A meta-analysis. *Personality and Individual Differences*, *55*(4), 367–374. https://doi.org/10.1016/j.paid.2013.03.019

Jones, J. (2025). *U.S. mental health ratings continue to worsen*. Gallup, Inc. https://news.gallup.com/poll/700079/mental-health-ratings-continue-worsen.aspx

Kalokerinos, E. K., Erbas, Y., Ceulemans, E., & Kuppens, P. (2019). Differentiate to regulate: Low negative emotion differentiation is associated with ineffective use but not selection of emotion-regulation strategies. *Psychological Science*, *30*(6), 863–879. https://doi.org/10.1177/0956797619838763

Kashdan, T. B., Barrett, L. F., & McKnight, P. E. (2015). Unpacking emotion differentiation: Transforming unpleasant experience by perceiving distinctions in negativity. *Current Directions in Psychological Science*, *24*(1), 10–16. https://doi.org/10.1177/0963721414550708

Ke, G., Meng, Q., Finley, T., Wang, T., Chen, W., Ma, W., Ye, Q., & Liu, T.-Y. (2017). LightGBM: A highly efficient gradient boosting decision tree. *Proceedings of the 31st International Conference on Neural Information Processing Systems, NIPS'17*, 3149–3157. https://dl.acm.org/doi/10.5555/3294996.3295074

Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., & Walters, E. E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, *62*(6), 593–602. https://doi.org/10.1001/archpsyc.62.6.593

Kircanski, K., Lieberman, M. D., & Craske, M. G. (2012). Feelings into words: Contributions of language to exposure therapy. *Psychological Science*, *23*(10), 1086–1091. https://doi.org/10.1177/0956797612443830

Kopp, D., Maudet, B., Sun-Hosoya, L., Bennett, K. P., & Paris-Saclay, U. (2025). Encoding Matters: Impact of Categorical Variable Encoding on Performance and Bias. *ESANN 2025 Proceedings, European Symposium on Artificial Neural Networks, Computational Intelligence and Machine Learning*.

Kuehner, C. (2017). Why is depression more common among women than among men? *The Lancet Psychiatry*, *4*(2), 146–158. https://doi.org/10.1016/S2215-0366(16)30263-2

Kumar, P., Garg, S., & Garg, A. (2020). Assessment of anxiety, depression and stress using machine learning models. *Procedia Computer Science, Third International Conference on Computing and Network Communications (CoCoNet'19)*, *171*, 1989–1998. https://doi.org/10.1016/j.procs.2020.04.213

Levkovich, I., Yatzkar, U., & Shenaar-Golan, V. (2025). Age and gender differences in emotional and behavioral functioning among youth referred to a psychiatric outpatient clinic at a public hospital. *Children*, *12*(6), 683. https://doi.org/10.3390/children12060683

Levy-Gigi, E., & Shamay-Tsoory, S. (2022). Affect labeling: The role of timing and intensity. *PLOS ONE*, *17*(12), e0279303. https://doi.org/10.1371/journal.pone.0279303

Li, G., Li, Y., Zhang, Z., Chen, Y., Li, B., Hao, D., Yang, L., Yang, Y., Li, X., & Li, C.-S. R. (2023). Sex differences in externalizing and internalizing traits and ventral striatal responses to monetary loss. *Journal of Psychiatric Research*, *162*, 11–20. https://doi.org/10.1016/j.jpsychires.2023.04.013

Lippa, R. A. (2010). Gender differences in personality and interests: When, where, and why?: Gender differences in personality and interests. *Social and Personality Psychology Compass*, *4*(11), 1098–1110. https://doi.org/10.1111/j.1751-9004.2010.00320.x

Liu, Y., María-Itatí, P., Taha, B., Toledo, C., Ouyang, Y., Zhongzheng, L., Wang, Z., Toledo, F., Zeng, H., & María-Trinidad, H. (2025). Machine learning, physiological signals, and emotional stress/anxiety: Pitfalls and challenges. *Applied Sciences*, *15*(21), 11777–11806. https://doi.org/10.3390/app152111777

Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states: Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behaviour Research and Therapy*, *33*(3), 335–343. https://doi.org/10.1016/0005-7967(94)00075-U

Lovibond, S. H., & Lovibond, P. F. (1995). *Manual for the Depression Anxiety Stress Scales* (2nd ed.). Psychology Foundation. https://www2.psy.unsw.edu.au/dass/

Martin, L. A., Neighbors, H. W., & Griffith, D. M. (2013). The experience of symptoms of depression in men vs women: Analysis of the national comorbidity survey replication. *JAMA Psychiatry*, *70*(10), 1100–1106. https://doi.org/10.1001/jamapsychiatry.2013.1985

Matud, M. P. (2004). Gender differences in stress and coping styles. *Personality and Individual Differences*, *37*(7), 1401–1415. https://doi.org/10.1016/j.paid.2004.01.010

Mayer, J. D., & Salovey, P. (1995). Emotional intelligence and the construction and regulation of feelings. *Applied and Preventive Psychology*, *4*(3), 197–208. https://doi.org/10.1016/S0962-1849(05)80058-7

McLean, C. P., Asnaani, A., Litz, B. T., & Hofmann, S. G. (2011). Gender differences in anxiety disorders: Prevalence, course of illness, comorbidity and burden of illness. *Journal of Psychiatric Research*, *45*(8), 1027–1035. https://doi.org/10.1016/j.jpsychires.2011.03.006

Microsoft. (2026). *Visual Studio Code* (Version 1.128.0) [Windows]. Microsoft. https://code.visualstudio.com

Nembrini, S., König, I. R., & Wright, M. N. (2018). The revival of the Gini importance? *Bioinformatics*, *34*(21), 3711–3718. https://doi.org/10.1093/bioinformatics/bty373

Nook, E. C., Stavish, C. M., Sasse, S. F., Lambert, H. K., Mair, P., McLaughlin, K. A., & Somerville, L. H. (2020). Charting the development of emotion comprehension and abstraction from childhood to adulthood using observer-rated and linguistic measures. *Emotion (Washington, D.C.)*, *20*(5), 773–792. https://doi.org/10.1037/emo0000609

Open Psychometrics. (2019). *Answers to the depression anxiety stress scales* [Dataset]. https://openpsychometrics.org/_rawdata/

Open-Source Psychometrics Project. (2019, September 17). *About the Open-Source Psychometrics Project*. https://openpsychometrics.org/about/

Pandas via NumFOCUS, Inc. (2023). *Categorical data*. Pandas 2.1.4 Documentation. https://pandas.pydata.org/pandas-docs/version/2.1/user_guide/categorical.html

Pavani, S., Gulhare, K. K., Handa, R., & Kushwaha, S. (2024). "Enhancing mental health assessments: The role of voting classifiers in evaluating depression's impact on quality of life." *International Journal of Communication Networks and Information Security (IJCNIS)*, *16*(4), 727–735.

Pedersen, T. L. (2025). *patchwork: The composer of plots* (Version 1.3.2) [Computer software]. https://cran.r-project.org/web/packages/patchwork/index.html

Pedregosa, F., Pedregosa, F., Varoquaux, G., Varoquaux, G., Org, N., Gramfort, A., Gramfort, A., Michel, V., Michel, V., Fr, L., Thirion, B., Thirion, B., Grisel, O., Grisel, O., Blondel, M., Prettenhofer, P., Prettenhofer, P., Weiss, R., Dubourg, V., … Cournapeau, D. (n.d.). Scikit-learn: Machine Learning in Python. *MACHINE LEARNING IN PYTHON*.

Posit team. (2026). *RStudio: Integrated Development Environment for R* (Version 2026.01.1.403) [Computer software]. Posit Software, PBC. http://www.posit.co/

Prokhorenkova, L., Gusev, G., Vorobev, A., Dorogush, A. V., & Gulin, A. (2018). CatBoost: Unbiased boosting with categorical features. *Proceedings of the 32nd International Conference on Neural Information Processing Systems, NIPS'18*, 6639–6649. https://dl.acm.org/doi/10.5555/3327757.3327770

Python Software Foundation. (2025, August 14). *Python 3.13.7*. Python.Org. https://www.python.org/downloads/release/python-3137/

R Core Team. (2026). *R: A Language and Environment for Statistical Computing* (Version 4.6.0) [Computer software]. R Foundation for Statistical Computing. https://www.R-project.org/

Robinson, D., Hayes, A., Couch, S., Hvitfeldt, E., Posit Software, PBC, Patil, I., Chiu, D., Gomez, M., Demeshev, B., Menne, D., Nutter, B., Johnston, L., Bolker, B., Briatte, F., Arnold, J., Gabry, J., Selzer, L., Simpson, G., … Reinhart, A. (2026). *broom: Convert*

*Statistical Objects into Tidy Tibbles* (Version 1.0.13) [Computer software]. https://cran.r-project.org/web/packages/broom/index.html

Rogers, F. (1994). *You are special: Words of wisdom for all of us from a beloved neighbor*. Viking Press.

Salk, R. H., Hyde, J. S., & Abramson, L. Y. (2017). Gender Differences in Depression in Representative National Samples: Meta-Analyses of Diagnoses and Symptoms. *Psychological Bulletin*, *143*(8), 783–822. https://doi.org/10.1037/bul0000102

Salovey, P., & Mayer, J. D. (1990). Emotional intelligence. *Imagination, Cognition and Personality*, *9*(3), 185–211. https://doi.org/10.2190/DUGG-P24E-52WK-6CDG

Sloan, D. M., & Sandt, A. R. (2006). Gender differences in depression. *Women's Health*, *2*(3), 425–434. https://doi.org/10.2217/17455057.2.3.425

Smidt, K. E., & Suvak, M. K. (2015). A brief, but nuanced, review of emotional granularity and emotion differentiation research. *Current Opinion in Psychology, Emotion Regulation*, *3*, 48–51. https://doi.org/10.1016/j.copsyc.2015.02.007

Soria-Reyes, L. M., Alarcón, R., Cerezo, M. V., & Blanca, M. J. (2024). Psychometric properties of the Depression Anxiety Stress Scales (DASS-21) in women with breast cancer. *Scientific Reports*, *14*, 20473. https://doi.org/10.1038/s41598-024-68814-9

Speklé, R. F., & Widener, S. K. (2018). Challenging issues in survey research: Discussion and suggestions. *Journal of Management Accounting Research*, *30*(2), 3–21. https://doi.org/10.2308/jmar-51860

Sterner, P., Goretzko, D., & Pargent, F. (2025). Everything has its price: Foundations of cost-sensitive machine learning and its application in psychology. *Psychological Methods*, *30*(1), 112–127. https://doi.org/10.1037/met0000586

Strobl, C., Boulesteix, A.-L., Zeileis, A., & Hothorn, T. (2007). Bias in random forest variable importance measures: Illustrations, sources and a solution. *BMC Bioinformatics*, *8*(1), 25. https://doi.org/10.1186/1471-2105-8-25

Sujon, K. M., Hassan, R., Choi, K., & Samad, M. A. (2025). Accuracy, precision, recall, f1-score, or MCC? Empirical evidence from advanced statistics, ML, and XAI for evaluating business predictive models. *Journal of Big Data*, *12*(1), 268. https://doi.org/10.1186/s40537-025-01313-4

Taskynbayeva, M., & Gutoreva, A. (2025). Machine learning approaches to anxiety detection: Trends, model evaluation, and future directions. *Frontiers in Artificial Intelligence*, *8*. https://doi.org/10.3389/frai.2025.1630047

The Pandas Development Team. (2020). *Pandas* (Version 2.3.3) [Computer software]. Zenodo. https://doi.org/10.5281/zenodo.17229934

Tugade, M. M., Fredrickson, B. L., & Barrett, L. F. (2004). Psychological resilience and positive emotional granularity: Examining the benefits of positive emotions on coping and health. *Journal of Personality*, *72*(6), 1161–1190. https://doi.org/10.1111/j.1467-6494.2004.00294.x

Uliaszek, A. A., Zinbarg, R. E., Mineka, S., Craske, M. G., Sutton, J. M., Griffith, J. W., Rose, R., Waters, A., & Hammen, C. (2010). The role of neuroticism and extraversion in the stress–anxiety and stress–depression relationships. *Anxiety, Stress, & Coping*, *23*(4), 363–381. https://doi.org/10.1080/10615800903377264

Waskom, M. L. (2021). Seaborn: Statistical data visualization. *Journal of Open Source Software*, *6*(60), 3021. https://doi.org/10.21105/joss.03021

Weisberg, Y. J., DeYoung, C. G., & Hirsh, J. B. (2011). Gender differences in personality across the ten aspects of the Big Five. *Frontiers in Psychology*, *2*. https://doi.org/10.3389/fpsyg.2011.00178

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., … Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, *4*(43), 1686. https://doi.org/10.21105/joss.01686

Witters, D. (2026, April 21). *U.S. depression rate remains elevated*. Gallup, Inc. Gallup.Com. https://news.gallup.com/poll/708221/depression-rate-remains-elevated.aspx

Yesudas, A. (2023). *A Machine Learning Framework to Predict Depression, Anxiety and Stress* [Masters, Dublin, National College of Ireland]. https://norma.ncirl.ie/6346/

Yue, P., An, Z., & Lin, R. (2025). How do “words poorly expressed emotion” affect mental health? The mediating role of affect labelling effect. *Cognition and Emotion*, *39*(7), 1703–1718. https://doi.org/10.1080/02699931.2024.2362377

Zhu, W., Qiu, R., & Fu, Y. (2024). Comparative study on the performance of categorical variable encoders in classification and regression tasks. *arXiv*. https://doi.org/10.48550/ARXIV.2401.09682