


# Predicting plasticity outcomes based on LFP magnitude

Sebastiano Bariselli &  
Bridget Matikainen-Ankney  
BIOF309, Spring 2018

# Background

## Neural plasticity:

*The ability of connections between neurons to change*



# Question

*Can we predict plasticity outcomes based on local field potential magnitude during PRE period?*

# Project goals


- Import data
- Clean data (outliers)
- Data visualization
- Use prediction model

# IMPORT DATA

```
import numpy as np
import pandas as pd
import matplotlib.pyplot as plt
import scipy.stats as stats
from sklearn.linear_model import LinearRegression
from sklearn.metrics import mean_squared_error
from sklearn.model_selection import train_test_split

# Retrieve '.csv' files containing electrophysiology data in TBS and Cocaine conditions.
LFP_data_TBS = pd.read_csv('/Users/barisellisebastiano/Desktop/Excel_Practice4.csv')
LFP_data_COC = pd.read_csv('/Users/barisellisebastiano/Desktop/Excel_practice_Cocaine.csv')

# Plot a boxplot for the data to observe data points in 4 distributions: PreTBS, PostTBS, PreCOC, PostCOC
data1 = LFP_data_TBS['PRE'].values.reshape(-1, 1)
data2 = LFP_data_TBS['POST'].values.reshape(-1, 1)
data3 = LFP_data_COC['PRE'].values.reshape(-1, 1)
data4 = LFP_data_COC['POST'].values.reshape(-1, 1)
data = [data1, data2, data3, data4]
plt.boxplot(data)
plt.xticks([1, 2, 3, 4], ['PreTBS', 'PostTBS', 'PreCOC', 'PostCOC'])
plt.ylabel('LFP magnitude (mV)')
plt.xlabel('Plasticity protocol')
plt.show()
```



# CLEANING THE DATAFRAME

```
# Calculate mean and standard deviation for preTBS, postTBS, preCocaine and postCocaine and
LFP_data_TBS['MeanPre'] = LFP_data_TBS['PRE'].mean()
LFP_data_TBS['StDevPre'] = LFP_data_TBS['PRE'].std()
LFP_data_TBS['MeanPost'] = LFP_data_TBS['POST'].mean()
LFP_data_TBS['StDevPost'] = LFP_data_TBS['POST'].std()
LFP_data_COC['MeanPre'] = LFP_data_COC['PRE'].mean()
LFP_data_COC['StDevPre'] = LFP_data_COC['PRE'].std()
LFP_data_COC['MeanPost'] = LFP_data_COC['POST'].mean()
LFP_data_COC['StDevPost'] = LFP_data_COC['POST'].std()


# Write a function to detect 3 standard deviation (SD) outliers from 'PRE' column of a DataFrame
def remove_outliers_fromPRE(row):
    Pre = row['PRE']
    if Pre < (row['MeanPre'] - (3 * row['StDevPre'])) or Pre > (row['MeanPre'] + (3 * row['StDevPre'])):
        return NaN
    else:
        return Pre

# Write a function to detect 3 standard deviation (SD) outliers from 'POST' column of a DataFrame
def remove_outliers_fromPOST(row):
    Post = row['POST']
    if Post < (row['MeanPost'] - (3 * row['StDevPost'])) or Post > (row['MeanPost'] + (3 * row['StDevPost'])):
        return NaN
    else:
        return Post

# Apply a function to TBS and Cocaine dataframe create columns
LFP_data_TBS['cPRE'] = LFP_data_TBS.apply(remove_outliers_fromPRE, axis=1)
LFP_data_COC['cPRE'] = LFP_data_COC.apply(remove_outliers_fromPRE, axis=1)
LFP_data_TBS['cPOST'] = LFP_data_TBS.apply(remove_outliers_fromPOST, axis=1)
LFP_data_COC['cPOST'] = LFP_data_COC.apply(remove_outliers_fromPOST, axis=1)


# Add a new column 'cCHANGE' to the TBS Dataframe that contains the change in LFP magnitude
for row_index, row in LFP_data_TBS.iterrows():
    LFP_data_TBS['cCHANGE'] = LFP_data_TBS['cPOST'] / LFP_data_TBS['cPRE']

# Drop the rows in TBS DataFrame that contain 'NaN'
LFP_data_TBS_cleaned = LFP_data_TBS.dropna(axis=0, h
```



# Is there a plasticity?


```
# Execute a paired t-test to assess changes between pre- and post- for TBS and Cocaine conditions.  
print('paired t-test for TBS = ' + str(stats.ttest_rel(LFP_data_TBS_cleaned['cPRE'], LFP_data_TBS_cleaned['cPOST'])))  
print('paired t-test for COCAINE = ' + str(stats.ttest_rel(LFP_data_COC_cleaned['cPRE'], LFP_data_COC_cleaned['cPOST'])))
```



```
paired t-test for TBS = Ttest_relResult(statistic=-5.7939485436530358, pvalue=1.8844506960029739e-08)  
paired t-test for COCAINE = Ttest_relResult(statistic=-9.7858542513832418, pvalue=8.5423497024256997e-18)
```

# Regression analysis

- Regression: % Change = [ 6.55694474 ] \* Pre-stim response amplitude + 0.954975708918
- R squared = 0.471091816409





# Conclusions

- 1) LFP magnitude at PRE does predict plasticity outcomes given a linear models ( $R^2 = 0.47$ ).
- 2) This code will be used to organize and analyze neural data.