

# Even- $n$ Torus Knots in CDFD: The Exclusion of $T(2, 2)$ , the Extension to $T(2, 2m)$ , and the Anti-Phase Mass Scaling Law

Steve Bico Mujjabi, MD\*  
VuraLabs, Kampala, Uganda  
Independent researcher  
ORCID: 0009-0001-0556-5516

<https://github.com/bampita-bico/CDFD-Part-I-Release>

July 2026

## Abstract

Paper VIII left four open problems, of which this paper resolves three: the physical interpretation of the universal anti-phase modes, the extension of the CDFD torus-knot hierarchy to even family numbers, and the mass scaling of the lightest modes.

The central structural result is the Fourier exclusion theorem. The universal mass sum rule depends on a Fourier identity that holds exactly for family numbers three and higher but fails for family number two. The Hopf fibration is therefore excluded from the CDFD hierarchy by algebraic necessity rather than by an adjustable physical preference.

For even family numbers of four and greater, the Fourier identity holds and the machinery of Papers VI-VIII extends without modification. The paper computes self-consistency solutions, amplitude splits, mode spectra, and the seven-fourths mass hierarchy for family numbers four, six, eight, and ten. The amplitude-split rule from Paper VIII applies to even families with one identified exception at family number eight.

The anti-phase modes are identified as boundary modes. Their masses are set by how close the self-consistency orientation places the anti-phase Fourier modes to an amplitude zero, which depends on discrete number-theoretic alignment rather than on a simple power law. This explains why the two lightest modes are not Goldstone bosons and have no simple topology-independent mass formula. A complete mode-mass census for family numbers three through thirteen is assembled as a unified hierarchy table.

Across the series, the public CDFD Runtime expresses this equilibrium vacuum limit as a reusable flow-constraint state grammar. The calculations below use that equilibrium limit only; adaptive departures are left for later dynamical work rather than fitted in this manuscript. The paper-local supplementary scripts remain the audited reproduction layer for the figures and tables in this Part I archive.

**Keywords:** Constraint-Driven Field Theory, Constraint-Driven Flux Dynamics, adaptive operating ratio, vacuum structure, Mujjabi tests, reproducible physics, even- $n$  torus knots, topological exclusion

---

\*msbico@gmail.com

## Project Context

This manuscript constitutes Part I (Paper 09) of the multi-part series *Constraint-Driven Flux Dynamics (CDFD)*. The underlying mathematical architecture builds directly upon the concepts established in Paper 08 of this series [11].

## Symbol Conventions

### CDFD physical-vacuum symbol convention.

| Symbol              | Part I meaning                                                                     | Physical reading                                                                                             |
|---------------------|------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------|
| $J$ or $\Phi$       | Driving flux or throughput                                                         | Transport current, field throughput, circulation drive, or generic flux; a subscript fixes the exact channel |
| $C$                 | Active constraint or capacity burden                                               | Vacuum transport capacity, geometric confinement, stiffness, boundary resistance, or load-bearing limit      |
| $S$                 | Surface responsiveness                                                             | Local ability of the vacuum or modeled surface to reconfigure under drive                                    |
| $M_s$               | Structural memory                                                                  | Retained geometry, phase history, stiffness history, or accumulated constraint state                         |
| $\Psi_s$            | Adaptive operating ratio $(J/C) \cdot S \cdot M_s$ or $(\Phi/C) \cdot S \cdot M_s$ | Local bookkeeping gauge for constrained operation; it is not a new universal constant                        |
| $R, a, \chi$        | Ring radius, core radius, and aspect ratio $R/a$                                   | Geometry used to connect vortex structure to dimensionless electromagnetic scales                            |
| $\rho_0, c, \kappa$ | Vacuum density scale, propagation speed, and transport stiffness                   | Medium parameters used in stability, pressure, and self-consistency calculations                             |

### Greek-symbol convention.

| Symbol                 | Local role                                                       | Interpretation rule                                                                                       |
|------------------------|------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------|
| $\alpha$               | Fine-structure constant unless a local equation states otherwise | In the Part I physics papers, un-subscripted $\alpha$ means the electromagnetic fine-structure constant   |
| $\beta, \gamma$        | Local response, stiffness, coupling, or correction coefficients  | Equation-local coefficients; subscripts bind them to the named mode, channel, or approximation            |
| $\theta, \phi, \Phi_c$ | Phase, orientation, or chirality angles                          | Used for torus-knot orientation, vacuum phase, or chiral phase geometry as defined in the nearby equation |
| $\delta, \epsilon$     | Perturbation, residual, tolerance, or small-offset scales        | Local stability margins, numerical residuals, or experimental tolerances                                  |
| $\lambda, \mu, \nu$    | Rate, mass, mode, or frequency labels                            | Meaning is fixed by the equation, subscript, and physics context where the symbol appears                 |
| $\rho, \sigma, \tau$   | Density, spread/noise, or time-scale terms                       | Local density, variance, stress, relaxation, or transport-memory quantities                               |
| $\Omega, \Lambda$      | State/domain space or integrated measure where defined           | Reserved for explicitly defined state spaces, spectra, or synthesis-level quantities                      |

## 1 Introduction

The CDFD series [4–11] has so far treated only odd- $n$  torus knot families  $T(2, n)$ . Paper VIII identified even- $n$  families as an open problem and noted that the universal anti-phase light modes

require interpretation.

The present paper addresses both, plus the status of  $T(2, 2)$ . The torus knot classification [12] and the Faddeev–Niemi energy hierarchy [2] underpin the entire  $T(2, n)$  framework; knot polynomial invariants [3] distinguish families that share the same crossing number, and numerical knot soliton spectra [1] provide independent checks on the energy ordering. Section 2 proves that  $n = 2$  is excluded from the CDFD framework by a purely algebraic obstruction: the Fourier identity that underpins the mass sum rule fails for  $n = 2$  and only  $n = 2$ . Section 3 extends the full hierarchy to even  $n \geq 4$ . Section 4 identifies the anti-phase light modes as boundary modes and derives their mass formula. Section 5 presents the complete  $T(2, n)$  mass census for  $3 \leq n \leq 13$ .

## 1.1 Mujjabi boundary principle

This paper is the first hard boundary test of the hierarchy. A serious theory must say no. The Mujjabi Boundary Principle is:

A candidate CDFD state is physical only if it survives topology, capacity, phase, and stability constraints simultaneously.

The exclusion of  $T(2, 2)$  is therefore not a footnote. It is proof that the framework is not allowed to accept every attractive pattern. If later candidate states require exceptions to this boundary principle, the hierarchy loses predictive force.

## 2 The Fourier Exclusion Theorem: Why $n = 2$ is Inadmissible

### 2.1 The Fourier identity and where it fails

The universal mass sum rule (Paper VII, Theorem 1) rests on a single algebraic identity:

$$\sum_{k=0}^{n-1} \cos^2\left(\theta + \frac{2\pi k}{n}\right) = \frac{n}{2} \quad \text{for all } \theta, \quad (1)$$

which gives  $\sum_k A_k^2 = 2n$  and hence  $\sum_k m_k = 2nM_n$ .

**Theorem 1** (Fourier exclusion). *Identity (1) holds for all  $\theta$  if and only if  $n \geq 3$ . For  $n = 2$  it fails by a theta-dependent residual  $\cos(2\theta)$ .*

*Proof.* Writing  $\cos^2 x = \frac{1}{2} + \frac{1}{2} \cos(2x)$ :

$$\sum_{k=0}^{n-1} \cos^2\left(\theta + \frac{2\pi k}{n}\right) = \frac{n}{2} + \frac{1}{2} \sum_{k=0}^{n-1} \cos\left(2\theta + \frac{4\pi k}{n}\right).$$

The second sum is the real part of  $e^{2i\theta} \sum_{k=0}^{n-1} e^{4\pi i k/n}$ . This geometric sum vanishes if and only if  $e^{4\pi i/n} \neq 1$ , i.e.  $4\pi/n \notin 2\pi\mathbb{Z}$ , i.e.  $n \nmid 2$ . The only integer  $n \geq 2$  that divides 2 is  $n = 2$  itself. For  $n = 2$ :  $e^{4\pi i/2} = e^{2\pi i} = 1$ , so the sum equals  $\sum_{k=0}^1 e^{2i\theta} = 2e^{2i\theta}$ , giving residual  $\text{Re}(2e^{2i\theta}) = 2\cos(2\theta)$ . Hence  $\sum_k \cos^2 = 1 + \cos(2\theta) \neq 1$  in general.  $\square$




Figure 1: The sum  $\sum_k \cos^2(\theta + 2\pi k/n)$  plotted against orientation  $\theta$ . For  $n = 3$  and  $n = 5$ , the sum is perfectly flat at  $n/2$ , guaranteeing the universal mass sum rule. For  $n = 2$ , the sum oscillates, algebraically excluding the  $T(2, 2)$  Hopf link from the mass framework.

## 2.2 Physical consequence: $T(2, 2)$ is excluded

For  $n = 2$ , the sum  $\sum_k A_k^2 = 2 + 2 \cos(2\theta)$  is  $\theta$ -dependent, ranging from 0 (at  $\theta = \pi/2$ ) to 4 (at  $\theta = 0$ ). The universal mass sum rule  $\sum m_k = 2nM_n = 4M_2$  does not hold at a general orientation; it holds only at  $\theta = 0$  (a measure-zero special case).

The self-consistency equation  $2\theta_2 = \tilde{Q}_2(\theta_2)$  does have a numerical solution near  $\theta_2 \approx 22.7^\circ$ , but at this  $\theta$ ,  $\sum A_k^2 \approx 5.40 \neq 4 = 2n$ . The sum rule is violated, and the Brannen parametrisation is self-contradictory: the mass formula  $m_k = M_2 A_k^2$  is no longer anchored by the  $2n$ -normalisation that gives  $M_n$  its physical meaning.

**Corollary 1.** *The Hopf fibration  $T(2, 2)$  is not a member of the CDFD torus knot hierarchy. The minimal valid family is  $T(2, 3)$  (the trefoil, the lepton sector).*

The physical interpretation is natural:  $T(2, 2)$  has only two strands and no braiding complexity; it is topologically an unknot (the two strands can be isotoped apart). The CDFD framework requires genuine knotting, which begins at  $T(2, 3)$ .

### 3 Extension to Even $n \geq 4$

#### 3.1 Validity

For  $n \geq 4$ , the divisibility condition  $n \mid 2$  is not satisfied, so Theorem 1 confirms the Fourier identity holds and the entire CDFD machinery extends without modification.

The self-consistency condition  $n\theta_n = \tilde{Q}_n(\theta_n)$  (Paper VIII, Eq. (2)) applies verbatim. Proposition 1 of Paper VIII (existence and uniqueness) extends to even  $n \geq 4$  by the same intermediate-value and monotonicity argument.

#### 3.2 Self-consistency solutions for even $n$

Table 1 gives the solutions for  $n = 4, 6, 8, 10$ .

Table 1: Self-consistency solutions for even- $n$  torus knot families. The  $n^{7/4}$  hierarchy and sum rule hold for all entries.

| $n$ | Knot       | $\theta_n$ (deg) | $M_n$ (MeV) | $\Sigma_n$ (MeV) | Split   |
|-----|------------|------------------|-------------|------------------|---------|
| 4   | $T(2, 4)$  | 4.9366           | 389.44      | 3115.51          | $3 + 1$ |
| 6   | $T(2, 6)$  | 2.4595           | 527.85      | 6334.15          | $5 + 1$ |
| 8   | $T(2, 8)$  | 1.4540           | 654.96      | 10479.28         | $6 + 2$ |
| 10  | $T(2, 10)$ | 0.8811           | 774.27      | 15485.45         | $7 + 3$ |

#### 3.3 Even- $n$ mode spectra

Table 2:  $T(2, 4)$  mode masses,  $\theta_4 = 4.937^\circ$ ,  $M_4 = 389.44$  MeV.

| Mode  | $m_k$ (MeV) | Sector     |
|-------|-------------|------------|
| $m_0$ | 65.135      | anti-phase |
| $m_1$ | 300.418     | co-phase   |
| $m_2$ | 489.995     | co-phase   |
| $m_3$ | 2259.960    | co-phase   |
| Sum   | 3115.51     | $= 8M_4$   |

Table 3:  $T(2, 6)$  mode masses,  $\theta_6 = 2.459^\circ$ ,  $M_6 = 527.85$  MeV.

| Mode  | $m_k$ (MeV) | Notes                  |
|-------|-------------|------------------------|
| $m_0$ | 30.655      | anti-phase             |
| $m_1$ | 63.229      | anti-phase (near zero) |
| $m_2$ | 89.995      | anti-phase (near zero) |
| $m_3$ | 1443.860    | co-phase               |
| $m_4$ | 1633.222    | co-phase               |
| $m_5$ | 3073.194    | co-phase (heaviest)    |
| Sum   | 6334.15     | $= 12M_6$              |

Table 4:  $T(2, 8)$  mode masses,  $\theta_8 = 1.454^\circ$ ,  $M_8 = 654.96$  MeV. This family has two extremely light modes ( $< 1$  MeV), the lightest in the entire hierarchy.

| Mode  | $m_k$ (MeV) |
|-------|-------------|
| $m_0$ | 0.411       |
| $m_1$ | 0.432       |
| $m_2$ | 112.126     |
| $m_3$ | 608.793     |
| $m_4$ | 702.804     |
| $m_5$ | 2552.933    |
| $m_6$ | 2685.863    |
| $m_7$ | 3815.918    |
| Sum   | 10479.28    |

Table 5:  $T(2, 10)$  mode masses,  $\theta_{10} = 0.881^\circ$ ,  $M_{10} = 774.27$  MeV.

| Mode  | $m_k$ (MeV) |
|-------|-------------|
| $m_0$ | 13.329      |
| $m_1$ | 19.029      |
| $m_2$ | 132.737     |
| $m_3$ | 227.749     |
| $m_4$ | 263.816     |
| $m_5$ | 1553.077    |
| $m_6$ | 1645.125    |
| $m_7$ | 3516.772    |
| $m_8$ | 3601.652    |
| $m_9$ | 4512.167    |
| Sum   | 15485.45    |

### 3.4 Amplitude-split formula for even $n$

The formula of Paper VIII, Theorem 2:  $n_- = \lfloor 5n/8 \rfloor - \lfloor 3n/8 \rfloor + 1$ , extends to even  $n$  with one exception. Table 6 gives results.

The  $n = 8$  exception arises because  $3n/8 = 3$  and  $5n/8 = 5$  are exact integers: the modes  $k = 3$  and  $k = 5$  sit exactly at the amplitude-zero boundaries. At the self-consistency orientation  $\theta_8 \approx 1.45^\circ > 0$ , these boundary modes are shifted slightly positive, giving  $n_- = 2$  rather than 3. The formula requires refinement at integer values of  $3n/8$  and  $5n/8$ : the count is  $n_-^{\text{corrected}} = n_-^{\text{formula}} - 1$  when  $\{3n/8\} = 0$  or  $\{5n/8\} = 0$  (i.e. when  $8 \mid n$ ).

## 4 The Anti-Phase Modes as Boundary Modes

### 4.1 The light modes are not Goldstone bosons

A natural hypothesis for the near-zero modes would be that they are pseudo-Nambu–Goldstone bosons (PNGBs): massless modes of a broken continuous symmetry that acquire a small mass from an explicit symmetry-breaking term proportional to some small parameter. If so, their masses




Figure 2: Mode mass spectra for the even- $n$  torus knot families  $n = 4, 6, 8, 10$ . The characteristic CDFD mode clustering (light boundary modes, intermediate co-phase modes, and heavy outlier modes) is clearly visible across all families.

should scale as  $m \propto M_n \theta_n^2$  (the square of the symmetry-breaking angle).

This is *not* the case. Numerically,  $m_{\min}/(M_n \theta_n^2)$  varies from 1.9 ( $n = 5$ ) to 782 ( $n = 11$ ) with no discernible universal law. The hypothesis is refuted.

## 4.2 The boundary mode identification

The amplitude  $A_k = 1 + \sqrt{2} \cos \phi_k$  vanishes at  $\phi_k = 3\pi/4$  and  $\phi_k = 5\pi/4$ . The anti-phase modes are those whose phase  $\phi_k = \theta_n + 2\pi k/n$  falls closest to one of these zeros.

**Theorem 2** (Boundary mode mass). *The minimum mode mass in any  $T(2, n)$  family ( $n \geq 5$ ) is:*

$$m_{\min} = M_n \cdot A_{\min}^2, \quad A_{\min} = 1 + \sqrt{2} \cos(\theta_n + 2\pi \lceil 3n/8 \rceil / n), \quad (2)$$

where  $k^* = \lceil 3n/8 \rceil$  is the mode index whose phase is closest to the lower amplitude zero at  $3\pi/4$ .

The key non-universality is the fractional part  $\{3n/8\}$ , which cycles through the values  $\{1/8, 3/8, 5/8, 7/8, 1/8, \dots\}$  as  $n$  runs through odd multiples of 1, and through different values for even  $n$ . This fractional offset controls how far the boundary mode sits from the zero, and hence how light it is.

## 4.3 Interpretation

The two anti-phase modes are topological boundary modes of the torus knot field configuration. In the  $\mathbb{Z}_n$  Fourier decomposition of the knotted soliton, the modes at  $k^*$  and  $n - k^*$  are the ones whose

Table 6: Amplitude splits for even- $n$  families. The formula matches numerical results except at  $n = 8$ .

| $n$ | Formula $n_-$ | Numerical $n_-$ | Match | Note                                                                                                        |
|-----|---------------|-----------------|-------|-------------------------------------------------------------------------------------------------------------|
| 4   | 1             | 1               | Yes   | —                                                                                                           |
| 6   | 1             | 1               | Yes   | —                                                                                                           |
| 8   | 3             | 2               | No    | Boundary integer: $\lfloor 5 \cdot 8/8 \rfloor = 5$ , $\lceil 3 \cdot 8/8 \rceil = 3$ ; mode at boundary is |
| 10  | 3             | 3               | Yes   | —                                                                                                           |

phase most nearly cancels the background condensate. They are structurally enforced by the sum rule: because  $\sum A_k^2 = 2n$  is fixed and the three (or more) co-phase modes carry large amplitude, the anti-phase pair must carry correspondingly small amplitude.

This is geometrically analogous to the *nodal modes* of a vibrating string: a string of length  $L$  has a mode at every  $2L/n$  wavelength, and the mode whose wavelength is closest to  $2L/3$  (the tripling resonance) is the one with lowest energy. The CDFD anti-phase modes play this role in the torus knot field.

They are not observable as free particles; they are the Fourier “ghost modes” that enforce the global sum rule. This resolves Paper VIII open problem 2.

## 5 The Complete $T(2, n)$ Mass Census

Table 7 assembles the complete CDFD torus knot hierarchy for  $3 \leq n \leq 13$ , combining the odd- $n$  results of Papers VI–VIII with the even- $n$  results of this paper. All energies derive from the single input  $m_e = 0.51100$  MeV, via  $M_3$  (Paper V) and the Faddeev–Niemi scaling  $M_n = M_3(n/3)^{3/4}$  (Paper IV).

Table 7: Complete CDFD torus knot mass census,  $3 \leq n \leq 13$ .  $\Sigma_n = 2nM_n$  (exact, Paper VII).  $\theta_n$ : algebraic for  $n = 3$ ; self-consistency fixed point for  $n \geq 4$ . Dagger ( $\dagger$ ): the split formula requires a boundary correction (see Section 3).

| $n$ | Knot       | $\theta_n$ (deg) | $M_n$ (MeV) | $\Sigma_n$ (MeV) | Split           | Status         |
|-----|------------|------------------|-------------|------------------|-----------------|----------------|
| 3   | $T(2, 3)$  | 12.732           | 313.86      | 1883.2           | $3 + 0$         | Verified (PDG) |
| 4   | $T(2, 4)$  | 4.937            | 389.44      | 3115.5           | $3 + 1$         | Predicted      |
| 5   | $T(2, 5)$  | 3.698            | 460.39      | 4603.9           | $3 + 2$         | Predicted      |
| 6   | $T(2, 6)$  | 2.459            | 527.85      | 6334.2           | $5 + 1$         | Predicted      |
| 7   | $T(2, 7)$  | 1.749            | 592.54      | 8295.6           | $5 + 2$         | Predicted      |
| 8   | $T(2, 8)$  | 1.454            | 654.96      | 10479.3          | $6 + 2^\dagger$ | Predicted      |
| 9   | $T(2, 9)$  | 1.077            | 715.44      | 12878.0          | $7 + 2$         | Predicted      |
| 10  | $T(2, 10)$ | 0.881            | 774.27      | 15485.5          | $7 + 3$         | Predicted      |
| 11  | $T(2, 11)$ | 0.742            | 831.65      | 18296.2          | $9 + 2$         | Predicted      |
| 12  | $T(2, 12)$ | 0.608            | 886.62      | 21279.0          | $9 + 3$         | Predicted      |
| 13  | $T(2, 13)$ | 0.528            | 942.65      | 24509.0          | $9 + 4$         | Predicted      |




Figure 3: The minimum mode mass  $m_{\min}$  vs the pseudo-Goldstone scaling expectation  $M_n \theta_n^2$ . The lack of correlation visually refutes the hypothesis that the light modes are pseudo-Nambu-Goldstone bosons of a broken continuous symmetry.

### 5.1 The $n^{7/4}$ hierarchy: extended

The scaling law  $\Sigma_n = 2M_3 \cdot 3^{-3/4} \cdot n^{7/4}$  (Paper VII) holds for all  $n \geq 3$ , odd and even. The even families fill in the gaps of the odd hierarchy, producing a smooth, monotonically increasing sequence. The dominant pattern is that  $\Sigma_n$  grows by roughly 1100–1500 MeV for each unit step in  $n$  across the entire range.

### 5.2 Theta convergence: the full sequence

The supplementary script computes  $\theta_n$  for  $3 \leq n \leq 13$ . The convergence  $\theta_n \rightarrow 0$  is monotone and well-described by the asymptotic  $2\pi^2/[(\pi + 4)^2 n^2]$  derived in Paper VIII, confirming that the large- $n$  limit applies equally to odd and even families.

## 6 Open Problems Closed and Remaining

### 6.1 Closed

(a) **Even- $n$  families (Paper VIII open problem 4):** resolved by extension of the CDFD framework to all  $n \geq 4$ .

(b) **Anti-phase light mode identification (Paper VIII open problem 2):** identified as




Figure 4: Theta convergence for the full sequence  $3 \leq n \leq 13$ . The exact numerical solutions (even and odd) align tightly with the  $1/n^2$  continuous vacuum asymptotic prediction.

boundary modes whose mass is set by the fractional offset  $\{3n/8\}$ , not by any symmetry-breaking mechanism.

(c) **Exclusion of  $T(2, 2)$ :** proved algebraically by Theorem 1.

## 6.2 Remaining

The single surviving open problem from Papers VII–IX is:

1. **Physical principle selecting the self-consistency condition.** Papers VII and VIII established that  $n\theta_n = \tilde{Q}_n(\theta_n)$  has a unique solution for every  $n \geq 3$ , with an exact closed-form asymptotic. For  $n = 3$ , the condition is motivated by the constancy of  $Q_3$ . For  $n \geq 4$ , a derivation from the CDFD vacuum equation of state is needed. This is the final open problem of the series.

## 7 Conclusion

The Fourier exclusion theorem (Theorem 1) identifies  $n = 2$  as the unique excluded family: the Fourier identity  $\sum \cos^2(\theta + 2\pi k/n) = n/2$ , which anchors the entire CDFD mass framework, fails for  $n = 2$  and no other  $n$ . The  $T(2, 2)$  Hopf fibration is topologically an unknot and is algebraically excluded.

For all  $n \geq 4$  (even and odd), the framework is valid. Complete mode spectra for  $n = 4, 6, 8, 10$  are tabulated (Tables 2–5) and the amplitude-split formula is verified with a single boundary-integer correction at  $n = 8$ .

The anti-phase light modes present in every family  $n \geq 5$  are identified as boundary modes: their mass is determined by the fractional proximity  $\{3n/8\}$  of the nearest Fourier mode to the amplitude zeros at  $\cos \phi = -1/\sqrt{2}$ . They are not Goldstone bosons, not confinement-driven zero modes, and not artifacts: they are the modes that enforce the global sum rule  $\sum m_k = 2nM_n$  when the co-phase modes carry large amplitude.

The complete  $T(2, n)$  mass census for  $3 \leq n \leq 13$  (Table 7) constitutes the full CDFD torus knot prediction table, derived entirely from  $m_e$  with zero additional free parameters.

## Supplementary Material

This manuscript is accompanied by release-archive supplementary material in the canonical Part I tree. The relevant paper-local Python scripts, numerical checks, generated figure outputs, tabulated data, figure assets, and environment notes are maintained under `notebooks/`, `outputs/`, `figures/`, and `REPRODUCIBILITY.md` in `Part_I_Fundamental_Physics/`.

## Data and Code Availability

The release-archive supplementary material is included in the archived Part I release on Zenodo at <https://doi.org/10.5281/zenodo.20250820>. The current version DOI is assigned by Zenodo after each GitHub release. Zenodo is the permanent release archive, and the public source archive is available on GitHub at <https://github.com/bampita-bico/CDFD-Part-I-Release>. The broader public CDFD Runtime is separate reusable software infrastructure; the numerical values reported here are reproduced from this paper’s Supplementary Material.

## Acknowledgments

The author used AI-assisted tools for formatting checks and code-verification support. The scientific claims, interpretations, and final manuscript decisions are the author’s responsibility.

The author is indebted to the foundational work of D. Rolfsen [12] (torus knot classification), V.F.R. Jones [3] (knot polynomial invariants), R.A. Battye and P.M. Sutcliffe [1] (knot soliton energy spectra), and L. Faddeev and A.J. Niemi [2] (knots and particles) — whose work underpins the extension to even- $n$  torus knots.

## Declarations

### Funding

This research received no external institutional funding.

### Author Contributions

Steve Bico Mujjabi: conceptualization, methodology, formal analysis, software, validation, visualization, writing—original draft, and writing—review and editing.

## Conflicts of Interest

The author declares no conflicts of interest.

## References

## References

- [1] Richard A. Battye and Paul M. Sutcliffe. Knots as stable soliton solutions in a three-dimensional classical field theory. *Physical Review Letters*, 81:4798–4801, 1998. doi: 10.1103/PhysRevLett.81.4798. Numerical construction of stable knotted solitons; mass–energy spectrum of torus knot configurations.
- [2] Ludwig Faddeev and Antti J. Niemi. Stable knot-like structures in classical field theory. *Nature*, 387:58–61, 1997. doi: 10.1038/387058a0.
- [3] Vaughan F. R. Jones. A polynomial invariant for knots via von Neumann algebras. *Bulletin of the American Mathematical Society*, 12:103–111, 1985. doi: 10.1090/S0273-0979-1985-15304-2. The Jones polynomial; distinguishes torus knots  $T(2, n)$  that share the same crossing number.
- [4] Steve Bico Mujjabi. Vortex Stability in a Constrained Transport Vacuum and the Origin of the Fine-Structure Constant. Zenodo, 2026. URL <https://doi.org/10.5281/zenodo.21420457>. <https://doi.org/10.5281/zenodo.21420457>.
- [5] Steve Bico Mujjabi. Lepton Masses from Z3-Symmetric Vortex Modes and the Origin of the Koide Formula. Zenodo, 2026. URL <https://doi.org/10.5281/zenodo.21421496>. <https://doi.org/10.5281/zenodo.21421496>.
- [6] Steve Bico Mujjabi. Topology, Chirality, and the Vacuum Density: Closing All Open Problems of the CDFT Series. Zenodo, 2026. URL <https://doi.org/10.5281/zenodo.21421906>. <https://doi.org/10.5281/zenodo.21421906>.
- [7] Steve Bico Mujjabi. The Vacuum Equation of State and the Self-Consistency of the CDFT Lepton Sector. Zenodo, 2026. URL <https://doi.org/10.5281/zenodo.21425048>. <https://doi.org/10.5281/zenodo.21425048>.
- [8] Steve Bico Mujjabi. Zero Parameters: Deriving the Vacuum Chirality Phase from the Koide Invariant. Zenodo, 2026. URL <https://doi.org/10.5281/zenodo.21427696>. <https://doi.org/10.5281/zenodo.21427696>.
- [9] Steve Bico Mujjabi. The Universal Torus Knot Hierarchy: Zn Symmetry, the Cinquefoil Family, and the General Structure of CDFT Particle Families. Zenodo, 2026. URL <https://doi.org/10.5281/zenodo.21431810>. <https://doi.org/10.5281/zenodo.21431810>.
- [10] Steve Bico Mujjabi. The Universal Mass Sum Rule: A Theta-Independent Prediction for All CDFT Torus Knot Families. Zenodo, 2026. URL <https://doi.org/10.5281/zenodo.21434735>. <https://doi.org/10.5281/zenodo.21434735>.
- [11] Steve Bico Mujjabi. The CDFT Torus Knot Spectrum: Closing the Four Open Problems of Paper VII. Zenodo, 2026. URL <https://doi.org/10.5281/zenodo.21434949>. <https://doi.org/10.5281/zenodo.21434949>.

206 [12] Dale Rolfsen. *Knots and Links*. Publish or Perish, 1976. Standard reference on torus knot  
207 classification  $T(p, q)$ , knot invariants, and linking numbers.