

Paper XI Addendum: A Constrained-Transport Hypothesis Program for Cosmological Analogies and Falsification Criteria

Steve Bico Mujjabi, MD*
VuraLabs, Kampala, Uganda
Independent researcher
ORCID: 0009-0001-0556-5516

<https://github.com/bampita-bico/CDFD-Part-I-Release>

July 2026

Abstract

This speculative addendum records cosmological analogies expressed in the Constraint-Driven Flux Dynamics (CDFD) vocabulary. It asks whether dark-matter phenomenology, dark energy, black-hole horizons, and the proposed pre-geometric “Blancken layer” suggest useful questions about constrained transport; it does not offer a cosmological theory or a derivation of those phenomena.

The preceding papers provide selected conditional mathematical identities and model constructions, not established lepton, vacuum, or cosmological laws. Every dark-sector and Blancken-layer statement here remains a hypothesis until it supplies a quantitative discriminant beyond established theory. The shared runtime state grammar is used only to organize possible tests, not as cosmological evidence.

The public CDFD Runtime implements the same flow-constraint grammar for reusable simulations. The paper-local supplementary script remains the audited reproduction layer for the figures and tables in this Part I archive.

Keywords: Constraint-Driven Field Theory, Constraint-Driven Flux Dynamics, adaptive operating ratio, vacuum structure, Mujjabi tests, reproducible physics, dark matter, dark energy, black holes

Project Context

This manuscript constitutes Part I (Paper 11) of the multi-part series *Constraint-Driven Flux Dynamics (CDFD)*. The underlying mathematical architecture builds directly upon the concepts established in Paper 10 of this series [4].

Symbol Conventions

CDFD physical-vacuum symbol convention.

*msbico@gmail.com

Symbol	Part I meaning	Physical reading
J or Φ	Driving flux or throughput	Transport current, field throughput, circulation drive, or generic flux; a subscript fixes the exact channel
C	Active constraint or capacity burden	Vacuum transport capacity, geometric confinement, stiffness, boundary resistance, or load-bearing limit
S	Surface responsiveness	Local ability of the vacuum or modeled surface to reconfigure under drive
M_s	Structural memory	Retained geometry, phase history, stiffness history, or accumulated constraint state
Ψ_s	Adaptive operating ratio $(J/C) \cdot S \cdot M_s$ or $(\Phi/C) \cdot S \cdot M_s$	Local bookkeeping gauge for constrained operation; it is not a new universal constant
R, a, χ	Ring radius, core radius, and aspect ratio R/a	Geometry used to connect vortex structure to dimensionless electromagnetic scales
ρ_0, c, κ	Vacuum density scale, propagation speed, and transport stiffness	Medium parameters used in stability, pressure, and self-consistency calculations

8 Greek-symbol convention.

Symbol	Local role	Interpretation rule
α	Fine-structure constant unless a local equation states otherwise	In the Part I physics papers, un-subscripted α means the electromagnetic fine-structure constant
β, γ	Local response, stiffness, coupling, or correction coefficients	Equation-local coefficients; subscripts bind them to the named mode, channel, or approximation
θ, ϕ, Φ_c	Phase, orientation, or chirality angles	Used for torus-knot orientation, vacuum phase, or chiral phase geometry as defined in the nearby equation
δ, ϵ	Perturbation, residual, tolerance, or small-offset scales	Local stability margins, numerical residuals, or experimental tolerances
λ, μ, ν	Rate, mass, mode, or frequency labels	Meaning is fixed by the equation, subscript, and physics context where the symbol appears
ρ, σ, τ	Density, spread/noise, or time-scale terms	Local density, variance, stress, relaxation, or transport-memory quantities
Ω, Λ	State/domain space or integrated measure where defined	Reserved for explicitly defined state spaces, spectra, or synthesis-level quantities

10 1 Motivation

Modern cosmology contains several successful but conceptually separate descriptions. Dark matter is inferred from galaxy dynamics, cluster lensing, and precision cosmology [2, 6, 8]. Dark energy is inferred from supernova acceleration and the cosmological parameter ladder [5–7]. Black holes are described by general relativity, horizon mechanics, and semiclassical thermodynamics [1, 3]. Particle masses are treated within quantum field theory. CDFD does not replace these mature descriptions. Its proposed contribution is a unifying transport ontology:

$$\Psi_{s,\text{vac}} = \frac{\Phi_{\text{vac}}}{C_{\text{vac}}}, \quad (1)$$

where Φ_{vac} is vacuum flux or activity and C_{vac} is the effective transport capacity, stiffness, or constraint of the vacuum medium.

19 The question is whether apparent anomalies arise where vacuum flux and vacuum constraint
 20 are forced into extreme configurations. The answer is not asserted as established physics; it is
 21 formulated as a sequence of hypotheses with explicit failure criteria.




Figure 1: Cosmological bridge map. Dark matter, dark energy, black-hole saturation, and the Blancken-layer proposal are shown as hypothesis classes attached to the same constrained-transport operator. The lower bar states the required falsification standard.

22 2 Vacuum Constraint Geometry

23 The common mathematical object is the constrained transport operator:

$$\mathcal{T}[\Phi, C] = \nabla \cdot (C^{-1} \nabla \Phi). \quad (2)$$

24 In ordinary diffusion, transport depends only on gradient and conductivity. In CDFD, conductivity
 25 is itself a dynamical field. Large flux gradients alter constraint; altered constraint bends subsequent
 26 transport. This feedback is the mechanism by which geometry can emerge from transport.

27 **Hypothesis 1** (Vacuum constraint feedback). *The vacuum behaves, at an effective level, as a*
 28 *transport medium whose constraint field responds to persistent flux stress:*

$$\partial_t C_{\text{vac}} = \alpha_{\text{vac}} |\Phi_{\text{vac}}| - \beta_{\text{vac}} C_{\text{vac}} + \gamma_{\text{vac}} \nabla^2 C_{\text{vac}}. \quad (3)$$

29 Here α_{vac} is the vacuum constraint-reinforcement coefficient, β_{vac} is the vacuum relaxation
 30 coefficient, and γ_{vac} is the spatial smoothing or diffusion coefficient for the effective constraint field.

31 This is the bridge assumption. If no such effective feedback exists, the rest of this paper collapses
 32 into metaphor.

3 Archive Source, Reproducibility, and Claim Discipline

This manuscript expands the archived physical-mysteries synthesis rather than discarding it. The archive contained five compact source claims:

1. dark matter as geometric vacuum hardening;
2. dark energy as adaptive metric expansion;
3. black holes as transport-capacity rupture;
4. the mass spectrum as vacuum topology;
5. the Blancken layer as a pre-geometric substrate.

Those claims are retained, but their status is corrected. The paper does not claim to have replaced general relativity, particle dark matter searches, or precision cosmology. It gives a constraint-transport research program whose cosmological claims must produce independent discriminants before they can be called physical explanations.

The original synthesis figures were archive assets. In this version they are regenerated by `supplementary_XI.py` into `outputs/paper_XI/`, with CSV tables and check files recording the toy assumptions behind each conceptual panel. The figures remain schematics; their value is that the bridge claims now have an auditable source trail rather than orphaned images.

Archive claim	Retained kernel	Correct status here
Dark matter	constraint gradients can mimic excess lensing	hypothesis requiring residual-lensing tests
Dark energy	vacuum self-consistency can leave pressure residuals	hypothesis linked to Paper X
Black holes	horizons are capacity saturation boundaries	interpretive bridge to GR
Mass topology	stable masses are topological constraint states	inherited from Papers I–VI
Blancken layer	pre-geometric substrate below spacetime	speculative until invariant is derived

4 Dark Matter as Constraint-Induced Lensing

The CDFD dark-matter interpretation is not that ordinary matter is secretly missing from observations. It is that some portion of the gravitationally inferred anomaly could be modeled as vacuum constraint geometry induced by persistent flux aggregation.

Let ρ_b be baryonic mass density and let ΔC_{vac} be the induced constraint deformation. The effective lensing potential is written schematically:

$$\Phi_{\text{lens}} = \Phi_{\text{GR}}[\rho_b] + \Phi_{\text{CDFD}}[\Delta C_{\text{vac}}]. \quad (4)$$

The second term is the hypothesis. It says that the vacuum remembers or stiffens around long-lived flux structures, producing lensing behavior not accounted for by visible matter alone.

Prediction 1 (Geometry-memory signature). *If dark matter is partly vacuum-constraint geometry, lensing residuals should correlate not only with present baryonic mass but also with indicators of long-term flux history: merger pathways, rotational shear, plasma history, or persistent high-throughput structure.*

Criterion 1 (Dark-matter falsification). *The CDFD dark-matter claim fails if lensing residuals require no history-dependent or constraint-like term once baryons, standard dark matter models, and modified-gravity alternatives are properly fitted.*




Figure 2: Reproducible Paper XI conceptual output for the dark-matter claim: persistent flux aggregation hardens the surrounding constraint geometry, producing lensing-like curvature. In this manuscript the image is treated as a conceptual model, not as observational evidence.

4.1 Observable Residual Protocol

The dark-matter claim becomes scientific only if it specifies what to fit. The proposed observational object is a residual field

$$R_{\text{lens}}(\mathbf{x}) = \Phi_{\text{obs}}(\mathbf{x}) - \Phi_{\text{GR+baryon}}(\mathbf{x}) - \Phi_{\text{standard DM}}(\mathbf{x}|\Theta), \quad (5)$$

where Θ denotes the best-fit parameters of a conventional dark-matter model. A CDFD contribution must then correlate with a measurable history functional

$$H_C(\mathbf{x}, t) = \int_{-\infty}^t K(t - t') [\alpha_{\text{vac}} |\Phi(\mathbf{x}, t')| + \gamma_{\text{vac}} \nabla^2 C(\mathbf{x}, t')] dt', \quad (6)$$

not merely with current baryonic mass. The falsifiable statement is: clusters or galaxies with similar baryonic mass but different merger, plasma, or shear histories should show different residual constraint patterns if the CDFD term is real.

4.2 Relation to Modified Gravity

CDFD is not identical to modified Newtonian dynamics or a generic modified-gravity ansatz. In this paper the modification is path-dependent: the constraint field has memory. A static algebraic relation between acceleration and baryonic matter would not be sufficient to confirm the CDFD interpretation. The distinguishing feature must be hysteresis, relaxation, or spatial lag in the inferred constraint field.

5 Dark Energy as Residual Vacuum Self-Consistency

The vacuum equation-of-state paper [4] develops the condition $\Psi_{s,\text{vac}} \rightarrow 1$ as a stability point. In the cosmological bridge, dark energy is interpreted as the residual pressure of a vacuum that approaches, but has not globally completed, self-consistency.

At perfect local balance:

$$\Phi_{\text{vac}} \approx C_{\text{vac}}. \quad (7)$$

The residual term is:

$$\delta_{\text{vac}} = \frac{\Phi_{\text{vac}} - C_{\text{vac}}}{C_{\text{vac}}}. \quad (8)$$

If δ_{vac} remains nonzero on cosmological scales, it may appear as a large-scale pressure term.

Hypothesis 2 (Residual expansion). *Cosmic acceleration is a residual of incomplete vacuum self-consistency at Hubble scale rather than an independent substance.*

Prediction 2 (Scale dependence). *The effective dark-energy term should be close to constant where the vacuum is already equilibrated, but may show subtle scale or epoch dependence where $\nabla^2 C_{\text{vac}}$ remains significant.*




Figure 3: Reproducible Paper XI conceptual output for adaptive expansion. The revised claim is narrower: cosmic acceleration is modeled as a residual of incomplete vacuum self-consistency, not as proof that the residual has already been derived from first principles.

5.1 Connection to Paper X

Paper X gives the local vacuum condition

$$\Psi_{s,\text{vac}} = \frac{\Phi_{\text{vac}}}{C_{\text{vac}}} \rightarrow 1. \quad (9)$$

The cosmological extension is not automatic. It requires a scale-separated residual:

$$\rho_{\Lambda, \text{eff}} = \mathcal{R}[\Phi_{\text{vac}} - C_{\text{vac}}, \nabla C_{\text{vac}}, \nabla^2 C_{\text{vac}}, H(t)], \quad (10)$$

where $\mathcal{R}$ is the still-unfinished residual map. The archive correctly identified the near-cancellation idea, but the mature manuscript must state the open calculation: derive $\mathcal{R}$ with units, boundary conditions, and cosmological parameters fixed before comparison with data.

6 Black Holes as Constraint Saturation Boundaries

In CDFD language, a black hole is not merely high density. It is the limit where the transport system reaches a local boundary condition:

$$J \rightarrow J_{\text{crit}}, \quad \Psi_s \rightarrow \infty, \quad (11)$$

because flux demand overwhelms outward transport capacity. The horizon is interpreted as the surface at which constraint geometry becomes causally dominant.

This does not remove the general relativistic horizon. It reframes the horizon as the macroscopic expression of a transport-capacity limit. The useful test is whether the CDFD variables add predictive structure to black-hole thermodynamics, ringdown deviations, accretion-state transitions, or information-flow constraints.

Criterion 2 (Black-hole falsification). *The CDFD black-hole interpretation fails if it yields no measurable distinction from ordinary general relativity or from known effective-field corrections.*




Figure 4: Reproducible Paper XI conceptual output for black-hole rupture. The figure is retained as a transport-capacity visualization; the actual physics claim must be tested through horizon, ringdown, or thermodynamic discriminants.

6.1 Capacity Boundary Formulation

Let J denote outward information/energy transport through a surface $\mathcal{S}$ and let $J_{\text{crit}}[C]$ denote the local capacity allowed by the constraint geometry. A CDFD horizon is the boundary at which

$$\frac{J}{J_{\text{crit}}[C]} \rightarrow 1^-, \quad \partial_r \left(\frac{J}{J_{\text{crit}}[C]} \right) > 0. \quad (12)$$

This is not meant to erase the relativistic event horizon. It reframes the horizon as a saturation surface in a transport medium. A useful next calculation would show whether $J_{\text{crit}}[C]$ reproduces the Schwarzschild radius in the classical limit and whether it predicts any controlled deviation in rotating or charged geometries.

6.2 Where the Claim Can Fail

The black-hole bridge fails if the capacity-boundary language cannot recover ordinary GR in regimes where GR is already verified. It also fails if it produces no additional observable beyond a relabeling of established geometry. Candidate discriminants are ringdown mode shifts, accretion-state hysteresis, entropy-area corrections, or information-flow bounds. Without one of these, this section remains a conceptual translation.

7 The Blancken Layer

The Blancken layer is the most speculative part of the physics sequence. It is proposed as a pre-geometric substrate below ordinary spacetime description. In this view, spacetime is not fundamental but emerges as a stable constraint phase of a deeper continuous medium.

The role of the Blancken layer is conceptual: it provides a place for the continuous field before metric geometry, particle topology, or classical spacetime have appeared. A responsible formulation must keep it hypothesis-level until it produces mathematical consequences.

Hypothesis 3 (Pre-geometric substrate). *There exists a sub-Planckian continuous layer whose stable macroscopic excitations appear as vacuum geometry, particle topology, and cosmic expansion.*

Criterion 3 (Blancken-layer falsification). *The Blancken-layer claim remains non-scientific unless it yields a calculable invariant, scaling law, or observational discriminant not already supplied by established theory.*




Figure 5: Reproducible Paper XI conceptual output for the Blancken layer. In the revised manuscript the layer is explicitly marked as speculative until it yields a calculable invariant or boundary condition.

7.1 Minimum Mathematical Requirement

The Blancken layer cannot remain a name for “whatever is below spacetime.” For the proposal to become physics, it must produce at least one of the following:

1. a dimensionless invariant preserved under emergence of metric geometry;
2. a boundary condition that fixes the vacuum self-consistency selector used across Papers VI–X;
3. a scaling law connecting sub-Planckian constraint to an observed constant or cosmological residual;
4. a no-go theorem showing which effective geometries cannot emerge from constrained transport.

The most valuable target is item 2: if the Blancken layer can derive the fixed-point condition $n\theta_n = \tilde{Q}_n(\theta_n)$, it would strengthen both the torus-knot hierarchy and the cosmological bridge.

8 Cosmological Mujjabi Tests

The cosmology bridge is the large-scale version of the same question asked in Paper XII: does constraint history leave an observable residual after standard models are fitted? Three targets follow from the source notes and the Part I grammar:

1. **Alpha drift as response drift.** Here un-subscripted α denotes the electromagnetic fine-structure constant. If S or M_s varies over cosmological environments, high-precision spectral data should bound or reveal a correlated shift in effective α .
2. **Lensing hysteresis.** If dark-matter-like effects are constraint-history geometry, residual lensing should correlate with a history functional, not only with instantaneous baryonic mass.
3. **Horizon information bounds.** If black holes are capacity saturation boundaries, ring-down, entropy-area, or information-flow observables must recover GR first and then state a controlled correction.

These are Mujjabi Tests at cosmological scale. They are not evidence until they produce a discriminant that standard cosmology cannot absorb with fewer assumptions.

9 Connection to the Mass Spectrum

The physics series treats particle masses as self-consistent topological states of constrained vacuum transport. The dark-sector synthesis depends on the same premise: stable objects appear when continuous flux becomes locked by constraint topology. The difference is scale. In the lepton papers, the topology is microscopic and quantified through mass ratios and knot families. In the cosmological bridge, the topology is macroscopic and expressed through lensing, horizons, and expansion.

This scale analogy is not proof. It is a proposed research program:

1. derive microscopic self-consistency conditions;
2. test mass-spectrum predictions;
3. extend the same constraint operator to vacuum geometry;
4. search for dark-sector observables that require a constraint-history term.

10 Submission-Grade Falsification Program

For this paper to stand beside the more mathematical physics papers, each cosmological bridge must be tied to a concrete failure route:

Bridge	Primary test	Failure condition
Dark matter	residual lensing vs history functional H_C	no history-dependent residual after standard fits
Dark energy	derivation of $\rho_{\Lambda, \text{eff}}$ from $\Psi_{s, \text{vac}} \rightarrow 1$	wrong units, free tuning, or w incompatible with data
Black holes	capacity-boundary recovery of GR plus controlled correction	no recovery of GR or no distinct prediction
Blancken layer	invariant or boundary condition below metric geometry	no calculable quantity beyond metaphor
Mass topology bridge	derivation of fixed-point selector from vacuum EOS	spectra need fitted selectors per family

This is deliberately stricter than the archive language. The older draft captured the conceptual map; this version states what would make the map scientific and what would break it.

11 Reproducibility Artifacts

The accompanying script `supplementary_XI.py` generates the figures and tables used by this manuscript. The notebook `notebooks/paper_XI_fullstack.ipynb` calls the same function for an interactive reproduction path.

Artifact	Role
<code>table1_cosmology_bridge_map.csv</code>	bridge taxonomy and failure criteria
<code>table2_dark_matter_residual_sample.csv</code>	toy residual sample used for the lensing-history gate
<code>table3_dark_energy_residual_curve.csv</code>	epoch-dependent residual-pressure proxy
<code>table4_black_hole_capacity_boundary.csv</code>	transport-saturation boundary proxy
<code>table5_blancken_layer_requirements.csv</code>	minimum mathematical requirements for the substrate claim
<code>checks_summary_XI_gate.csv</code>	consolidated reproducibility gate

12 Summary of Status

Claim	Current status	Required next step
Vacuum $\Psi_s = 1$ stability	derived within CDFD assumptions	independent mathematical review
Lepton mass topology	model-derived and numerically checkable	precision comparison and external critique
Dark matter as constraint geometry	hypothesis	lensing residual discriminant
Dark energy as residual self-consistency	hypothesis	scale/epoch-dependent prediction
Black holes as transport saturation	hypothesis	ringdown or thermodynamic discriminant
Blancken layer	speculative hypothesis	calculable invariant

13 Scope Boundary and Conclusion

This addendum is not a foundation for a physics program. It supplies no established self-consistency, mass-spectrum, dark-matter, dark-energy, black-hole, or pre-geometric derivation. Its value is limited to making speculative mappings falsifiable: each must mature into a quantitative, independently testable discriminant beyond established theory or be discarded.

Supplementary Material

This manuscript is accompanied by release-archive supplementary material in the canonical Part I tree. The relevant paper-local Python scripts, numerical checks, generated figure outputs, tabulated data, figure assets, and environment notes are maintained under `notebooks/`, `outputs/`, `figures/`, and `REPRODUCIBILITY.md` in `Part_I_Fundamental_Physics/`.

Data and Code Availability

The release-archive supplementary material is included in the archived Part I release on Zenodo at <https://doi.org/10.5281/zenodo.20250820>. The current version DOI is assigned by Zenodo after each GitHub release. Zenodo is the permanent release archive, and the public source archive is available on GitHub at <https://github.com/bampita-bico/CDFD-Part-I-Release>. The broader public CDFD Runtime is separate reusable software infrastructure; the numerical values reported here are reproduced from this paper's Supplementary Material.

Acknowledgments

The author used AI-assisted tools for formatting checks and code-verification support. The scientific claims, interpretations, and final manuscript decisions are the author's responsibility.

Declarations

Funding

This research received no external institutional funding.

Author Contributions

Steve Bico Mujjabi: conceptualization, methodology, formal analysis, software, validation, visualization, writing—original draft, and writing—review and editing.

Conflicts of Interest

The author declares no conflicts of interest.

References

References

- [1] James M. Bardeen, Brandon Carter, and Stephen W. Hawking. The four laws of black hole mechanics. *Communications in Mathematical Physics*, 31:161–170, 1973. doi: 10.1007/BF01645742.
- [2] Douglas Clowe, Marusa Bradac, Anthony H. Gonzalez, Maxim Markevitch, Scott W. Randall, Christine Jones, and Dennis Zaritsky. A direct empirical proof of the existence of dark matter. *Astrophysical Journal Letters*, 648:L109–L113, 2006. doi: 10.1086/508162. Weak-lensing and X-ray separation in the Bullet Cluster.
- [3] Stephen W. Hawking. Particle creation by black holes. *Communications in Mathematical Physics*, 43:199–220, 1975. doi: 10.1007/BF02345020.
- [4] Steve Bico Mujjabi. The Vacuum Equation of State: Deriving Torus Knot Self-Consistency from the Public CDFD Balance Law $\Psi_{LS} = \Phi / C$. Zenodo, 2026. URL <https://doi.org/10.5281/zenodo.21435391>. <https://doi.org/10.5281/zenodo.21435391>.
- [5] S. Perlmutter et al. Measurements of Ω and Λ from 42 high-redshift supernovae. *Astrophysical Journal*, 517:565–586, 1999. doi: 10.1086/307221.
- [6] Planck Collaboration, N. Aghanim, et al. Planck 2018 results. VI. cosmological parameters. *Astronomy & Astrophysics*, 641:A6, 2020. doi: 10.1051/0004-6361/201833910.
- [7] Adam G. Riess et al. Observational evidence from supernovae for an accelerating universe and a cosmological constant. *Astronomical Journal*, 116:1009–1038, 1998. doi: 10.1086/300499.
- [8] Vera C. Rubin and Jr. Ford, W. Kent. Rotation of the Andromeda nebula from a spectroscopic survey of emission regions. *Astrophysical Journal*, 159:379–403, 1970. doi: 10.1086/150317. Classic galaxy rotation-curve measurement motivating dark-matter phenomenology.