

Paper OOL-5: Oscillating Constraints and the Polymerization Ratchet: Environmental Cycling as Chemical Memory

Steve Bico Mujjabi, MD

Independent Researcher

Founder, Vura Labs

Kampala, Uganda

ORCID: 0009-0001-0556-5516

May 2026

Abstract

This manuscript addresses the water paradox in origin-of-life chemistry: aqueous environments enable transport and encounter, while condensation chemistry is opposed by hydrolysis in bulk water. CDFD reframes the problem as a temporal constraint problem. Wet–dry, freeze–thaw, tidal, geothermal, or other cycling environments can alternately mix reactants, concentrate them, and then preserve a fraction of the resulting products [1, 2]. The claim is not that one setting is proven to be universal. The claim is that an origin sequence needs some mechanism that periodically changes effective water activity, concentration, retention, and degradation pressure so that polymer-length or pattern-memory gains can accumulate.

Symbol Conventions

CDFD origins-of-life symbol convention.

Symbol	Meaning in Part II	Origins-of-life reading
Φ	Available driving flux	Redox, thermal, photochemical, proton/ion, or chemical potential drive supplied by the environment
C	Effective constraint or resistance	Mineral geometry, pore confinement, diffusion barrier, permeability, activation barrier, or maintenance burden
S	Responsive organization	Surface, boundary, catalytic network, coacervate, or protocellular structure that routes flux into retained function
M_s	Retained structural memory	Composition, sequence, topology, compartment state, mineral imprint, or boundary history that persists across renewal cycles
Ψ_s	Local adaptive ratio $(\Phi/C) \cdot S \cdot M_s$	Bookkeeping gauge for whether drive, constraint, responsiveness, and retention are co-present; not a universal constant
Λ_{life}	Integrated Life Number where used	Capstone surplus ratio for decay, near-critical proto-biology, and sustained self-maintenance

Greek-symbol convention.

Symbol	Local role	Interpretation rule
α	Accumulation or reinforcement coefficient	Defines how fast a constraint, pathway, or retained structure grows under load
β	Relaxation, decay, or washout coefficient	Defines recovery, loss, clearance, hydrolysis, or return toward baseline
γ	Coupling, diffusion, or redistribution coefficient	Defines spread across pores, surfaces, compartments, networks, or chemical phases
δ, ϵ	Perturbation, tolerance, or small offset scales	Used only as local thresholds, numerical guards, or experimental tolerances
η, λ, μ, ν	Efficiency, rate, baseline, or frequency terms	Meaning is set by the equation and subscript where they occur
ρ, σ, τ, ϕ	Density/correlation, conductance/noise, time-scale, or phase/field terms	Never universal constants unless a paper explicitly defines them that way
Ω, Λ	Domain/state space; Life Number where used	Ω names a modeled state space; Λ is reserved for life-number notation only where defined

1 Introduction

Papers 2–4 established geochemical redox, electron transport, and proton/ion coupling as the first concrete substrate. That is not enough. A system also needs a route from small molecules to longer-lived chemical patterns. Polymerization is difficult in bulk water because condensation reactions compete with hydrolysis and product loss. CDFD therefore asks what environmental topology can raise the forward polymerization opportunity without permanently destroying

transport. This includes wet–dry cycling, mineral-surface oligomerization, and activated nucleotide chemistry as different ways to change the same constraint landscape [6, 5].

2 Oscillating Adaptive Ratios

The shared adaptive ratio is

$$\Psi_s = \left(\frac{\Phi}{C} \right) S M_s. \quad (1)$$

In a cycling environment, the relevant quantities are time dependent:

$$\Psi_s(t) = \left(\frac{\Phi_{\text{chem}}(t)}{C_{\text{hydro}}(t)} \right) S(t) M_s(t). \quad (2)$$

Wet phases can increase encounter rates and transport. Dry or concentrated phases can raise monomer activity and reduce the effective hydrolytic penalty for selected reactions. Rehydration then tests whether any products survive long enough to seed another cycle.

3 The Polymerization Ratchet

A minimal water-volume cycle can be written

$$V_{\text{H}_2\text{O}}(t) = V_0 + \Delta V \cos(\omega t). \quad (3)$$

Here $V_{\text{H}_2\text{O}}$ is the available water volume, V_0 is the mean volume, ΔV is the cycle amplitude, and ω is the angular frequency. As $V_{\text{H}_2\text{O}}$ falls, monomer concentration rises and the system can enter a condensation-favoring window. As water returns, hydrolysis and dilution resume. A ratchet exists only if the retained product fraction after one cycle is greater than the lost fraction over many cycles:

$$M_s(t + nT) > M_s(t) \quad (4)$$

for at least some range of cycle periods $T = 2\pi/\omega$, temperatures, solute mixtures, surfaces, and degradation rates. The memory variable M_s should therefore be measured as retained product length, sequence/composition bias, catalytic activity, or surface-bound product fraction after rehydration, not as a single best-cycle yield.




Figure 1: Wet–dry polymerization-ratchet diagnostic. The diagnostic separates product formation from retained memory across several cycling regimes and adds a two-dimensional period–hydrolysis response surface. The model supports a conditional ratchet claim: activation, concentration, retention, and survival must align over repeated cycles.

4 Two-Dimensional Ratchet Window

The final release expands the companion script from a few named scenarios to a response surface. The scan varies the cycle period T and hydrolysis pressure h while keeping the feed chemistry fixed. For each grid point, the model runs repeated wet–dry substeps and records final retained memory. The retained region is not monotonic. Short periods fail because chemistry has too little time to complete; long periods can fail because products spend too much time in destructive phases; high hydrolysis erases the memory even when product is made.

This is the central CDFD lesson of Paper 5: the ratchet is a window, not a slogan. Saying “wet–dry cycling helps” is too weak and often misleading. The testable claim is that a finite region in $(T, h, \text{surface}, \text{pH}, \text{activation})$ space should outperform steady controls in retained product distribution after rehydration.

Status: Derived diagnostic

5 Recent Experimental Anchors

The cycling argument has strengthened since older wet–dry sketches. Heat-flow crack experiments show that geological non-equilibria can concentrate and purify many prebiotically relevant molecules while enhancing reaction yields [3]. Related work on phosphate shows that heat flows can solubilize apatite-derived phosphate and support trimetaphosphate formation under conditions relevant to prebiotic activation [4]. Meanwhile, aqueous-origin reviews emphasize that condensation polymer growth is not solved by saying “remove water”; useful chemistry must store activation energy in precursors, release it selectively, and survive hydrolysis pressure [7]. In CDFD terms, the ratchet is not only wet versus dry. It is the repeated alignment of activation, concentration, retention, and survival.

Status: Inferred

6 Validation Targets and Controls

A serious validation program must test cycling as a repeated-selection process, not as a one-shot concentration trick. The primary observables are (i) retained product mass after rehydration, (ii) length distribution or compositional bias across cycles, (iii) hydrolysis loss per wet phase, and (iv) whether products from one cycle seed the next cycle. Controls should include continuously wet mixtures, dry-only mixtures without remixing, inert-surface controls, mineral-surface variants, and identical feedstocks with different cycle periods.

The CDFD prediction is narrow: cycling improves memory only inside a window. Too little dehydration gives dilution and hydrolysis; too much dehydration traps or destroys reactants; too rapid cycling prevents reaction completion; too slow cycling allows degradation to dominate. A future experiment should therefore report the full response surface over T , temperature, ionic strength, pH, surface type, and activated precursor chemistry.

The strongest falsification would be a broad negative result: across realistic precursor chemistries and surfaces, no cycling protocol improves retained length, composition, catalytic activity, or seeding capacity relative to matched steady controls. A weaker but still important

falsification would show that cycling increases gross product only by producing chemically dead material that does not seed later cycles. Either outcome would force the Part II chain to replace wet–dry cycling with another mechanism that solves the same memory problem.

Status: Open experimental program

7 Claim Boundary

The claim is not that static submerged vents are impossible or that hot springs are mandatory. The claim is that a complete CDFD origin chain must include an environmental or material mechanism that solves the same functional problem:

- periodic concentration or compartmental concentration,
- partial dehydration or equivalent activation chemistry,
- product retention against washout,
- survival of useful products across the next destructive phase.

Wet–dry cycling is one strong implementation of that function; mineral surfaces, eutectic phases, gels, and compartment interiors may supply related effects.

8 Bridge to Paper 6

The output of this step is not life. It is a population of longer-lived patterns, oligomers, and surface-bound products. Paper 6 asks when such patterns become self-reinforcing through autocatalytic closure and chemical memory.

9 Conclusion

Oscillating constraints provide a plausible way to turn environmental fluctuation into retained chemical structure. In the Part II chain, this paper supplies the transition from geochemical throughput to memory-bearing chemistry while keeping the historical setting open to empirical testing.

10 Reproducibility Note

The companion script `supplementary-ool-05.py` writes the figure `outputs/paper_05/wet_dry_polymerization_ratchet.png` plus CSV/JSON summaries, including `outputs/paper_05/ratchet_response_surface.csv`. No notebook is required.

References

- [1] Bruce Damer and David Deamer. The hot spring hypothesis for an origin of life. *Astrobiology*, 20(4):429–452, 2020.

- [2] James P. Ferris, Aubrey R. Hill, Rihe Liu, and Leslie E. Orgel. Synthesis of long prebiotic oligomers on mineral surfaces. *Nature*, 381:59–61, 1996.
- [3] Thomas Matreux, Paula Aikkila, Bettina Scheu, Dieter Braun, and Christof B. Mast. Heat flows enrich prebiotic building blocks and enhance their reactivity. *Nature*, 628(8006):110–116, 2024.
- [4] Thomas Matreux, Almuth Schmid, Mechthild Rappold, et al. Heat flows solubilize apatite to boost phosphate availability for prebiotic chemistry. *Nature Communications*, 16:1809, 2025.
- [5] Bhavesh H. Patel, Claudia Percivalle, Dougal J. Ritson, Colm D. Duffy, and John D. Sutherland. Common origins of RNA, protein and lipid precursors in a cyanosulfidic protometabolism. *Nature Chemistry*, 7:301–307, 2015.
- [6] Matthew W. Powner, Béatrice Gerland, and John D. Sutherland. Synthesis of activated pyrimidine ribonucleotides in prebiotically plausible conditions. *Nature*, 459:239–242, 2009.
- [7] Daniel Whitaker and Matthew W. Powner. On the aqueous origins of the condensation polymers of life. *Nature Reviews Chemistry*, 8:817–832, 2024.