

Master Synthesis of Origins-of-Life CDFD: Tri-Regime Bioenergetics, Life Number, and Falsification Program

Steve Bico Mujjabi, MD*
VuraLabs, Kampala, Uganda
Independent researcher
ORCID: 0009-0001-0556-5516

August 2026

Abstract

This capstone manuscript synthesizes the revised twelve-paper Part II origins-of-life sequence. The central claim is not that life is proven inevitable, nor that a single molecule or environment is mandatory. The claim is that life-like persistence requires a coupled dependency graph: sustained input, electron transport, proton/ion coupling, retention, autocatalytic closure, information stability, replication with bounded error, defense against parasitic drains, and stabilization against overload. We define the Life Number Λ_{life} as a release-level bookkeeping gauge for transitions from decay ($\Lambda_{\text{life}} < 1$), to near-critical proto-biology ($\Lambda_{\text{life}} \approx 1$), to sustained self-maintenance ($\Lambda_{\text{life}} > 1$).

Keywords: tri-regime bioenergetics; life number; falsification; origins of life; systems synthesis.

Preprint note

This is a computational research article. The companion script produces numerical diagnostics reported in the Computational Results section. It is not a narrative review and does not claim historical proof of a unique abiogenesis pathway.

Symbol Conventions

See Paper 1, §Symbol Conventions, for the base CDFD and Greek-symbol tables. This capstone reserves Λ_{life} for the integrated Life Number and B_{stab} for stabilization burden, distinct from the local responsiveness symbol S . The capstone variables I_E , σ_e , σ_p , τ_{relax} , B_{stab} , and $E_{\text{maintenance}}$ are defined in §2 below.

*msbico@gmail.com

1 Introduction

The narrative mapped in this series is a dependency graph rather than a rigid geological clock. Each step supplies a function that the next step uses or stresses.

1. **Thermodynamic framing:** sustained disequilibrium creates the possibility of organized dissipation.
2. **Geochemical crucibles:** mineral redox, mixed-valence transport, and interfacial water localize electron and proton/ion work.
3. **Oscillating constraints:** cycling environments provide a route to retained polymers and pattern memory.
4. **Autocatalytic closure:** reaction loops begin to reinforce their own production.
5. **Topological boundaries:** mineral pores, films, coacervates, and soft compartments raise retention and create adaptive surfaces.
6. **Protocell integration:** copying, energy flow, and container persistence must covary.
7. **Parasitic threshold:** successful copying creates exploitative failure modes and pressure for boundary logic.
8. **Photochemical expansion:** light or other stronger inputs can expand the energy channel without being required at the first origin step.
9. **Master synthesis:** the chain is evaluated as a tri-regime persistence criterion.

2 Notation and Symbol Discipline

Part II uses CDFL symbols as measurement targets, not decoration. The local CDFD adaptive ratio remains

$$\Psi_s = \left(\frac{\Phi}{C} \right) SM_s, \tag{1}$$

where Φ is local drive, C is effective resistance or constraint, S is responsive organization, and M_s is retained structural memory. In this capstone we reserve Λ_{life} for the integrated Life Number. We reserve B_{stab} for stabilization burden so that it is not confused with the local responsiveness symbol S .

The main capstone variables are:

Symbol	Meaning
I_E	Usable environmental energy input after coupling to a chemical or mineral pathway.
σ_e	Effective electron-transport conductance.
σ_p	Effective proton or ion coupling conductance.
τ_{relax}	Time over which structure persists after perturbation.
B_{stab}	Stabilization burden from radical load, heat, photodamage, hydrolysis, or uncoupled side reactions.
$E_{\text{maintenance}}$	Energy cost of repair, boundary maintenance, and retained organization.
Λ_{life}	Integrated surplus ratio for decay, near-critical, and sustained regimes.

3 Tri-Regime Bioenergetics

All of these transitions motivate a shared modelling question: does usable coupled throughput exceed the cost of maintaining organization? We define the Life Number (Λ_{life}) as an integrated bookkeeping gauge constructed from the local adaptive ratio:

$$\Lambda_{\text{life}} = \frac{I_E \cdot \sigma_e \cdot \sigma_p \cdot \tau_{\text{relax}}}{B_{\text{stab}} \cdot E_{\text{maintenance}}}. \quad (2)$$

The reference value one and the labels below are conventions of this toy diagnostic, not calibrated biological boundaries or demonstrated historical transition points. They acquire empirical meaning only after variables, units, and decision thresholds have been independently specified for a stated system.

The classical CDFD parameters map to bioenergetic requirements as follows:

- **Flux** (Φ): represented by energy input (I_E) captured through geochemical, photochemical, or hydrothermal means.
- **Constraint** (C): The maintenance energy ($E_{\text{maintenance}}$) required to repair or preserve boundaries, catalysts, polymers, and inherited organization.
- **Responsiveness** (S): structural elasticity and routing capacity, amplified by electron transport (σ_e) and proton/ion coupling (σ_p).
- **Memory** (M_s): Reflected in the relaxation time (τ_{relax}) of the structural network. Stable, high-memory polymers allow the system to maintain its configuration without instantly degrading back into chaos.
- **Stabilization burden** (B_{stab}): The damage and dissipation load that must be buffered before extra input becomes useful work.

3.1 The Three Regimes of Emergence

1. **Decay-dominated** ($\Lambda_{\text{life}} < 1$): Energy flows, but structures cannot maintain themselves against hydrolysis, dilution, side reactions, and transport loss.

2. **Near-critical proto-biological** ($\Lambda_{\text{life}} \approx 1$): A compartmental or surface-bound system marginally maintains itself and is sensitive to cycling, feedstock, and boundary perturbations.
3. **Sustained/self-maintaining** ($\Lambda_{\text{life}} > 1$): Coupled input, transport, retention, memory, and stabilization leave surplus capacity for repair, growth, division, or complexity accumulation.

4 The Corrected Synthesis

The Part II sequence is clearest when written as a dependency graph rather than a substance list. The earliest transition does not require literal chlorophyll, literal melanin, or any single privileged molecule. It requires a system that can keep useful work localized long enough for chemistry to become self-reinforcing.

The corrected one-line thesis is:

Life begins when constrained flux becomes coupled, retained, self-amplifying, information-stabilizing, and repairable above maintenance cost.

Earlier manuscripts in this programme establish the mineral, chemical, protocellular, replication, and boundary-defense side of that transition [19, 9, 7, 22]. Related photochemical work clarifies that photochemical energy capture expands the energy-input channel but should not be confused with the first emergence of proto-life. This paper joins both tracks into the four-channel/three-regime model.

Relation to the closest existing thermodynamic-dissipation literature. This thesis sits inside an established lineage rather than standing apart from it. Prigogine’s dissipative-structures framework [16] is the origin of the dissipative-structuring language used throughout this series (see Paper 1). England’s statistical-physics account of dissipation-driven adaptation [3] makes a closely related quantitative claim using different formal machinery — that driven systems can be statistically favored toward configurations that dissipate more of the available work, provided those configurations also persist. Michaelian’s thermodynamic dissipation theory [11, 12] proposes a directly competing account centered on UV photon dissipation as the primary selective pressure at the origin of life, rather than the more general coupled-channel account developed across this series. Where this synthesis differs from all three is in the specific multi-channel (I_E , σ_e , σ_p , τ_{relax} , B_{stab} , $E_{\text{maintenance}}$) bookkeeping and the twelve-stage dependency ladder — not in the underlying thermodynamic intuition, which these three sources established first.

5 Functional Ontology

Definition 1 (Functional Material Class). *A functional material class is any set of substrates that performs a required dynamical role in the CDFD Part II model, regardless of whether the modern exemplar existed at the origin of life.*

The main classes are:

Role	Symbol	Function	Example implementations
Energy input	I_E	Supplies usable drive	Fe-S redox, electron donors, photoredox minerals, pigments
Electron transport	σ_e	Moves reducing equivalents	Fe-S clusters, magnetite-like phases, quinones, ETC cofactors
Proton/ion coupling	σ_p	Couples charge flow to work	Interfacial water, pH gradients, proton relays, membranes
Stabilization burden	B_{stab}	Penalizes damage/dissipation	Radical load, heat, hydrolysis, photodamage, repair burden

Modern names are mnemonic only. “Chlorophyll-like” means energy capture. “Magnetite-like” means distributed electron conductance. “Water-like” means proton or ion coherence. “Melanin-like” means dissipation buffering.

The model is wrong if these labels are read as mandatory historical substances.

6 Revised Part II Flow

Stage	Main bottleneck	CDFD resolution	Current paper
1	No common language for persistence	Thermodynamic notation defines flux, constraint, responsiveness, and memory	Thermodynamic mandate
2	Redox work is not localized	Fe-S redox interfaces localize chemical drive	Fe-S scaffolds
3	Redox sites are isolated	Mixed-valence networks distribute electron flow	Magnetite networks
4	Electrons lack proton/ion partner	Structured interfacial fluids couple charge channels	Proton/ion coupling
5	Products wash out or hydrolyze	Cycling raises concentration, activation, and retained product memory	Polymerization ratchet
6	Chemistry is not self-reinforcing	Autocatalytic closure beats diffusive loss	Autocatalytic closure
7	Patterns decay or scramble	Aromatics, alphabets, and chiral amplification improve pattern persistence	Alphabets/homochirality
8	Chemistry lacks movable retention	Mineral pores, interfaces, coacervates, and vesicles create adaptive boundaries	Boundaries/compartments
9	Replication loses information	Flux-dependent fidelity bounds error	Protocell/error threshold

Stage	Main bottleneck	CDFD resolution	Current paper
10	Exploiters drain copied systems	Specificity and boundary logic suppress parasitic access	Parasitic threshold
11	External energy input can expand	Photochemical/material amplifiers raise I_E when coupled and buffered	Photochemical capture
12	Claims need one falsifiable synthesis	Λ_{life} organizes decay, near-critical, and sustained regimes	Tri-regime synthesis

Table 1: Corrected dependency graph for the Part II origins-of-life sequence. The order is logical rather than a rigid geological timeline.

This table restores the flow: energy capture comes before overload stabilization, while modern plant biochemistry remains a later example.

7 Computational Results

The companion master-synthesis diagnostic is reported in Figure 1.




Figure 1: Capstone numerical diagnostic. The left panel shows the three-regime Life Number interpretation. The middle panel shows how environmental cycling can move a localized system across the persistence threshold. The right panel illustrates LLPS-like heterogeneity as a compartment-forming constraint field.

8 Notation and independence

Part II uses CDFD/CDFL symbols as a coarse-grained bookkeeping layer for origin-of-life dependency order. Evaluation of this paper does not require accepting Part I vacuum-physics

claims; chemical and geological arguments stand on their own.

9 The Four-Channel Throughput Model

Let I_E be usable input energy after environmental coupling. It may come from geochemical redox gradients, electron donors, mineral photochemistry, or mature photosynthetic pigments. Let $\bar{\sigma}_e$ be effective electron transport conductance, $\bar{\sigma}_p$ effective proton/ion coupling, and B_{stab} the stabilization burden.

The mean usable throughput is:

$$\bar{J} = I_E \frac{\bar{\sigma}_e \bar{\sigma}_p}{B_{\text{stab}}}. \quad (3)$$

The weakest-link constraint remains:

$$J_{\text{max}} \leq \min(I_E, \bar{\sigma}_e, \bar{\sigma}_p, 1/B_{\text{stab}}). \quad (4)$$

Equation (4) is why the series cannot be reduced to one favored substance. A better energy absorber does not help if electron transport is absent. Better electron transport does not help if proton coupling is incoherent. Better coupling does not help if products wash out. Better flux does not help if overload destroys the system.

10 The Three Regimes and the Life Number

Definition 2 (Life Number).

$$\Lambda_{\text{life}} = \frac{I_E \bar{\sigma}_e \bar{\sigma}_p \tau_{\text{relax}}}{B_{\text{stab}} E_{\text{maintenance}}}, \quad (5)$$

Here τ_{relax} is the characteristic time for preserved organization. $E_{\text{maintenance}}$ is the energetic cost of maintaining coherent structure over that interval.

Λ_{life} is treated as a surplus ratio whose numerator is usable coupled throughput retained over time and whose denominator is the cost of not falling apart. Whether the ratio is genuinely dimensionless after a full unit analysis of I_E , σ_e , σ_p , τ_{relax} , B_{stab} , and $E_{\text{maintenance}}$ remains open (see Limitations).

Condition	Regime	Interpretation
$\Lambda_{\text{life}} < 1$	Decay-dominated	Dissipation, dilution, and damage exceed useful work
$\Lambda_{\text{life}} \approx 1$	Near-critical/proto-biological	Marginal self-maintenance; perturbation-sensitive
$\Lambda_{\text{life}} > 1$	Sustained/self-maintaining	Surplus supports repair, replication, and complexity accumulation

The phrase “tri-regime” therefore refers to three dynamical regimes, not three materials.

11 Relation to $\Psi_s = (\Phi/C)SM_s$

The local CDFD adaptive ratio is:

$$\Psi_s = \left(\frac{\Phi}{C}\right) SM_s. \quad (6)$$

In a single-channel system, higher driving flux Φ , lower effective constraint C , stronger responsiveness S , or longer memory M_s increases the chance of persistent structure. The Life Number is the multi-channel generalization:

$$\Psi_s \longrightarrow \Lambda_{\text{life}} = \frac{I_E \bar{\sigma}_e \bar{\sigma}_p \tau_{\text{relax}}}{B_{\text{stab}} E_{\text{maintenance}}}. \quad (7)$$

The mapping is not a claim that all channels are literally identical. It says that both quantities compare usable drive and retained organization against constraining cost. Ψ_s is the local adaptive form; Λ_{life} is the integrated bioenergetic form.

12 Geochemistry to Biology

12.1 Working structural comparison

The following is a modelling comparison, not a theorem, derivation, or identity claim. Similar low-order differential equations can describe many systems with production, loss, and spatial transport. Shared notation or operator form does not establish that geochemistry and biology have one mechanism, one ontology, or a common natural law.

For a geochemical system, write:

$$\frac{\partial C_{\text{geo}}}{\partial t} = \alpha_{\text{geo}} |\Phi_{\text{geo}}| - \beta_{\text{geo}} C_{\text{geo}} + \gamma_{\text{geo}} \nabla^2 C_{\text{geo}}. \quad (8)$$

Here C_{geo} is mineral or transport constraint, Φ_{geo} is environmental forcing, α_{geo} captures precipitation or channel formation, β_{geo} captures dissolution or degradation, and γ_{geo} captures spatial smoothing.

For a biological or proto-biological maintenance system, write:

$$\frac{\partial C_{\text{bio}}}{\partial t} = \alpha_{\text{bio}} |\Phi_{\text{bio}}| - \beta_{\text{bio}} C_{\text{bio}} + \gamma_{\text{bio}} \nabla^2 C_{\text{bio}}. \quad (9)$$

Here C_{bio} is maintenance burden, damage state, or adaptive constraint; Φ_{bio} is metabolic throughput; and the remaining coefficients describe repair, degradation, and spatial propagation.

The equations share the same operator structure under the substitution

$$\begin{aligned} C_{\text{geo}} &\leftrightarrow C_{\text{bio}}, \\ \Phi_{\text{geo}} &\leftrightarrow \Phi_{\text{bio}}, \\ \alpha_{\text{geo}}, \beta_{\text{geo}}, \gamma_{\text{geo}} &\leftrightarrow \alpha_{\text{bio}}, \beta_{\text{bio}}, \gamma_{\text{bio}}. \end{aligned}$$

The equations have a formally similar production–loss–diffusion shape under this chosen correspondence. That similarity is useful only as a working bookkeeping analogy: it can suggest measurements and comparisons, but it does not show that a biological maintenance process is a refined geochemical process or that either equation is sufficient for origin-of-life history.

Working implication. One possible modelling hypothesis is that a localized chemical system must maintain coupled input, retention, repair, and bounded error against dilution and damage. Λ_{life} is only a coarse bookkeeping gauge for organizing such comparisons; it neither proves a historical origin pathway nor defines life.

13 Where Photochemistry and Stabilization Belong

Photochemical capture is an energy-input expansion, not the first mandatory step. In early settings, I_E may be supplied by vent redox gradients, Fe^{2+} , H_2 , H_2S , pH gradients, or mineral surface chemistry. Photochemical materials become important when they allow the system to harvest light and decouple from a purely local geochemical source.

Modern chlorophyll illustrates the high-performance endpoint:

$$\Phi_{\text{RC}} = \eta_{\text{transfer}} N_{\text{antenna}} \Phi_{\text{photon-per-pigment}}. \quad (10)$$

But the prebiotic claim is broader: any material that increases I_E while remaining coupled to $\bar{\sigma}_e$ and $\bar{\sigma}_p$ can push Λ_{life} upward. This includes weaker and earlier candidates such as photoredox minerals, porphyrinoid precursors, flavins, quinones, and other chromophores.

Stabilization follows energy capture in the causal order. Once I_E rises, damage risk rises too:

$$\Psi_{s,\text{overload}} = \frac{I_E}{C_{\text{use}} + C_{\text{diss}}}. \quad (11)$$

If the usable and dissipative capacities cannot absorb input, side reactions, radical chemistry, photobleaching, and thermal disorder increase S .

Melanin-like language means any process that reduces effective S without collapsing useful channels. Modern examples include carotenoids, non-photochemical quenching, melanins, polyphenols, enzymatic antioxidants, and repair cycles. Prebiotic analogues could include disordered aromatic polymers, mineral screens, sacrificial redox buffers, or radical sinks. Literal melanin is not required for the earliest origin.

Feedstock provenance is orthogonal to the Life Number. Exogenous aromatic delivery can raise the available aromatic reservoir, but it does not itself raise Λ_{life} unless the material is retained, coupled, stabilized, and integrated into closure or replication gates. In the series logic, meteoritic organics are therefore a possible supply term, not a substitute for localization, transport, heredity, or overload control.

14 Phosphate Quantization and Phase Separation

Two later transitions fit naturally into the same model.

First, phosphate chemistry discretizes energy handling:

$$E_{\text{packet}} = n\Delta E_P, \quad n \in \mathbb{Z}^+. \quad (12)$$

Discrete packets reduce uncontrolled dissipation and make reaction coupling more precise. In Eq. (5), this can lower B_{stab} and improve effective $\bar{\sigma}_p$.

Second, phase separation and soft compartments provide dynamic localization before full lipid-cell architecture. LLPS, coacervates, gels, and mineral-organic films all increase residence time and co-localization [4, 22]. These mechanisms sit between mineral pores and modern membranes in the dependency graph.

15 Additional Origins-of-Life Transitions Now Captured

The current twelve-paper spine should explicitly preserve several ideas that are easy to lose when separate mechanism threads are merged.

15.1 Mineral to soft compartment

Mineral pores localize chemistry first, but soft compartments matter later because they can grow, divide, deform, and inherit internal composition. In CDFD notation this is a transition from externally imposed constraint to partially self-maintained constraint:

$$C_{\text{boundary}} = C_{\text{mineral}} + \eta(t)C_{\text{organic}} + \chi(t)C_{\text{amphiphile}}. \quad (13)$$

When η and χ increase, the boundary becomes less purely geological and more proto-biological.

15.2 Phosphate and energy packets

Continuous flux is not enough for heredity. The system also needs reusable energetic packets or coupling currencies. Phosphate chemistry is one modern endpoint, but the broader claim is quantized coupling: energy must become deliverable to specific reactions instead of remaining diffuse heat or bulk redox disequilibrium.

15.3 Environmental cycling

Drying, rewetting, freezing, heating, vent pulsation, and day–night illumination can act as external clocks. A system that survives only at a steady optimum is fragile. A more realistic proto-biological candidate should persist through cycles:

$$\Psi_{s,\text{cycle}} = \prod_{k=1}^N \frac{I_E^{(k)} \sigma_e^{(k)} \sigma_p^{(k)} \tau_k}{B_{\text{stab}}^{(k)} E_{\text{maintenance}}^{(k)}}. \quad (14)$$

The product form is schematic, but it captures the idea that repeated sub-events can either accumulate resilience or erase the system.

15.4 Material agnosticism

Named materials are examples. Fe-S, magnetite, structured water, chlorophyll, and melanin are not mandatory steps. The required sequence is functional: input, transport, coupling, retention, closure, information, replication, capture expansion, and overload stabilization.

16 Reference-Layer Synthesis

The expanded bibliography clarifies that CDFD is not choosing a single traditional origin scenario. It is integrating constraints from several partially compatible literatures and from field-level reviews that treat origin-of-life research as a plural, planetary problem [15, 20]:

- alkaline hydrothermal vents provide a fed redox-gradient and mineral-compartment architecture [19, 9, 21];
- critiques of vent-gradient sufficiency force explicit coupling tests instead of narrative certainty [6];
- hot-spring and wet-dry settings emphasize cycling, concentration, and dehydration chemistry [2];
- cyanosulfidic photochemistry links nucleotide, amino-acid, and lipid precursor chemistry under shared environmental assumptions [14];
- model protocells show that compartments, growth, division, and nonenzymatic replication can be experimentally coupled [4, 8, 1];
- coacervates and phase-separated droplets provide non-lipid soft-compartment routes [10, 13];
- UV-environment studies define which surface photochemistry is plausible [18, 17].

The series therefore reads as a constraint map across environments rather than as a single-site narrative.

17 Master Falsification Matrix

The master paper states what would weaken or falsify each layer:

Layer	CDFD expectation	Strong disconfirming result
Mineral redox	Fed mineral interfaces localize usable work	No product or redox coupling advantage over bulk controls
Transport	Mixed-valence networks improve source-sink coupling	Conductive phases do not alter downstream chemistry
Proton/ion coupling	Electron and proton/ion currents must co-localize	Robust proto-metabolism without local charge compensation
Compartments	Retention changes reaction-network fate	Open dilution performs as well as confined/cycled systems
Information	Durable alphabets must persist through noise	No plausible molecular class can carry persistent patterns
Replication	Bounded error is required for heredity	Long templated heredity persists below error-threshold constraints
Photochemistry	Light helps only when coupled and buffered	Photochemical input only damages, never expands usable throughput

This matrix keeps the series ambitious but falsifiable.

18 Validation Program and Future Experiments

The framework becomes scientifically useful only if it generates experiments that can fail. The following program translates the twelve-paper chain into test classes. Each test must pre-register the variable being changed, the matched controls, and the failure condition; otherwise the CDFD notation risks becoming descriptive rather than predictive.

1. **Matched mineral-reactor tests.** Build paired fed reactors with identical pH, temperature, feedstock, residence time, and surface area. Vary only the connected Fe-S or mixed-valence bridge. The expected signal is not “life,” but a reproducible change in downstream redox state, product distribution, and persistence time relative to disconnected, inert, or poisoned-surface controls.
2. **Electron-proton co-localization tests.** Measure redox current, pH microgradients, ion flux, and product formation in the same pore network. CDFD predicts that chemistry improves when electron and proton/ion channels overlap in space and time, not merely when either channel exists alone. A failure condition is robust product improvement in geometries where charge channels are deliberately separated.
3. **Cycling-retention tests.** Compare wet-dry, thermal-gradient, freeze-thaw, and continuously wet controls using the same monomer pool. The score is the retained product distribution after many cycles, including hydrolysis losses, not the best single-cycle yield. The response should be windowed over period, amplitude, temperature, surface, and activation chemistry.
4. **Boundary-covariance tests.** In coacervates, vesicles, gels, or mineral-organic films, ask whether internal chemistry changes boundary growth, division, permeability, or survival. A container becomes protocellular only when internal state and boundary fate covary.

Empty-boundary controls and free-template controls are required to show that selection is acting on the coupled unit.

5. **Autocatalytic-closure tests.** Track reaction-network composition through serial transfer or repeated environmental cycles. The required signal is not a transient product spike, but mutual reinforcement: products from one cycle increase the probability of regenerating the same network in later cycles. Failure is amplification without retained network identity.
6. **Parasitic-drain tests.** Add noncontributing templates or catalytic freeloaders to compartmental copying systems. The CDFD expectation is collapse unless specificity, exclusion, degradation, or boundary partitioning is introduced. The falsifiable readouts are host fraction, compartment survival, leakage, and host-boundary covariance after serial passage.
7. **Photochemical-overload tests.** Raise light or photoredox input while independently varying electron transport, proton/ion coupling, and buffering materials. Useful capture should appear only in the coupled-and-buffered region; otherwise light should increase damage, side products, or bleaching. The control is total absorption without productive charge coupling.
8. **Cross-environment Life Number ranking.** Estimate I_E , σ_e , σ_p , τ_{relax} , B_{stab} , and $E_{\text{maintenance}}$ for vents, ponds, hot springs, ice/eutectic settings, mineral films, and soft compartments. The model predicts ranking by coupled surplus, not by a preferred location name. Ranking fails if uncoupled high-input environments outperform coupled environments after maintenance and damage costs are included.

These tests would not prove one historical path by themselves. They would identify which environments solve the required gates and which attractive stories fail under controlled comparison. The practical standard is therefore comparative: same feedstock, same measurement window, one changed CDFD channel, and an explicit criterion for weakening the claim.

19 CDFD Runtime Discovery Triage

The public CDFD Runtime is useful here only if it sharpens experiments rather than inflating claims. The root-level discovery sweep `experiments/notebooks/discovery_ool_phase.py` was rerun for this release using the current ontology adapter. It scanned a 50×50 grid of energy input and constraint burden and wrote `experiments/outputs/discovery_ool_phase_results.h5`. The Life Number range was 0.144 to 28.515, with four grid points in the near-critical window $0.98 \leq \Lambda_{\text{life}} \leq 1.02$.

That result should be treated as triage, not proof. It says the current runtime ontology has a narrow transition band that can be used to select follow-up parameter windows for paper-local models and wet-lab proposals. It does not say that those four grid points correspond to real prebiotic settings. The near-critical window’s own width (0.98–1.02) has *not* been tested for sensitivity to alternative reasonable choices — the same open item flagged in Paper 1 for the

0.85–1.15 window applies here; a sweep across several window widths, confirming the same four (or a similar number of) near-critical grid points recur qualitatively, is identified as required future work rather than already performed. The correct publication rule is: runtime discovery outputs may nominate candidate regimes; manuscript claims require paper-local scripts, cited chemistry, and falsifiable controls.

20 Environmental Cycling as Boundary Condition

Static chemistry is not enough. Prebiotic systems are driven by environmental cycles that repeatedly create, erase, and re-create gradients. The model therefore treats the environment as a time-dependent boundary condition:

$$T(t) = T_0 + \Delta T \sin(\omega_T t), \quad (15)$$

$$W(t) = W_0 + \Delta W \sin(\omega_W t), \quad (16)$$

$$E_{\text{redox}}(t) = E_0 + \Delta E \sin(\omega_E t), \quad (17)$$

$$J_{\text{ext}}(t) = J_0 + A \sin(\omega_J t). \quad (18)$$

Thermal cycling changes rates and solubility. Wet–dry cycling can favor condensation during low-water phases and mobility during wet phases. Redox cycling alternates available electron donors and acceptors. Hydrodynamic cycling injects nutrients and removes waste.

In CDFD notation these forcings modulate constraints:

$$C(x, t) = C_0(x) + C_{\text{env}}(x, t), \quad (19)$$

and therefore modulate throughput:

$$\bar{J}(t) = I_E(t) \frac{\bar{\sigma}_e(t) \bar{\sigma}_p(t)}{B_{\text{stab}}(t)}. \quad (20)$$

Without forcing, many chemical systems relax toward equilibrium. With repeated forcing, systems can be tested over cycles: compartments form and dissolve, polymers synthesize and hydrolyze, and only more persistent networks remain coherent across the cycle.

21 Autonomy, Persistence, and Death Thresholds

The transition toward individuality is not merely high flux, but persistence of a localized flux loop through environmental change. For a localized domain Ω , define a flux-to-constraint efficiency proxy:

$$\Psi_{s, \Omega} = \frac{\int_{\Omega} |\mathbf{J}| dV}{\int_{\Omega} C dV}. \quad (21)$$

Early domains are expected to be phase-locked to the environment. A more autonomous domain stores energy and maintains internal circulation after external forcing declines.

One can measure this with an autonomy index:

$$A_{\Omega} = 1 - |\text{corr}(\Psi_{s,\Omega}(t), J_{\text{ext}}(t))|. \quad (22)$$

This is only a proposed diagnostic. $A_{\Omega} \approx 0$ means the domain simply follows the environment; $A_{\Omega} \rightarrow 1$ means the internal flux state has partly decoupled from external oscillation.

A localized system dies, in this model, when internal circulation and maintenance fall below the losses needed to preserve identity:

$$\Gamma_{\Omega} = \oint_{\partial\Omega} \mathbf{J} \cdot d\mathbf{l} \leq \Gamma_{\text{crit}}. \quad (23)$$

Equivalently, the Life Number falls below unity for longer than the relaxation time:

$$\Lambda_{\text{life}}(t) < 1 \quad \text{for} \quad \Delta t \gtrsim \tau_{\text{relax}}. \quad (24)$$

This provides a simulation criterion for persistence and collapse without claiming that a toy domain is already biologically alive.

22 Mature Plant Systems as Later Examples

Plants, algae, and cyanobacteria are not the origin mechanism; they are mature demonstrations of the same logic. Their channels can be mapped as:

Channel	Early/geochemical implementation	Mature phototroph implementation
I_E	Redox donors, mineral photochemistry	Chlorophyll antennae, photosystems
$\bar{\sigma}_e$	Fe-S/magnetite-like conductors	Plastoquinone, cytochrome b_6f , ferredoxin
$\bar{\sigma}_p$	Mineral-water pH gradients	Thylakoid proton motive force
S	Radical sinks, mineral screens, polymers	Carotenoids, NPQ, repair, antioxidants

C_3 , C_4 , and CAM are therefore downstream examples: spatial and temporal solutions to carbon, water, light, and photorespiration constraints.

23 Predictions and Falsifiability

The corrected model is falsifiable at multiple levels:

1. Systems with higher coupled $I_E \bar{\sigma}_e \bar{\sigma}_p / B_{\text{stab}}$ should maintain organization longer than systems with high input but poor coupling.
2. Mineral or soft compartments that increase retention should shift autocatalytic networks closer to closure thresholds.

3. Photochemical materials should raise Λ_{life} only when electron and proton/ion channels are available; uncoupled absorption should increase damage or dissipation instead.
4. Stabilizing materials should improve survival most strongly under high-input overload, not under starvation.
5. Replacing chlorophyll-like or melanin-like examples with different materials should preserve the model if the same functions are preserved.

The model would be weakened if self-maintaining replication-like chemistry is demonstrated without persistent energy input, without co-localized charge coupling, without retention, and without any mechanism controlling damage or error.

24 Limitations

- Λ_{life} is currently a coarse-grained diagnostic, not a calibrated universal constant.
- No dimensional analysis has been performed showing Λ_{life} is genuinely unitless — this is identified as required future work: a unit table for I_E , σ_e , σ_p , τ_{relax} , B_{stab} , and $E_{\text{maintenance}}$, done once for the series, is needed to confirm the ratio cancels to dimensionless as claimed.
- The material implementation space is broad; assigning historical priority requires experiments and geochemical evidence.
- The structural comparison above is a bookkeeping analogy, not evidence that geochemistry and biology are identical or that either shares a unique governing equation.
- Photochemical routes may be late, parallel, or environment-specific rather than universal.
- Stabilization burden B_{stab} bundles many processes that should eventually be decomposed.
- The near-critical window widths used in this paper and Paper 1 (0.98–1.02 and 0.85–1.15 respectively) have not been sensitivity-tested against alternative widths.

25 Conclusion

The CDFD origins-of-life framework is strongest when it is functional. The series should be read as:

$$\begin{aligned}
 &\text{redox work} \rightarrow \text{transport coupling} \rightarrow \text{cycling retention} \\
 &\quad \rightarrow \text{closure} \rightarrow \text{information persistence} \rightarrow \text{bounded replication} \\
 &\quad \rightarrow \text{parasitic defense} \rightarrow \text{photochemical expansion} \rightarrow \text{overload stabilization.}
 \end{aligned} \tag{25}$$

Chlorophyll enters as a mature and powerful energy-input amplifier, not as a required first molecule. Melanin enters as one possible stabilization exemplar, not as a required origin material. This preserves the original intuition while making the papers clearer and more falsifiable: CDFD predicts the coupled functions a living transition must solve, then leaves the historical

material assignment open to evidence — an evidence base that, per the corrections in this release, is now explicitly positioned relative to the Prigogine/England/Michaelian dissipation-based origin-of-life literature [16, 3, 11, 12] rather than presented as a standalone framework.

26 Reproducibility Note

Supplementary material should stay script-first. Paper 1 now has a dependency-gate diagnostic, so every active Part II paper has an explicit script or output path. Papers with generated figures should store only matching script outputs under `outputs/paper_XX/`. Do not include placeholder notebooks solely to preserve one-to-one numbering.

Related literature

Part II is a self-contained origins-of-life dependency graph. Shared CDFD/CDFL symbols are bookkeeping only and do not require Part I physics claims. Adjacent mechanism papers in this archive are cited where used. Additions required by the missing-references review: England [3]; Michaelian [11, 12]; Prigogine and Nicolis [16]; Hordijk and Steel [5] (RAF set theory, most directly Paper 6).

Competing Interests

The author declares no competing interests.

Funding

This research received no external funding.

Data and Code Availability

Companion scripts, generated figures, and numerical outputs for this manuscript are included in the public CDFD Part II release archive (<https://github.com/bampita-bico/CDFD-Part-II-Release>; Zenodo DOI 10.5281/zenodo.20264779). See the Reproducibility Note above for the paper-local script path.

Author Contributions

Conceptualization, formal analysis, computational diagnostics, and manuscript preparation: Steve Bico Mujjabi

References

- [1] Katarzyna Adamala and Jack W. Szostak. Nonenzymatic template-directed RNA synthesis inside model protocells. *Science*, 342:1098–1100, 2013.

- [2] Bruce Damer and David Deamer. The hot spring hypothesis for an origin of life. *Astrobiology*, 20(4):429–452, 2020.
- [3] Jeremy L. England. Statistical physics of self-replication. *Journal of Chemical Physics*, 139(12):121923, 2013.
- [4] Martin M. Hanczyc, Shelly M. Fujikawa, and Jack W. Szostak. Experimental models of primitive cellular compartments: encapsulation, growth, and division. *Science*, 302:618–622, 2003.
- [5] Wim Hordijk and Mike Steel. Detecting autocatalytic, self-sustaining sets in chemical reaction systems. *Journal of Theoretical Biology*, 227(4):451–461, 2004.
- [6] J. Baz Jackson. Natural pH gradients in hydrothermal alkali vents were unlikely to have played a role in the origin of life. *Journal of Molecular Evolution*, 83:1–11, 2016.
- [7] Nick Lane, John F. Allen, and William Martin. How did LUCA make a living? chemiosmosis in the origin of life. *BioEssays*, 32:271–280, 2010.
- [8] Sheref S. Mansy, Jason P. Schrum, Mathangi Krishnamurthy, et al. Template-directed synthesis of a genetic polymer in a model protocell. *Nature*, 454:122–125, 2008.
- [9] William Martin and Michael J. Russell. On the origin of biochemistry at an alkaline hydrothermal vent. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 362(1486):1887–1925, 2007.
- [10] Muneyuki Matsuo and Kensuke Kurihara. Proliferating coacervate droplets as the missing link between chemistry and biology in the origins of life. *Nature Communications*, 12:5487, 2021.
- [11] Karo Michaelian. Thermodynamic dissipation theory for the origin of life. *Earth System Dynamics*, 2:37–51, 2011.
- [12] Karo Michaelian. Microscopic dissipative structuring and proliferation at the origin of life. *Heliyon*, 3(10):e00424, 2017.
- [13] Karina K. Nakashima, Merlijn H. I. van Haren, Alain A. M. André, et al. Active coacervate droplets are protocells that grow and resist Ostwald ripening. *Nature Communications*, 12:3819, 2021.
- [14] Bhavesh H. Patel, Claudia Percivalle, Dougal J. Ritson, Colm D. Duffy, and John D. Sutherland. Common origins of RNA, protein and lipid precursors in a cyanosulfidic protometabolism. *Nature Chemistry*, 7:301–307, 2015.
- [15] Martina Preiner et al. The future of origin of life research: Bridging decades-old divisions. *Life*, 10(3):20, 2020.
- [16] Ilya Prigogine and Grégoire Nicolis. *Self-Organization in Nonequilibrium Systems*. Wiley, New York, 1977.

- [17] Sukrit Ranjan, Charles L. Kufner, Georgina G. Lozano, Zoe R. Todd, Arda Haseki, and Dimitar D. Sasselov. UV transmission in natural waters on prebiotic Earth. *Astrobiology*, 22(3):242–262, 2022.
- [18] Sukrit Ranjan and Dimitar D. Sasselov. Constraints on the early terrestrial surface UV environment relevant to prebiotic chemistry. *Astrobiology*, 17(3):169–204, 2017.
- [19] Michael J. Russell, Roy M. Daniel, Allan J. Hall, and John A. Sherringham. A hydrothermally precipitated catalytic iron sulphide membrane as a first step toward life. *Journal of Molecular Evolution*, 39:231–243, 1994.
- [20] Dimitar D. Sasselov, John P. Grotzinger, and John D. Sutherland. The origin of life as a planetary phenomenon. *Science Advances*, 6:eaax3419, 2020.
- [21] Victor Sojo, Barry Herschy, Alexandra Whicher, Eloi Camprubí, and Nick Lane. The origin of life in alkaline hydrothermal vents. *Astrobiology*, 16(2):181–197, 2016.
- [22] Jack W. Szostak. The origins of cellular life. *Cold Spring Harbor Perspectives in Biology*, 2:a002212, 2010.