

Paper BIO-03: Adaptive Flux Limitation in Erythropoiesis and Anaemia: A CDFD Model of Blood Production Under Constraint

Steve Bico Mujjabi
Independent Researcher

May 22, 2026

Abstract

This paper presents **Adaptive Flux Limitation (AFL)** as the Part III biology-and-medicine model inside Constraint-Driven Flux Dynamics (CDFD). CDFD supplies the flow-constraint-memory grammar: physiological flow Φ , constraint C , surface responsiveness S , and structural memory M_s . The Runtime citation anchors the CDFL implementation, while clinical interpretation remains separate from software output and bedside validation.

Anemia of chronic disease, including in chronic kidney disease (CKD), is often explained by erythropoietin deficiency, iron restriction, and inflammation (system-wide constraint C amplification). Clinical responses are nonlinear: erythropoiesis-stimulating agents (ESAs) show diminishing returns, and single-modality therapy frequently underperforms.

We model erythropoiesis as an AFL system: **throughput** Φ is erythropoietic output (red-cell production intensity), **constraint** C aggregates iron availability, inflammatory load, and marrow capacity, and $\Psi_s = (\Phi/C) \cdot S \cdot M_s$ is a dimensionless adequacy ratio in normalized units. An active-numbered public supplement should contrast healthy, CKD, ESA-only, and multi-modality scenarios, illustrating ESA resistance when C remains elevated.

Status: Inferred

Within the extended framework of this series, biological networks are modeled as *Adaptive Surfaces* that dynamically integrate physiological load and retain structural memory. The system's health is governed by the adaptive ratio $\Psi_s = (\Phi/C) \cdot S \cdot M_s$, where homeostasis ($\Psi_s \approx 1.0$) is maintained near the stable attractor ($S \cdot M_s \approx 1.0$), and disease emerges as surface dysregulation when maladaptive memory locks tissues into chronic constraint.

Greek-symbol convention. Unless a manuscript states a tighter equation-local meaning, Greek symbols are local CDFD variables or coefficients, not universal constants. Here Φ denotes driving flux, Ψ_s denotes the adaptive operating ratio, Ω denotes a domain or state space, and Λ denotes the Life Number where used. The coefficients α , β , and γ denote accumulation or reinforcement, relaxation or decay, and diffusion, coupling, or redistribution. The symbols δ and ϵ denote perturbation or tolerance scales, while η , λ , μ , ν , ρ , σ , τ , and ϕ denote local efficiency, rate, baseline, frequency, density or correlation, noise or variance, time-scale, and phase or field terms. Subscripts restrict any coefficient to the named pathway, tissue, domain, or experiment.

Clinical Reader's Guide

This paper is the first direct bridge from notation to clinical reasoning. It frames disease trajectories as movements through a state space rather than as isolated abnormal laboratory values. In practice,

a clinician sees this every day: patients worsen because a compensatory loop that once protected them becomes exhausted, rigid, or misdirected.

In AFL terms, the bedside question becomes: what is the patient trying to move, what is blocking it, what response surface is still flexible, and what previous stress has become retained memory? This is why a single input escalation can fail. More erythropoietin, more insulin, more vasopressor, or more immunosuppression can be rational in one state and harmful or futile in another.

The paper’s translational claim is deliberately falsifiable. If Φ , C , S , and M_s proxies do not improve risk stratification, timing, or treatment matching beyond standard clinical variables, then AFL remains only a metaphor. The value of the framework depends on measurable trajectories.

1 Introduction

Adaptive Flux Limitation (AFL) is the named model used in this paper; CDFD is the parent framework that supplies the variables, closure logic, and claim discipline. The biological or medical question is posed first as AFL, then interpreted as a CDFD model of flow, constraint, surface responsiveness, and structural memory.

1.1 Formal Symbol Rigor: The Extended CDFD Paradigm

In strict alignment with the Fundamental Physics (Part I) and Origins of Life (Part II) archives, this biological translation formally preserves the exact Mujjabi transport tensor structure:

- Φ (**Flow Intensity**): The physiological or biochemical drive (e.g., nutrient flux, signaling rate).
- C (**Constraint / Pathway Resistance**): The biological resistance regulating the flow. It dynamically evolves via the standard closure: $dC/dt = \alpha\Phi - \beta C$.
- S (**Surface Responsiveness**): The active response capability of the biological medium.
- M_s (**Surface Memory**): Structural hysteresis or maladaptive drift (e.g., chronic scarring, epigenetic locking) with a finite recovery time τ_M .
- $\Psi_s = (\Phi/C) \cdot S \cdot M_s$ (**Operating State Ratio**): The dimensionless index of biological health. $\Psi_s \approx 1$ defines homeostasis ($\Psi_s \approx 1.0$); $\Psi_s \gg 1$ defines pathological overload.

Erythropoiesis integrates EPO signaling, iron trafficking, marrow reserve, oxygen demand, and energy supply. Reductionist single-factor explanations miss coupled failures: ESA escalation with partial hemoglobin response, functional iron deficiency despite adequate stores, and inflammatory blunting.

AFL-2 argued that diverse biological domains admit a shared $(\Phi, C, S, M_s, \Psi_s)$ skeleton. **Open gap addressed here:** whether a *clinical* pathway exhibits the same qualitative coupling—especially ESA resistance driven by non-EPO constraints.

Status: Hypothesis

2 Model and Assumptions

2.1 Variables

- Φ : erythropoietic throughput (production intensity), driven in part by EPO analogue S_{EPO} in the supplement.
- C : effective constraint combining iron burden χ_{Fe} , inflammatory load χ_{inf} , and marrow cap χ_{mar} (positive numbers; larger means tighter limitation in the toy normalization).
- $\Psi_s = (\Phi/C) \cdot S \cdot M_s$: adequacy ratio; bands in the script mirror AFL-1–2 conventions for labeling only.

The scalar C is therefore an aggregate marrow constraint, not a single assay or single molecular pathway.

Status: Derived (definitions)

2.2 0D coupled dynamics used in the supplement

The script integrates

$$\dot{\Phi} = S_{\text{EPO}} - \Phi C, \quad (1)$$

$$\dot{C} = \alpha |\dot{\Phi}| - \beta C, \quad (2)$$

with nonnegative α, β and scenario-dependent initial effective C from $(\chi_{\text{Fe}}, \chi_{\text{inf}}, \chi_{\text{mar}})$. This is a minimal caricature, not a marrow agent-based model.

Status: Hypothesis

2.3 Assumptions C1–C3

- **C1** A single Φ represents net erythropoietic output adequate for comparing scenarios at cohort scale.
- **C2** Iron and inflammation (system-wide constraint C amplification) enter primarily through effective C , not separate state variables, in this toy.
- **C3** ESA acts by increasing S_{EPO} without automatically lowering C .

Status: Open

3 Deep Analysis: Biological Implications of the CDFT/AFL Framework

3.1 Homeostasis as C management

Within CDFT/AFL, homeostasis ($\Psi_s \approx 1.0$) is continuous management of C to keep $\Phi < 1$ across all constraint domains. AMPK [Hardie et al., 2012], mTOR, HIF-1 α [Semenza, 2012], and the baroreceptor reflex are distributed sensors of Φ_i that trigger the fast dynamics (Eq. 10) to reduce C or J_{req} when Φ_i rises. Disease begins when these mechanisms exhaust their capacity to keep $\Phi < 1$.

3.2 Resistance as adaptive C upregulation

AFL reframes biological resistance as the system’s correct response: adaptive C upregulation when flux approaches the constraint ceiling. ESA hyporesponsiveness [Macdougall, 1995, Vaziri, 2004], insulin resistance (chronic C_E upregulation) [DeFronzo, 2009, Petersen and Shulman, 2018], T-cell exhaustion [O’Neill et al., 2016, Buck et al., 2015], and antimicrobial resistance [Davies and Davies, 2010, Andersson and Hughes, 2010] all represent states where demand has approached or exceeded $J_{\max}$, and the system has raised C (through fast dynamics) or allowed C_i to erode (slow dynamics) to maintain structural integrity.

In ESA hyporesponsiveness: $J_{\text{erythropoiesis}} \leq \min\{J_{\text{methyl}}, J_{\text{energy}}, J_{\text{redox}}\}$. Escalating EPO dose increases J_{req} without expanding C_i or reducing C , deepening constraint stress without restoring output.

3.3 Cancer as C bypass

Malignant transformation represents partial escape from AFL through coordinated modification of the constraint architecture: the Warburg effect [Warburg, 1956, Vander Heiden et al., 2009] raises glycolytic C_{energy} ; NRF2 activation expands redox C_{redox} [Schieber and Chandel, 2014]; epigenetic deregulation reduces transcriptional C . In CDFT terms, cancer cells engineer themselves to have lower C for proliferative processes — operating near $\Phi \approx 1$ with minimal reserve, generating vulnerability to multi-constraint perturbation.

3.4 Disease dynamics across contexts

Figure 1 illustrates CDFT/AFL dynamics across four disease contexts.




Figure 1: **CDFT/AFL Dynamics across Disease Contexts.** All four panels can be read in CDFT language as $D_{\text{eff}} = 1/C$ changing over time. **(A)** Aging: C rises as C_i erodes ($dC_i/dt = -\lambda_i J + R_i$); D_{eff} falls; transport capacity declines. **(B)** insulin resistance (chronic C_E upregulation): $C_{\text{metabolic}}$ rises progressively; $D_{\text{eff,glucose}}$ falls; glucose uptake decouples from insulin signal. **(C)** Differential stress response: cancer cells (*engineered low C , high Φ*) collapse under dual-constraint perturbation; normal cells (high C reserve) tolerate it. **(D)** Pharmacological saturation: drug effect is limited by the system’s processing C ; escalating dose cannot overcome a raised C . **KEY:** All four panels are consequences of the same CDFT governing equation $F = \nabla X/C(\Omega, t)$ under domain-specific parameterization.

3.5 Aging as C drift

Aging is progressive, asymmetric drift of C upward — the slow dynamics operating without adequate repair:

$$\frac{dC_i}{dt} = -\lambda_i J + R_i < 0 \implies \frac{dC_i}{dt} > 0 \quad (3)$$

$D_{\text{eff}} = 1/C$ falls across all constraint domains. The rate of functional aging is determined not by chronological time but by the cumulative integral of $\Phi(t) \cdot \lambda_i$ — a prediction with direct implications for longevity interventions [Longo and Mattson, 2014, Mattson et al., 2017].

3.6 Embryological development as C -governed flux coordination

Embryological development requires sustained high flux across multiple constraint domains simultaneously [Levin, 2021, 2012, Tseng and Levin, 2013]:

$$J_{\text{dev}} = J_{\text{prolif}} + J_{\text{diff}} + J_{\text{migr}} + J_{\text{pattern}} \leq \frac{\nabla X_{\text{dev}}}{C_{\text{dev}}(\Omega, x, t)} \quad (4)$$

In CDFT terms, the embryo is an adaptive medium in which bioelectric fields (B_e component of Ω) establish spatial patterns of C that guide tissue organization. Bioelectric patterning is not separate from mechanical development — it is the mechanism by which the embryo sets its own $C(x)$ landscape, establishing which regions are “open” (low C , high flux permitted) and which are “closed” (high C , flux restricted). Neural Tube Defects reflect $\Phi_M \rightarrow 1$ in the methylation domain during the closure window, when C_M is too high for the flux required for coordinated proliferation.

3.7 Stem cell regeneration and neural plasticity as C -reserve systems

Stem cells maintain low C_{regen} as a latent capacity reserve. Age-related regenerative failure occurs as the slow dynamics erode this reserve:

$$J_{\text{regen}} \leq \frac{\nabla X_{\text{regen}}}{C_{\text{regen}}(\Omega, t)} \quad (5)$$

Neural plasticity is governed by $P(\Omega) = g(E_n, B_e, E_p) = 1/C_{\text{plasticity}}$: learning is biologically possible only when the medium is sufficiently “open” (low $C_{\text{plasticity}}$) for synaptic reorganization.

4 Bidirectional Constraint Dynamics in Clinical Settings

4.1 The two-timescale model

Classical AFL (previous versions of this framework) included only the slow structural erosion equation. This captured chronic disease and aging but missed the acute protective upregulation that is the first line of biological defence against constraint saturation. We now formally introduce both:

Fast timescale (seconds–hours): Protective upregulation

$$\frac{dC}{dt} = \alpha F - \beta C \quad (6)$$

When flux F rises acutely, the system actively increases C as a protective response — enzyme product inhibition, baroreceptor reflex, immune negative feedback, autophagy induction, stress-activated protein kinase pathways. βC is the relaxation term: when flux normalizes, C decays back toward baseline with time constant $1/\beta$.

At quasi-steady state of Equation (6):

$$C^* = \frac{\alpha}{\beta} F \quad (7)$$

Substituting into the CDFT law $F = \nabla X/C$: $F = \nabla X \beta / (\alpha F)$, giving $F^2 = (\beta/\alpha) \nabla X$, or $F = \sqrt{(\beta/\alpha) \nabla X}$. This is the **square-root flux law** — a nonlinear relationship between driving gradient and flux that emerges from the adaptive medium, absent in any classical transport theory.

Slow timescale (years–decades): Structural erosion

$$\frac{dC_i}{dt} = -\lambda_i \cdot J + R_i \quad (8)$$

When chronic flux damage $\lambda_i J$ exceeds repair capacity R_i , the constraint ceiling C_i declines progressively. This is the aging and chronic disease trajectory, predicting decompensation at time $t^* = (C_i^0 - C_i^{\text{threshold}})/(\lambda_i J - R_i)$.

Missing figure asset: figures/fig9_bidirectional.png

Figure 2: Bidirectional Constraint Dynamics: Fast Protective Upregulation and Slow Structural Erosion. **(A)** Fast timescale ($dC/dt = \alpha F - \beta C$): constraints rise protectively when flux spikes; they relax when flux normalizes (time constant $1/\beta$). In CDFT terms: the medium temporarily increases its own resistance to protect structural integrity. **(B)** Slow timescale ($dC_i/dt = -\lambda_i J + R_i$): when cumulative damage exceeds repair, the constraint ceiling erodes progressively. Predicts age of decompensation from the patient's chronic flux history. **(C)** Combined two-timescale system: fast acute homeostatic oscillations on slow structural drift. $D_{\text{eff}} = 1/C(t)$ integrates both timescales. **KEY:** The two equations describe different mechanisms at different timescales. They operate simultaneously, and together they account for both acute resilience and chronic vulnerability.

4.2 The combined two-equation system

Together, Equations (6) and (8) constitute the complete CDFT/AFL temporal dynamics:

$$\text{Flux law: } F = \frac{\nabla X}{C} \quad (9)$$

$$\text{Fast constraint adaptation: } \frac{dC}{dt} = \alpha F - \beta C \quad (10)$$

$$\text{Slow capacity erosion: } \frac{dC_i}{dt} = -\lambda_i F + R_i \quad (11)$$

This three-equation system (flux law + fast dynamics + slow dynamics) describes growth, stabilization, acute adaptation, chronic erosion, and ultimate collapse:

- **Homeostasis:** C quickly adjusts to sustain $\Phi < 1$; C_i stable when $\lambda_i F < R_i$
- **Acute stress:** C rises protectively (Eq. 10); flux temporarily suppressed; resolves when stress removed
- **Chronic disease:** C_i erodes slowly (Eq. 11); C^* must rise to compensate; Φ creeps toward 1
- **Decompensation:** C_i falls below critical threshold; C can no longer compensate; $\Phi \rightarrow 1$ and bifurcation occurs

4.3 Biological examples of fast protective upregulation

The fast dynamics (Eq. 10) describe mechanisms that are well-documented but not previously unified under a single framework:

- **Enzyme product inhibition:** High flux products inhibit upstream enzymes — classic negative feedback. In CDFT: the enzyme acts as a C transducer that increases resistance when output is high.
- **Baroreceptor reflex:** High arterial pressure triggers vasodilation and heart rate reduction — reducing F by increasing vascular C .

- **Immune negative feedback:** IL-10, TGF- β , regulatory T cells suppress immune activation when effector flux is high — increasing immunological C .
- **Metabolic AMPK activation:** High AMP/ATP ratio activates AMPK, which suppresses anabolic processes — reducing metabolic demand J_{req} to match declining J_{max} .
- **Autophagy induction under nutrient stress:** When substrate flux falls, autophagy recycles intracellular components to maintain energetic flux — an adaptive reduction of C_E through internal resource mobilization.

5 Main Results

5.1 Multi-constraint anemia

When χ_{inf} and χ_{Fe} elevate C , the same S_{EPO} yields lower Ψ_s : anemia as **regulated** low throughput under high constraint, not only “low EPO.”

Status: Inferred

5.2 ESA resistance mechanism in the model

Boosting S_{EPO} raises $\dot{\Phi}$ transiently, but if C adapts upward or remains large from iron/inflammation (system-wide constraint C amplification), Φ plateaus: diminishing returns consistent with clinical ESA resistance *in this toy parameterization*.

Status: Inferred

5.3 Multi-modality advantage

Lowering χ_{Fe} and χ_{inf} in tandem with raised S_{EPO} increases final Ψ_s more than ESA alone in the script’s Table 2 comparison.

Status: Derived (from supplied numerics)

6 Numerical Verification

Use an active-numbered public supplement. Table 1 lists scenarios (Healthy; CKD stages; ESA; ESA+iron; ESA+iron+anti-inflammatory) with final Ψ_s and a discrete regime tag. Table 2 quantifies the percentage improvement in final Ψ_s for combination vs. ESA-only under the stated parameters.

Status: Derived

7 Interpretation

The framework reframes anemia of chronic disease as **coupled** throughput–constraint pathology: therapies that only move Φ while leaving C high are predicted to underperform. This is consistent with guideline emphasis on iron repletion and inflammation (system-wide constraint C amplification) control, but the model does not replace trials or individualized physiology.

Status: Inferred

8 Limitations and Open Problems

- No patient-level calibration; units are normalized.
- Hepcidin dynamics, hypoxia sensing, and stem-cell niches are collapsed into C .
- Safety (ESA cardiovascular risk) is not modeled.

Status: Open

9 Biological Mujjabi Laws & Discoveries

9.1 The Mujjabi Boundary-Covariance Test [Mujjabi \[2026\]](#)

Derived from the Origin of Life boundary conditions, the **Mujjabi Boundary-Covariance Test** [Mujjabi \[2026\]](#) states that tissue health requires internal metabolic flux (Φ) to strictly covary with boundary integrity (S). If flux scales up while boundary responsiveness drops (e.g., leaky gut, endothelial dysfunction), the adaptive ratio Ψ_s decouples. Diagnosis of disease must verify this covariance, treating the boundary not as a passive wall, but as an active regulatory tensor.

10 Translational Medicine: The AFL Flux-Shunting Theorem in Sepsis and ICU Cascades

Ward-level medicine frequently encounters multi-organ failure. The CDFD framework mathematically models this via the **AFL Flux-Shunting Theorem**.

When a primary organ’s constraint hits a critical threshold (e.g., $C_{renal} \rightarrow \infty$), the unresolvable physiological drive (Φ) is not destroyed; it is topologically shunted to parallel vascular and tissue networks. This forced redirection of Φ into the cardiovascular system triggers systemic hypertension, vascular shearing, and subsequent heart failure. Intensive care unit (ICU) decompensation is thus modeled not as a sequence of independent organ failures, but as a topological cascade of overflowing Ψ_s forcing adjacent systems past their Mujjabi Capacity limits.

11 Conclusion

AFL-3 applies the AFL contract to erythropoiesis: ESA resistance emerges when constraints remain high. The supplement provides deterministic contrasts aligned with Tables T1–T2.

Status: Inferred

12 Reproducibility Note

- **Script packaging:** active-numbered public supplement.
- **Dependencies:** NumPy; runtime $\ll 1$ s.
- **Outputs:** console Tables 1–2; improvement percentage in Table 2 footer.

Status: Derived

13 Clinical Translation Layer

The clinical translation of AFL begins with patient-level mapping. A disease state is not represented as one number, but as coupled axes: flux demand, constraint reserve, relaxation capacity, and compensatory cost. In anaemia of chronic disease, for example, erythropoietic flux cannot be interpreted without iron handling, inflammation (system-wide constraint C amplification), renal signaling, marrow reserve, and oxygen demand. In metabolic disease, glucose flux cannot be interpreted without mitochondrial capacity, adipose buffering, inflammatory tone, and hepatic handling.

14 Patient-Level Workflow

A practical workflow has four steps. First, identify the dominant flux: oxygen delivery, glucose throughput, inflammatory signaling, phosphate handling, drug exposure, or cell-proliferation drive. Second, identify the active constraint: renal reserve, mitochondrial redox capacity, hepcidin-mediated iron restriction, receptor saturation, vascular stiffness, immune regulation, or tissue perfusion. Third, estimate whether the patient is in compensation, saturation, or collapse. Fourth, choose an intervention that changes the relevant axis rather than simply pushing more flux through a constrained system.

15 Clinical Prediction

The central clinical prediction is that failed therapy often reflects a mismatched axis. Increasing flux when constraint is the dominant problem should produce diminishing returns or toxicity. Reducing constraint when flux is absent should produce little benefit. Multi-modal therapy works when each intervention targets a different limiting axis: source reduction, constraint relief, relaxation enhancement, or redistribution of load.

16 Medical Falsifiability

AFL clinical translation fails if it cannot improve stratification beyond standard disease categories, if proposed constraint variables do not track outcome, or if predicted multi-axis interventions do not outperform single-axis escalation in appropriately matched patients. The framework should be tested against clinical trajectories, not anecdotes.

17 Clinical Mapping of Symbols

Symbol	Clinical meaning	ICU / clinical proxy	Invalidation condition
Φ	Physiological flux (e.g., cardiac output, oxygen delivery)	Lactate, vasopressor dose, urine output	If Φ proxy cannot be separated from haemodynamic drivers, mapping collapses
C	Multi-organ constraint load	CRP, creatinine, endothelial leak markers	If C is indistinguishable from severity scores (SOFA, APACHE), AFL adds no stratification
S	Organ responsiveness reserve	Pre-illness organ function; frailty index	If S is uncorrelated with recovery trajectory, the surface framing adds nothing
M_s	Post-ICU memory / PICS	Persistent inflammatory markers; cognitive testing at 6 months	If M_s decays to baseline within 24h, memory framing is not needed

Status: Hypothesis

18 Clinical Safety Boundary

This paper is a research framework for ICU cascade framing, not a clinical protocol.

- Sepsis management: SSC 2021 guidelines take precedence — antimicrobials, source control, fluid assessment, vasopressors. AFL framing does not replace any SSC 2021 recommendation.
- Multi-organ failure: organ support decisions (dialysis, ventilation, vasopressor titration) are made by specialist teams using established monitoring, not Ψ_s values.
- Post-ICU sequelae: PICS rehabilitation is an active clinical field; AFL memory framing is a hypothesis, not a rehabilitation protocol.

19 Runtime Diagnostic

Release-local script: `supplementary/supplementary_afl_03.py`. Outputs in `outputs/paper_03/`: `sepsis_cascade.csv`, `cardiorenal_transfer.csv`, `post_icu_recovery.csv`, `icu_organ_coupling.png`, `summary.json`. All outputs are toy diagnostics under stated assumptions.

Conflict of Interest

The author declares no conflict of interest.

Funding

This research received no external funding.

Author Contributions

Conceptualization, formal analysis, runtime engineering, and manuscript preparation: S.B.M.

Data Availability

All computational models and scripts are in the `supplementary/` directory.

20 Reference Layer

References

- Dan I. Andersson and Diarmaid Hughes. Antibiotic resistance and its cost: Is it possible to reverse resistance? *Nature Reviews Microbiology*, 8:260–271, 2010.
- Michael D. Buck, David O’Sullivan, and Erika L. Pearce. T cell metabolism drives immunity. *Journal of Experimental Medicine*, 212(9):1345–1360, 2015. doi: 10.1084/jem.20151159.
- Julian Davies and Dorothy Davies. Origins and evolution of antibiotic resistance. *Microbiology and Molecular Biology Reviews*, 74(3):417–433, 2010. doi: 10.1128/MMBR.00016-10.
- Ralph A. DeFronzo. From the triumvirate to the ominous octet: A new paradigm for the treatment of type 2 diabetes mellitus. *Diabetes*, 58(4):773–795, 2009.
- D. Grahame Hardie, Fiona A. Ross, and Simon A. Hawley. Ampk: A nutrient and energy sensor that maintains energy homeostasis. *Nature Reviews Molecular Cell Biology*, 13(4):251–262, 2012. doi: 10.1038/nrm3311.
- Michael Levin. Molecular bioelectricity in developmental biology: New tools and recent discoveries. *BioEssays*, 34(3):205–217, 2012. doi: 10.1002/bies.201100136.
- Michael Levin. Bioelectric signaling: Reprogrammable circuits underlying embryogenesis, regeneration, and cancer. *Cell*, 184:1971–1989, 2021.
- Valter D. Longo and Mark P. Mattson. Fasting: Molecular mechanisms and clinical applications. *Cell Metabolism*, 19(2):181–192, 2014. doi: 10.1016/j.cmet.2013.12.008.
- Iain C. Macdougall. Poor response to erythropoietin: Practical guidelines on investigation and management. *Nephrology Dialysis Transplantation*, 10:607–614, 1995.
- Mark P. Mattson, Valter D. Longo, and Michelle Harvie. Impact of intermittent fasting on health and disease processes. *Ageing Research Reviews*, 39:46–58, 2017. doi: 10.1016/j.arr.2016.10.005.
- Steve Bico Mujjabi. Cdfd part ii: Origins of life and tri-regime bioenergetics. 2026. doi: 10.5281/zenodo.20264779. URL <https://doi.org/10.5281/zenodo.20264779>.
- Luke A. J. O’Neill, Ryan J. Kishton, and Jeffrey Rathmell. A guide to immunometabolism for immunologists. *Nature Reviews Immunology*, 16:553–565, 2016.
- Kitt Falk Petersen and Gerald I. Shulman. Mechanisms of insulin action and insulin resistance. *Physiological Reviews*, 98(4):2133–2223, 2018.

- Michael Schieber and Navdeep S. Chandel. Ros function in redox signaling and oxidative stress. *Current Biology*, 24(10):R453–R462, 2014. doi: 10.1016/j.cub.2014.03.034.
- Gregg L. Semenza. Hypoxia-inducible factors in physiology and medicine. *Cell*, 148(3):399–408, 2012. doi: 10.1016/j.cell.2012.01.021.
- Ai-Sun Tseng and Michael Levin. Cracking the bioelectric code: Probing endogenous ionic controls of pattern formation. *Communicative and Integrative Biology*, 6(1):e22595, 2013. doi: 10.4161/cib.22595.
- Matthew G. Vander Heiden, Lewis C. Cantley, and Craig B. Thompson. Understanding the warburg effect: The metabolic requirements of cell proliferation. *Science*, 324:1029–1033, 2009.
- Nosratola D. Vaziri. Oxidative stress in chronic renal failure: The nature, mechanisms, and consequences. *Seminars in Nephrology*, 24(5):469–473, 2004. doi: 10.1016/j.semnephrol.2004.06.005.
- Otto Warburg. On the origin of cancer cells. *Science*, 123:309–314, 1956.