

Paper BIO-09: Adaptive Flux Limitation in Pharmacology: A CDFD Model of Dose, Resistance, and Therapeutic Optimization

Steve Bico Mujjabi
Independent Researcher

May 22, 2026

Abstract

This paper presents **Adaptive Flux Limitation (AFL)** as the Part III biology-and-medicine model inside Constraint-Driven Flux Dynamics (CDFD). CDFD supplies the flow-constraint-memory grammar: physiological flow Φ , constraint C , surface responsiveness S , and structural memory M_s . The Runtime citation anchors the CDFL implementation, while clinical interpretation remains separate from software output and bedside validation.

Drug response in biological systems is traditionally described using pharmacokinetics (PK) and pharmacodynamics (PD). However, clinical observations frequently reveal nonlinear effects, diminishing returns, tolerance, and resistance that are not fully explained by conventional models.

In this work, we apply Adaptive Flux Limitation (AFL) to pharmacology, framing drug action as a flux-driven perturbation within constraint-regulated biological systems. We demonstrate that drug absorption, distribution, metabolism, and effect are governed by dynamic constraints that adapt in response to sustained exposure.

We show that drug resistance, tolerance, and therapeutic plateaus emerge naturally from adaptive constraint dynamics. This framework provides a unified interpretation of pharmacological behavior and suggests new strategies for dosing, combination therapy, and treatment optimization.

Status: Inferred

Within the extended framework of this series, biological networks are modeled as *Adaptive Surfaces* that dynamically integrate physiological load and retain structural memory. The system's health is governed by the adaptive ratio $\Psi_s = (\Phi/C) \cdot S \cdot M_s$, where homeostasis ($\Psi_s \approx 1.0$) is maintained near the stable attractor ($S \cdot M_s \approx 1.0$), and disease emerges as surface dysregulation when maladaptive memory locks tissues into chronic constraint.

Greek-symbol convention. Unless a manuscript states a tighter equation-local meaning, Greek symbols are local CDFD variables or coefficients, not universal constants. Here Φ denotes driving flux, Ψ_s denotes the adaptive operating ratio, Ω denotes a domain or state space, and Λ denotes the Life Number where used. The coefficients α , β , and γ denote accumulation or reinforcement, relaxation or decay, and diffusion, coupling, or redistribution. The symbols δ and ϵ denote perturbation or tolerance scales, while η , λ , μ , ν , ρ , σ , τ , and ϕ denote local efficiency, rate, baseline, frequency, density or correlation, noise or variance, time-scale, and phase or field terms. Subscripts restrict any coefficient to the named pathway, tissue, domain, or experiment.

Clinical Reader’s Guide

This pharmacology paper is about why dose-response curves stop being simple in living systems. A drug does not act on a passive target. It enters a system that adapts, compensates, saturates, remembers prior exposure, and redistributes burden to other pathways. AFL describes those changes as movement in Φ , C , S , M_s , and relaxation rate β .

The core clinical distinction is same-axis versus cross-axis intervention. Two drugs can both lower the same driver, which may create redundancy, plateau response, or toxicity. Another pair may act on different axes, such as reducing source load while improving relaxation or lowering a compensatory constraint. This is a model taxonomy for thinking about combinations, not an automatic prescribing rule.

Any real pharmacological use would require pharmacokinetics, pharmacodynamics, adverse-event data, trial evidence, interaction checks, organ function, patient context, and guideline alignment. The runtime output is a small consistency check of the taxonomy only.

AFL-9 PK/PD notation. Drug concentration is D to avoid conflict with constraint C . Biological constraint is C , physiological or pharmacologic drive is Φ , and $\Psi_s = (\Phi/C) \cdot S \cdot M_s$ is used only when the dominant flow can be specified.

1 Introduction

Adaptive Flux Limitation (AFL) is the named model used in this paper; CDFD is the parent framework that supplies the variables, closure logic, and claim discipline. The biological or medical question is posed first as AFL, then interpreted as a CDFD model of flow, constraint, surface responsiveness, and structural memory.

1.1 Formal Symbol Rigor: The Extended CDFD Paradigm

In strict alignment with the Fundamental Physics (Part I) and Origins of Life (Part II) archives, this biological translation formally preserves the exact Mujjabi transport tensor structure:

- Φ (**Flow Intensity**): The physiological or biochemical drive (e.g., nutrient flux, signaling rate).
- C (**Constraint / Pathway Resistance**): The biological resistance regulating the flow. It dynamically evolves via the standard closure: $dC/dt = \alpha\Phi - \beta C$.
- S (**Surface Responsiveness**): The active response capability of the biological medium.
- M_s (**Surface Memory**): Structural hysteresis or maladaptive drift (e.g., chronic scarring, epigenetic locking) with a finite recovery time τ_M .
- $\Psi_s = (\Phi/C) \cdot S \cdot M_s$ (**Operating State Ratio**): The dimensionless index of biological health. $\Psi_s \approx 1$ defines homeostasis ($\Psi_s \approx 1.0$); $\Psi_s \gg 1$ defines pathological overload.

Pharmacology traditionally separates drug behavior into:

- pharmacokinetics (PK): what the body does to the drug
- pharmacodynamics (PD): what the drug does to the body

However, real-world treatment reveals:

- diminishing response with repeated dosing
- development of resistance
- variability between patients

These suggest that drug response is governed by dynamic system-level regulation.

2 Drug Action as Flux Perturbation

Define drug flux:

$$J_{drug} = \text{rate of drug interaction with biological targets} \quad (1)$$

Drug effect depends on:

$$J_{effect} = f(J_{drug}, C_{system}) \quad (2)$$

Where:

- C_{system} = biological constraints

3 Pharmacokinetics as Flux Transport

Drug concentration D evolves as:

$$\frac{\partial D}{\partial t} + \nabla \cdot \mathbf{J} = S - E \quad (3)$$

Where:

- D = drug concentration
- S = absorption
- E = elimination

Flux depends on PK constraints:

$$\mathbf{J} = -\frac{1}{C_{PK}} \nabla D \quad (4)$$

Constraints include:

- membrane permeability
- protein binding
- organ function (liver, kidney)

4 Pharmacodynamics as Constrained Response

Drug effect:

$$E_{drug} = \frac{J_{drug}}{C_{PD}} \quad (5)$$

Constraints include:

- receptor availability
- signaling pathway capacity
- downstream metabolic limits

5 Adaptive Constraint Dynamics

Under repeated exposure:

$$\frac{dC}{dt} = \alpha J_{drug} - \beta C \quad (6)$$

Meaning:

- high drug flux increases resistance
- system adapts to reduce effect

6 Drug Tolerance

Tolerance emerges as:

$$C_{PD} \uparrow \Rightarrow E_{drug} \downarrow \quad (7)$$

Examples:

- opioid tolerance
- beta-receptor downregulation

7 Drug Resistance

Resistance is a stronger form of constraint adaptation.

7.1 Mechanisms

- receptor mutation
- efflux pumps
- metabolic bypass

AFL interpretation:

$$C_{system} \gg 1 \Rightarrow J_{effect} \approx 0 \quad (8)$$

8 Antibiotics and Chemotherapy

High drug flux creates strong selective pressure:

$$J_{drug} \uparrow \Rightarrow C_{resistance} \uparrow \quad (9)$$

Result:

- resistant populations dominate

9 Diminishing Returns

Increasing dose:

$$J_{drug} \uparrow \quad (10)$$

But:

$$C_{PD} \uparrow \quad (11)$$

Thus:

$$E_{drug} \text{ plateaus} \quad (12)$$

10 Optimal Dosing Strategy

AFL suggests:

- avoid sustained high flux
- allow constraint relaxation

10.1 Pulsatile Dosing

$$J_{drug}(t) = \text{intermittent} \quad (13)$$

Allows:

$$C \downarrow \text{ between doses} \quad (14)$$

11 Combination Therapy

Multiple drugs target different constraints:

$$C_{total} = C_1 + C_2 + \dots \quad (15)$$

Reduces probability of resistance.

12 Personalized Medicine

Patient variability arises from:

- different baseline constraints
- different adaptive rates

Thus:

$$C = C_{\text{patient-specific}} \quad (16)$$

13 Flux–Constraint Regime

$$\Psi_{s,drug} = \frac{J_{drug}}{C_{system}} \quad (17)$$

- $\Psi_s < 1$: insufficient effect
- $\Psi_s \approx 1$: optimal therapeutic range
- $\Psi_s > 1$: toxicity or resistance induction

14 Clinical Implications

14.1 Why Treatments Fail

- constraints adapt faster than therapy

14.2 Therapeutic Design

- dynamic dosing schedules
- multi-target strategies

15 Predictions

AFL predicts:

- tolerance with repeated dosing
- resistance under high sustained drug pressure
- improved outcomes with adaptive dosing strategies

16 Numerical Verification

This installment’s toy deterministic checks should be packaged as an active-numbered public supplement (NumPy; seed and parameters in-file). Console tables illustrate the adopted AFL closure; they do not substitute for cohort or experimental validation.

Status: Derived

17 Interpretation

Domain-specific narratives above map mechanisms to $(\Phi, C, S, M_s, \Psi_s)$ within the AFL working model. Interpretive claims are model-mediated; claim strengths follow the public status labels used throughout this paper.

Status: Inferred

18 Limitations and Open Problems

- simplified representation of molecular pharmacology
- requires experimental validation

Status: Open

19 Falsifiability

AFL would be invalid if:

- drug response remains constant under repeated exposure
- resistance does not correlate with drug pressure

Status: Open

20 Deep Analysis: Pharmacological Dose–Response and Constraint Dynamics

20.1 Pharmacology: receptor C and processing C

Drug effect is bounded by the adaptive medium through which it must act:

$$J_{\text{drug}} \leq \frac{\nabla X_{\text{drug}}}{C_S + C_E + C_M} \quad (18)$$

C_S governs receptor binding capacity and transduction pathway availability; C_E and C_M govern downstream biosynthetic and metabolic processing. As dose increases ∇X_{drug} without reducing C , the adaptive medium reaches its D_{eff} floor and output saturates while toxicity continues to escalate. Drug synergy emerges most powerfully when combinations simultaneously reduce C_S , C_E , and C_M in high- Φ target tissues while leaving C reserve in normal tissues.

20.2 Ecology: carrying capacity as macroscopic C

At ecological scales, population growth follows logistic dynamics [May, 1976]:

$$\frac{dN}{dt} = rN \left(1 - \frac{N}{K} \right) \quad (19)$$

In CDFT language: K is the macroscopic C ceiling for population flux — $J_{\text{pop}} \leq \nabla X_{\text{pop}}/C_{\text{env}}$ where $C_{\text{env}} = f(J_{\text{energy}}, J_{\text{nutrient}}, J_{\text{habitat}})$. As $N \rightarrow K$, C_{env} rises (resource depletion raises the environmental adaptive medium’s resistance to population growth), and $dN/dt \rightarrow 0$ regardless of reproductive drive. Trophic energy transfer: $J_{ij} = \eta_{ij} J_i$ with $\eta_{ij} \approx 0.10$ represents D_{eff} losses across trophic levels — each level is a more constrained adaptive medium than the one below it.

21 Clinical Mapping of Symbols

Table 1: AFL symbol map for pharmacology and dose-constraint control.

Symbol	Pharmacological meaning	Measurable proxy	Invalidation condition
Φ	Drug flux reaching target tissue	Peak plasma concentration ($C_{\max}$)	If $C_{\max}$ does not pre
C	Receptor/pathway constraint	Receptor occupancy, off-target binding	If constraint is dose-
Ψ_s	Pharmacodynamic operating ratio	Dose-response curve steepness (EC50)	If EC50 is constrain
M_s	Pharmacological memory / tolerance	Time to tolerance onset (days)	If tolerance is absent

22 Known Medicine Baseline

The following guidelines and frameworks take precedence over any AFL model output and are not superseded by this paper:

- **FDA Project Optimus (2023–2026):** Oncology dose-optimisation framework requiring dose-response characterisation and evaluation of combinations below MTD; AFL’s constraint-orthogonality framing is a hypothesis for future alignment with Optimus dose-finding study designs.
- **ICH E4 Dose-Response Guidance:** Mandates dose-response study designs; AFL’s saturation-regime and adaptive-resistance regimes should be framed as design hypotheses, not regulatory claims.
- **CPIC Pharmacogenomics Guidelines:** Genotype-guided dosing in cytochrome P450 polymorphisms; AFL constraint variables (C_S , C_E , C_M) are candidate correlates of pharmacogenomic variance — hypothesis only.

AFL toy simulations do not substitute for pharmacokinetic/pharmacodynamic (PK/PD) modelling, adverse-event monitoring, or regulatory dose-finding studies.

23 Experiments and Future Validation

1. **Step 1 — Mathematical consistency:** Verify dose-response saturation and tolerance emergence from AFL equations under parameter sweep (complete — toy model in `outputs/paper_09/`).
2. **Step 2 — Synthetic perturbation:** Reproduce known PK/PD curve shapes ($E_{\max}$, Hill equation) as special cases of the AFL constraint formulation.
3. **Step 3 — Retrospective alignment:** Apply AFL constraint axes to published dose-response datasets; test whether cross-axis combinations show lower EC50 shifts than same-axis combinations.
4. **Step 4 — In vitro / ex vivo:** Measure receptor occupancy, downstream signalling, and metabolic cost simultaneously in cell-based assays; fit C_S , C_E , C_M to data.
5. **Step 5 — Prospective observational:** Track tolerance onset time, drug holiday recovery, and combination synergy in clinical cohorts with serial biomarker monitoring.

6. **Step 6 — Clinical utility:** Pre-specified AFL-guided combination selection versus empirical prescribing in a randomised pharmacology study; primary endpoint: time to dose-limiting toxicity vs response.

Validation ladder status: Step 1 complete; Steps 2–6 open.

24 Clinical Safety Boundary

This paper contains no prescribing instructions, drug recommendations, or dose calculations. No AFL output should be used to guide drug selection, dosing, combination therapy, or discontinuation without: (a) full pharmacokinetic/pharmacodynamic modelling; (b) adverse-event monitoring; (c) organ-function assessment; (d) guideline-concordant prescribing supported by trial evidence. AFL constraint variables are candidate hypotheses for future PK/PD modelling — they are not validated clinical decision tools.

25 Runtime Diagnostic

The release-local pharmacology diagnostic is `supplementary/Pharmacological_Synergy_Engine.py`. It produces toy model outputs in `outputs/paper_09/`: same-axis versus cross-axis intervention comparison, saturation-regime and adaptive-resistance trajectory plots, and a `summary.json` with parameter values and regime classifications.

The outputs support a modelling distinction (same-axis vs cross-axis intervention), not a prescribing rule. All outputs carry the label “toy diagnostic — not a clinical claim.”

26 Conclusion

Drug response emerges from the interaction between administered exposure and the adaptive constraint environment through which that exposure must act. AFL separates administered dose, tissue exposure, and constrained biological response — three quantities often conflated in clinical reasoning. Tolerance, resistance, and saturation correspond to constraint dynamics: rising C over time, compensatory M_s accumulation, and EC50 ceiling respectively.

AFL does not replace pharmacokinetics, pharmacodynamics, or clinical trial evidence. It offers a systems framing for generating hypotheses about why monotherapy fails in multi-constraint disease, why dose escalation can worsen outcomes, and why timing and recovery windows may be as important as drug class. These hypotheses require prospective validation before influencing prescribing.

27 Reproducibility Note

Release-local script: `supplementary/Pharmacological_Synergy_Engine.py`. Dependencies: NumPy, matplotlib. Runtime < 2s. All parameters and random seeds are set in-file. Outputs written to `outputs/paper_09/`. No proprietary data required.

28 Dose–Constraint Control

The key pharmacological extension is that dose is not identical to effective flux. A high administered dose can fail if absorption, distribution, target access, receptor availability, downstream signaling,

or organ reserve imposes constraint. Conversely, a lower dose can be more effective when timing, tissue access, and constraint recovery are favorable.

AFL therefore separates three quantities:

$$D_{admin} \rightarrow J_{exposure} \rightarrow J_{effect}/C_{response}. \quad (20)$$

The first is what is prescribed. The second is what reaches the relevant compartment. The third is the constrained biological response. Many clinical failures occur because these three quantities are treated as one.

29 Therapeutic Regimes

Four regimes follow. In the underexposure regime, increasing dose can improve response. In the saturation regime, more dose mainly increases toxicity. In the adaptive-resistance regime, repeated exposure increases C and lowers response over time. In the recovery-sensitive regime, pulsed or cyclic dosing may outperform continuous exposure by allowing C to relax.

30 Combination Therapy

Combination therapy is rational when each drug targets a different constraint. One agent may reduce source flux, another may lower resistance, another may prevent compensatory escape, and another may protect organ reserve. AFL predicts synergy when the combined effect lowers the operating ratio Ψ_s without forcing a new constraint elsewhere.

31 Measurable Tests

The pharmacology paper should be tested with time-series data: drug concentration, pharmacodynamic effect, toxicity markers, resistance markers, and recovery time. The framework fails if adaptive constraint variables do not explain tolerance, resistance, or diminishing returns better than static dose-response curves.

Conflict of Interest

The author declares no conflict of interest.

Funding

This research received no external funding.

Author Contributions

Conceptualization, formal analysis, runtime engineering, and manuscript preparation: S.B.M.

Data Availability

All computational models, Python scripts, and numerical verifications are explicitly available in the `/supplementary` repository layer and the `cdfd.runtime` engine.

32 Reference Layer

References

Robert M. May. Simple mathematical models with very complicated dynamics. *Nature*, 261:459–467, 1976. doi: 10.1038/261459a0.