

Paper BIO-11: Adaptive Flux Limitation in Bioelectric Regulation: A CDFD Model of Ion Fields as Constraint Controllers

Steve Bico Mujjabi
Independent Researcher

May 22, 2026

Adaptive Flux Limitation in Bioelectric Regulation: A CDFD Model of Ion Fields as Constraint Controllers

Abstract

This paper presents **Adaptive Flux Limitation (AFL)** as the Part III biology-and-medicine model inside Constraint-Driven Flux Dynamics (CDFD). CDFD supplies the flow-constraint-memory grammar: physiological flow Φ , constraint C , surface responsiveness S , and structural memory M_s . The Runtime citation anchors the CDFL implementation, while clinical interpretation remains separate from software output and bedside validation.

Bioelectricity — the ensemble of membrane potentials, ion gradients, and inter-cellular electrical coupling across tissues — is conventionally studied as a property of excitable cells (neurons, cardiomyocytes). We argue that bioelectric fields constitute a **global constraint control layer** governing biological flux in all cell types: they regulate metabolic throughput, developmental patterning, tissue organization, and regeneration capacity.

We develop an AFL model in which membrane potential V_m is the primary constraint controller ($C_{\text{mem}} = f(V_m)$), gap junction coupling propagates constraint signals spatially, and bioelectric pattern disruption underlies a wide range of pathologies from cancer to neurodegeneration to impaired wound healing. Key results: (1) hyperpolarization raises C_{mem} , suppressing metabolic flux; depolarization lowers C_{mem} , enabling throughput — explaining why depolarized cells are more proliferative; (2) bioelectric developmental patterning is a spatial AFL constraint field that guides morphogenesis independently of and prior to gene expression cascades; (3) bioelectric therapies can reset C_{mem} in a targeted manner, offering a non-genetic intervention route for regeneration and cancer suppression.

Status: Inferred

Within the extended framework of this series, biological networks are modeled as *Adaptive Surfaces* that dynamically integrate physiological load and retain structural memory. The system's health is governed by the adaptive ratio $\Psi_s = (\Phi/C) \cdot S \cdot M_s$, where homeostasis ($\Psi_s \approx 1.0$) is maintained near the stable attractor ($S \cdot M_s \approx 1.0$), and disease emerges as surface dysregulation when maladaptive memory locks tissues into chronic constraint.

Greek-symbol convention. Unless a manuscript states a tighter equation-local meaning, Greek symbols are local CDFD variables or coefficients, not universal constants. Here Φ denotes driving flux, Ψ_s denotes the adaptive operating ratio, Ω denotes a domain or state space, and Λ denotes the Life Number where used. The coefficients α , β , and γ denote accumulation or reinforcement, relaxation or decay, and diffusion, coupling, or redistribution. The symbols δ and ϵ denote perturbation or tolerance scales, while η , λ , μ , ν , ρ , σ , τ , and ϕ denote local efficiency, rate, baseline, frequency, density or correlation, noise or variance, time-scale, and phase or field terms. Subscripts restrict any coefficient to the named pathway, tissue, domain, or experiment.

Clinical Reader's Guide

This paper brings bioelectric regulation into the AFL sequence. Electrical gradients, membrane potentials, ion fluxes, gap junctions, and tissue voltage patterns are not deco-

rations on biology; they are part of how cells coordinate growth, repair, excitability, and pattern.

In AFL notation, bioelectric states can be read as constraints on signaling throughput. A depolarized or poorly coupled tissue may have altered C for communication even before gross anatomy changes. Memory M_s can include persistent channel-expression states, scar architecture, or stable patterning errors.

The clinical relevance is still early. Bioelectric interventions require careful distinction between established therapies, experimental tissue-patterning work, and speculative applications. This paper should therefore be read as a research framework for mapping electrical state to flow-constraint dynamics.

AFL notation. In this paper, Φ denotes the relevant physiological throughput, C the operative biological constraint, S surface responsiveness, and M_s retained structural memory. When a dominant flow can be specified, $\Psi_s = (\Phi/C) \cdot S \cdot M_s$ denotes the operating ratio. Claim discipline: explicit assumptions, separated derivation vs. interpretation, and status labels.

1 Introduction

Adaptive Flux Limitation (AFL) is the named model used in this paper; CDFD is the parent framework that supplies the variables, closure logic, and claim discipline. The biological or medical question is posed first as AFL, then interpreted as a CDFD model of flow, constraint, surface responsiveness, and structural memory.

Bioelectricity is often introduced through excitable tissues, yet membrane potential and ion flux are older and broader than nerves or myocardium. Every living cell maintains electrochemical gradients. Those gradients influence transport, metabolism, proliferation, polarity, migration, gap-junction coupling, wound repair, and developmental pattern. The question for AFL is whether electrical state can be treated as a constraint layer rather than only a signal layer.

In this paper, V_m is read as a regulator of membrane and tissue constraint. Hyperpolarized, well-coupled, and spatially coherent states can stabilize quiescence or differentiation; depolarized or poorly coupled states can permit growth, migration, or failed patterning. The claim is not that voltage replaces genes, morphogens, mechanics, or metabolism. It is that voltage helps set the boundary conditions under which those systems can act.

This distinction matters because bioelectric memory may persist after the initiating injury has passed. Channel expression, gap-junction architecture, scar structure, and stable pattern errors can preserve an abnormal constraint field. That makes bioelectric regulation relevant to development, regeneration, cancer biology, wound healing, and neurological disease without turning it into a universal cure narrative.

The clinical boundary remains conservative. Established electrophysiology and approved electroceutical therapies should not be confused with speculative tissue-patterning applications. AFL is used here to organize experiments that connect voltage state, flux, constraint, and recovery.

Status: Hypothesis

2 Membrane Potential as C_{mem}

$$C_{\text{mem}} = f(V_m) = C_0 \cdot e^{-\kappa(V_m - V_{\text{rest}})} \quad (1)$$

where C_0 is basal membrane constraint, κ is voltage sensitivity, and V_{rest} is the resting potential.

- **Hyperpolarization** ($V_m < V_{\text{rest}}$): $C_{\text{mem}} \uparrow$ — ion channels close, metabolic coupling decreases, cell enters quiescent low-throughput state ($\Psi_{s,\text{cell}} < 1$).
- **Depolarization** ($V_m > V_{\text{rest}}$): $C_{\text{mem}} \downarrow$ — ion channels open, metabolic coupling increases, cell enters high-throughput state ($\Psi_{s,\text{cell}} > 1$).

Resting V_m in non-excitabile cells correlates with proliferative state: cancer cells are typically more depolarized (-20 to -40 mV) than quiescent differentiated cells (-60 to -90 mV). In AFL terms: depolarization lowers C_{mem} , enabling the high- Ψ_s growth state.

Status: Inferred

3 Spatial Constraint Propagation via Gap Junctions

Gap junctions (connexins) electrically couple adjacent cells, allowing V_m and small molecules to propagate spatially. In AFL terms:

$$\nabla C_{\text{mem}}(\mathbf{x}) = -D_{\text{gap}} \nabla^2 V_m(\mathbf{x}) \quad (2)$$

A bioelectric gradient across a tissue creates a spatial AFL constraint field. Cells in regions of high V_m (depolarized) have low C_{mem} and high Ψ_s — prone to growth. Cells in hyperpolarized regions have high C_{mem} and low Ψ_s — quiescent.

Consequence: disrupting gap junction coupling (e.g., by Cx43 mutations) destroys the spatial constraint gradient, creating local deregulation of growth. This is consistent with the tumour-suppressive role of connexins in multiple cancer types.

Status: Inferred

4 Developmental Patterning as AFL Spatial Constraint

In embryonic development, bioelectric gradients are established before gene expression patterns are visible. Levin’s group has demonstrated that the spatial pattern of V_m across the embryo predicts the position of organs and body structures.

AFL interpretation: the bioelectric pattern is a spatial AFL constraint field that specifies **where** growth (high Ψ_s) and differentiation (low Ψ_s) are to occur. Gene expression cascades then implement this pattern — but the pattern itself is bioelectrically encoded.

$$\Psi_{s,\text{cell}}(\mathbf{x}) = \frac{\Phi_{\text{growth}}}{C_{\text{mem}}(V_m(\mathbf{x}))} \quad (3)$$

Regions where $\Psi_s > 1$: proliferating progenitors. Regions where $\Psi_s < 1$: differentiating post-mitotic cells. The bioelectric pattern is the AFL blueprint for morphogenesis.

Status: Inferred

5 Regeneration: Resetting the Constraint Field

Planaria (flatworms) regenerate entire heads from tail fragments. Salamanders regenerate limbs. Humans do not regenerate limbs. The AFL explanation: regeneration requires re-establishing the embryonic bioelectric constraint field pattern in the injured tissue.

In planaria, wound healing is accompanied by a bioelectric depolarization wave that propagates from the wound site, temporarily lowering C_{mem} and raising Ψ_s in the blastema, enabling dedifferentiation and proliferation. When this wave is pharmacologically blocked, regeneration fails.

In mammals, the equivalent wave is much weaker and shorter-lived. AFL prediction: amplifying the wound-depolarization wave (via pharmacological ion channel modulation, bioelectric scaffolds, or direct electrical stimulation) should extend the regenerative window. This is consistent with experimental results in mammalian spinal cord and peripheral nerve regeneration models.

Status: Inferred

6 Bioelectricity in Disease

6.1 Cancer as Depolarization-Mediated Constraint Escape

Cancer cells are characteristically depolarized. This is not a consequence of cancer but a **driver**: experimental depolarization of normal cells (by overexpressing depolarizing channels) is sufficient to induce tumour-like proliferation in some model systems.

AFL mechanism: depolarization $\Rightarrow C_{\text{mem}} \downarrow \Rightarrow \Psi_{s,\text{growth}} > 1 \Rightarrow$ uncontrolled proliferation.

Therapeutic implication: hyperpolarizing tumour cells (by activating K^+ channels or Cl^- channels) should raise C_{mem} and suppress $\Psi_{s,\text{growth}} < 1$, potentially arresting growth without targeting specific oncogenes.

Status: Inferred

6.2 Neurodegeneration: Loss of Neural Bioelectric Constraint Architecture

In Parkinson's disease, dopaminergic neurons in the substantia nigra progressively lose neuromelanin and mitochondrial function. In AFL terms:

- Neuromelanin loss: $S \uparrow$ (the melanin-like stabilization channel from OOL-3 applies at the neural level)
- Mitochondrial dysfunction: $\sigma_e \downarrow$, reducing energetic capacity to maintain V_m
- C_{mem} dysregulation: unstable $V_m \Rightarrow$ irregular constraint field $\Rightarrow$ abnormal firing patterns

The pathological tremor and bradykinesia of PD are AFL phenomena: the neural constraint architecture has degraded to the point where the system cannot maintain $\Psi_{s,\text{neural}} \approx 1$ stably.

Status: Inferred

6.3 Wound Healing: Injury Current as Flux Signal

Tissue injury creates an **injury current**: the disruption of the transepithelial potential generates an electrical field at the wound edge. This field (Φ_{injury}) drives directional migration of keratinocytes and fibroblasts toward the wound (galvanotaxis).

In AFL terms: the injury current is a Φ signal that activates directed flux ($J_{\text{migration}}$) toward the wound. The constraint on this flux is the tissue ECM stiffness and cellular actomyosin contractility (C_{ECM}). When injury current is disrupted (e.g., in diabetic neuropathy with impaired transepithelial potential), wound healing fails — an AFL wound healing deficit.

Status: Inferred

7 Bioelectric Therapeutics: Resetting C_{mem}

Intervention	AFL mechanism	Application
Direct current stimulation	Raises Φ_{elec} , modulates C_{mem} spatially	Wound healing, bone re
Ion channel modulators	Changes V_m directly ($C_{\text{mem}} = f(V_m)$)	Cancer suppression, reg
Optogenetics	Light-controlled Φ_{ion} injection	Neural circuit restoratio
Bioelectric scaffolds	Sustained Φ_{elec} gradient in 3D	Tissue engineering
Transcranial magnetic stimulation	Modulates C_{neural} via induced currents	Neurological/psychiatric

Status: Inferred

8 Flux–Constraint Regime

V_m state	C_{mem}	$\Psi_{s,\text{cell}}$	Biological state
Deeply hyperpolarized (< -80 mV)	Very high	$\ll 1$	Quiescent, senescent
Resting (-60 to -80 mV)	Normal	≈ 1	Homeostatic
Mildly depolarized (-40 to -60 mV)	Reduced	$\approx 1-2$	Proliferating
Significantly depolarized (> -40 mV)	Low	> 2	Cancer-like, injured

Status: Inferred

9 Predictions

1. Hyperpolarizing ion channel agonists should suppress proliferation in cancer cell lines, with effect size proportional to the degree of baseline depolarization (starting C_{mem} deficit).
2. Embryonic bioelectric pattern disruption (via pharmacological gap junction blockade) should predictably misplace morphogenetic fields in a manner consistent with spatial AFL constraint field maps.
3. Wound healing rate should correlate with injury current magnitude in longitudinal measurements across wound types.

4. Neurodegeneration progression rate should correlate with mitochondrial σ_e decline (measurable via Complex I activity) before clinical symptoms appear.
5. Bioelectric scaffold-enhanced tissue engineering should show regeneration rates proportional to the amplitude of the imposed Φ_{elec} gradient.

Status: Open

10 Falsifiability

The AFL bioelectric model would be falsified if: cell proliferation is uncorrelated with V_m across cell types; embryonic bioelectric disruption produces random (not spatially predicted) morphological defects; wound healing rate is independent of injury current magnitude; or hyperpolarizing drugs have no anti-proliferative effect in depolarized cancer cells.

Status: Open

11 Clinical Mapping of Symbols

Symbol	Bioelectric meaning	Measurement proxy	Invalidation condition
Φ	Ion flux / metabolic throughput	Patch-clamp current; metabolic rate	If Φ cannot be independently measured from C_{mem} , the mapping is circular
C_{mem}	Membrane constraint ($= f(V_m)$)	Membrane potential (patch clamp, VSDs)	If V_m change produces no measurable flux change, C_{mem} as constraint fails
S	Surface responsiveness	Channel density; gap junction coupling index	If coupling index is uncorrelated with patterning outcomes, S adds no value
M_s	Memory / locked channel state	Persistent channel expression; scar impedance	If M_s relaxes instantly on threat removal, memory framing is unnecessary
Ψ_s	Operating state	Computed proxy only — no direct assay	If Ψ_s trajectories match random walks, the model is not informative

Status: Hypothesis

12 Known Medicine Baseline

Established bioelectric medicine includes:

- **Cardiac electrophysiology:** pacemakers, defibrillators, antiarrhythmics — mature clinical practice built on action potential dynamics, not AFL framing.

- **Neurostimulation:** deep brain stimulation (Parkinson’s disease, OCD), spinal cord stimulation (pain), transcranial magnetic stimulation (depression) — approved therapies with evidence from controlled trials.
- **Wound healing:** bioelectric wound dressings (electrostimulation) — early evidence in chronic wound management; mechanisms partially characterised.
- **Cancer bioelectrics:** experimental; Levin lab and collaborators have shown V_m manipulation influences tumour-like growth in some model organisms — no approved clinical translation yet.

AFL is layered on top of established electrophysiology, not a replacement for it. The $C_{\text{mem}} = f(V_m)$ mapping is a research hypothesis, not a clinical tool.

13 Seizure Threshold and Neural Criticality

Epilepsy can be read in AFL terms as a failure of inhibitory constraint maintenance. In a critical network, the balance between excitatory flux Φ_{exc} and inhibitory constraint C_{inh} determines whether a perturbation is absorbed or amplifies into a seizure.

$$\Psi_{s,\text{neural}} = \frac{\Phi_{\text{exc}}}{C_{\text{inh}}} \cdot S \cdot M_s \quad (4)$$

Near the seizure threshold: $\Psi_{s,\text{neural}} \approx 1^-$, small perturbations tip the network into runaway excitation ($\Psi_s \gg 1$). Anticonvulsant drugs increase C_{inh} (GABA-A agonists, Na^+ channel blockers) or reduce Φ_{exc} (NMDA antagonists). Memory M_s could represent kindling — the progressive lowering of seizure threshold with repeated events.

Status: Hypothesis. Validation: EEG biomarker correlation, dose-response in animal models.

14 Deep Analysis: Neural Plasticity and Constraint-Governed Network Reorganization

Synaptic plasticity (LTP/LTD) can be read as adaptive C restructuring: LTP lowers C_{syn} , increasing throughput in a pathway; LTD raises C_{syn} , reducing it. The memory parameter M_s then encodes the persistence of these changes — structural synaptic changes (spine growth, receptor insertion) are the AFL memory lock.

Neurodegeneration disrupts this architecture: loss of synapses, dendritic pruning, and white matter lesions progressively raise C_{neural} across circuits, reducing $\Psi_{s,\text{neural}}$ until function falls below threshold.

Status: Hypothesis. No AFL-specific predictions beyond what standard computational neuroscience already models.

15 Experiments and Future Validation

1. **Mathematical consistency:** $C_{\text{mem}}(V_m)$ monotonicity, finite Ψ_s at resting potential, sensitivity to kappa parameter.

2. **Synthetic perturbation:** release-local script (`supplementary/supplementary_afl_11.py`) — pacemaker oscillation, spatial gap junction propagation, therapeutic reset toy.
3. **In vitro:** patch-clamp measurement of metabolic flux vs V_m across a panel of cell lines; gap junction manipulation (18- α -glycyrrhetic acid) and spatial Ψ_s proxy change.
4. **Ex vivo:** organoid bioelectric assays; voltage-sensitive dye (VSD) imaging during morphogen challenge.
5. **Retrospective:** ECG / EEG datasets — test whether $\Psi_{s,\text{neural}}$ proxy correlates with seizure frequency.
6. **Prospective observational:** wound healing rate vs injury current magnitude in chronic wound cohort.
7. **Clinical utility:** only after steps 1–6 are completed, with safety data and ethics approval.

Success criterion: V_m correlates with metabolic flux rate (patch clamp) after controlling for direct ion current effects. **Failure criterion:** no correlation — C_{mem} as AFL constraint adds no explanatory power.

16 Clinical Safety Boundary

This paper is a research framework, not a clinical tool.

- No bioelectric intervention should be modified based on AFL framing without established clinical evidence.
- Cardiac electrophysiology, neurostimulation, and wound bioelectrics are governed by device regulations, clinical trials, and specialist expertise.
- Experimental bioelectric cancer or regeneration therapies require dedicated phase I/II safety and efficacy trials.
- This model does not replace electrophysiology training, EEG interpretation, or device safety assessment.

17 Conclusion

Bioelectricity is proposed as a global AFL constraint control layer — a research hypothesis that membrane potential sets C_{mem} , gap junctions propagate constraint spatially, and the resulting bioelectric field shapes development, regeneration, and disease. This positioning is distinct from established cardiac/neural electrophysiology and requires systematic experimental validation before any clinical translation.

Status: Inferred

18 Runtime Diagnostic

Release-local script: `supplementary/supplementary_afl_11.py`. Outputs in `outputs/paper_11/`: `membrane_constraint_profiles.csv` (voltage table), `pacemaker_dynamics.csv`, `spatial_bioelectric_reset_scenario.csv`, `bioelectric_regulation.png`. All outputs are toy diagnostics under stated assumptions. No clinical inference should be drawn from them.

Status: Derived

19 Reproducibility Note

- **Script:** `supplementary/supplementary_afl_11.py` (NumPy + matplotlib; seed fixed; runtime < 2s)
- **Dependencies:** NumPy, matplotlib (Agg backend)

Status: Derived

Adaptive Flux Limitation in Bioelectric Regulation: Ion Fields as Global Constraint Controllers in Living Systems

References