

Paper BIO-17: Adaptive Flux Limitation in Therapy and Medical Synthesis: A CDFD Model for Testable Intervention Hypotheses

Steve Bico Mujjabi
Independent Researcher

May 22, 2026

Adaptive Flux Limitation in Therapy and Medical Synthesis: A CDFD Model for Testable Intervention Hypotheses

Abstract

This paper presents **Adaptive Flux Limitation (AFL)** as the Part III biology-and-medicine model inside Constraint-Driven Flux Dynamics (CDFD). CDFD supplies the flow-constraint-memory grammar: physiological flow Φ , constraint C , surface responsiveness S , and structural memory M_s . The Runtime citation anchors the CDFL implementation, while clinical interpretation remains separate from software output and bedside validation.

Conventional pharmacology targets specific molecules: receptor agonists increase flux, antagonists reduce it, enzyme inhibitors raise constraint at a specific step. The AFL framework provides a **meta-level intervention taxonomy** that classifies all treatments by their effect on the flux-constraint balance ($\Psi_s = (\Phi/C) \cdot S \cdot M_s$) rather than their molecular target.

Three AFL intervention strategies exist: (1) **Flux enhancement** for $\Psi_s < 1$ states (under-delivery, under-function); (2) **Constraint reduction** for $C \uparrow$ states (resistant, fibrotic, inflammatory diseases); (3) **Flux limitation** for $\Psi_s > 1$ states (overload, toxicity, cancer). A fourth strategy — **System stabilization** — is applicable when Ψ_s is oscillating (unstable) rather than statically displaced.

We demonstrate that this taxonomy explains why certain treatment combinations are synergistic (they target different AFL axes), why monotherapy fails in multi-constraint diseases (only one axis targeted), and how dietary, lifestyle, and pharmacological interventions can be systematically designed for specific Ψ_s states.

Status: Inferred

Within the extended framework of this series, biological networks are modeled as *Adaptive Surfaces* that dynamically integrate physiological load and retain structural memory. The system's health is governed by the adaptive ratio $\Psi_s = (\Phi/C) \cdot S \cdot M_s$, where homeostasis ($\Psi_s \approx 1.0$) is maintained near the stable attractor ($S \cdot M_s \approx 1.0$), and disease emerges as surface dysregulation when maladaptive memory locks tissues into chronic constraint.

Greek-symbol convention. Unless a manuscript states a tighter equation-local meaning, Greek symbols are local CDFD variables or coefficients, not universal constants. Here Φ denotes driving flux, Ψ_s denotes the adaptive operating ratio, Ω denotes a domain or state space, and Λ denotes the Life Number where used. The coefficients α , β , and γ denote accumulation or reinforcement, relaxation or decay, and diffusion, coupling, or re-distribution. The symbols δ and ϵ denote perturbation or tolerance scales, while η , λ , μ , ν , ρ , σ , τ , and ϕ denote local efficiency, rate, baseline, frequency, density or correlation, noise or variance, time-scale, and phase or field terms. Subscripts restrict any coefficient to the named pathway, tissue, domain, or experiment.

Clinical Reader's Guide

This synthesis paper gathers the therapeutic implications of the series while keeping a strict boundary: AFL can organize hypotheses about intervention axes, but it is not a

treatment protocol. The four broad axes are increasing useful flow, reducing pathological constraint, improving responsiveness, and clearing maladaptive memory.

The clinical value is in asking better questions. Is a patient under-driven, over-constrained, poorly responsive, or locked by history? Is a drug pushing the same axis as another drug, or does it relieve a different bottleneck? Is a plateau a sign of treatment failure, or the moment when the next constraint became dominant?

The paper should be read as a research synthesis. Future work would need prospective studies, safety analysis, clinical endpoints, and comparison with standard care before any algorithmic use.

AFL notation. In this paper, Φ denotes the relevant physiological throughput, C the operative biological constraint, S surface responsiveness, and M_s retained structural memory. When a dominant flow can be specified, $\Psi_s = (\Phi/C) \cdot S \cdot M_s$ denotes the operating ratio. Claim discipline: explicit assumptions, separated derivation vs. interpretation, and status labels.

1 Introduction

Adaptive Flux Limitation (AFL) is the named model used in this paper; CDFD is the parent framework that supplies the variables, closure logic, and claim discipline. The biological or medical question is posed first as AFL, then interpreted as a CDFD model of flow, constraint, surface responsiveness, and structural memory.

Therapy is the deliberate alteration of a living system’s operating state. A treatment may increase a missing flow, reduce a harmful driver, lower a pathological constraint, raise a protective constraint, restore responsiveness, or help a tissue shed maladaptive memory. The same drug can be useful or harmful depending on which axis is limiting the patient at that time.

AFL offers a synthesis language for that problem. It asks whether the target state is under-delivery, overload, poor responsiveness, oscillation, or memory lock. It then asks whether an intervention acts on Φ , C , S , M_s , or the recovery terms that move the system back toward stability. This is not a prescribing algorithm. It is a way to classify why combinations work, why monotherapy plateaus, and why an apparently logical escalation can fail.

The medical standard is unchanged. Treatment decisions require diagnosis, staging, contraindication review, evidence hierarchy, patient preferences, specialist judgement, and monitoring for harm. AFL is useful only if it generates testable hypotheses about intervention timing, synergy, plateau interpretation, and recovery trajectories.

This final paper therefore closes Part III as a research program rather than a therapeutic manifesto. It connects metabolism, CKD, inflammation, oncology, pharmacology, bioelectric regulation, biomarkers, and systems medicine under one question: which constraint axis is actually governing recovery?

Status: Derived

$$\text{Treatment goal: } \Psi_{s,\text{target}} \approx 1 \tag{1}$$

- If $\Psi_s < 1$ (under-delivered): increase Φ or decrease the limiting C .
- If $\Psi_s > 1$ (overloaded): reduce the harmful flow, redistribute load, or relieve the compartment-specific constraint that traps flux.

- If Ψ_s oscillates: stabilize the dynamical system by reducing excessive gain or improving recovery.

2 Strategy 1: Flux Enhancement ($\Psi_s < 1$ States)

2.1 Clinical contexts

$\Psi_s < 1$ arises when physiological function is inadequate for demand:

- Heart failure: $\Psi_{s,\text{cardiac}} = \text{CO}/\text{SVR} < 1$ — insufficient cardiac output
- Hypothyroidism: $\Phi_{\text{T}_3} \downarrow$ — insufficient metabolic flux
- Type 1 diabetes: $\Phi_{\text{insulin}} = 0$ — no insulin flux
- Anaemia (EPO deficiency): $\Phi_{\text{EPO}} \downarrow$ — insufficient erythropoietic drive
- Adrenal insufficiency: $\Phi_{\text{cortisol}} \downarrow$ — insufficient stress flux

2.2 Flux enhancement interventions

Intervention	Φ increased	Mechanism
Insulin replacement (T1DM)	Φ_{insulin}	Direct flux restoration
EPO/darbepoetin	Φ_{EPO}	Signal amplification
Levothyroxine	Φ_{T_3}	Hormone replacement
Inotropes (dobutamine)	Φ_{cardiac}	Myocardial flux drive
Vasopressors (in low SVR states)	$\Phi_{\text{perfusion}}$	Vascular pressure restoration
Blood transfusion	Φ_{O_2}	Direct flux delivery
IV iron	Φ_{Fe}	Removes constraint on erythropoiesis
Corticosteroid replacement	Φ_{cortisol}	Adrenal axis flux

Status: Derived

3 Strategy 2: Constraint Reduction ($C \uparrow$ States)

3.1 Clinical contexts

$C \uparrow$ arises when the system's resistance to normal flux has increased pathologically:

- insulin resistance (chronic C_E upregulation) (T2DM): $C_{\text{met}} \uparrow$
- CKD: $C_{\text{renal}} \uparrow$
- Hypertension: $C_{\text{vascular}} \uparrow$ (elevated SVR)
- Heart failure (HFpEF): $C_{\text{myocardial}} \uparrow$ (diastolic stiffness)
- Chronic inflammation (system-wide constraint C amplification): $C_{\text{system}} \uparrow$
- COPD: $C_{\text{airway}} \uparrow$ (airflow resistance)
- Fibrotic disease: $C_{\text{ECM}} \uparrow$

3.2 Constraint reduction interventions

Intervention	C reduced	Mechanism
ACE inhibitors / ARBs	C_{vascular}	Vasodilation, RAAS blockade
Metformin	C_{met} (partial)	AMPK activation, reduces hepatic g
GLP-1 agonists	$C_{\text{met}}, C_{\text{liver}}$	Reduces liver fat, appetite, gastric e
Anti-inflammatory agents	C_{inflam}	Removes inflammation (system-wide
Bronchodilators	C_{airway}	Reduces airflow resistance
Diuretics (in oedema)	C_{preload}	Reduces ventricular wall stress
Antifibrotic agents (pirfenidone)	C_{ECM}	Reduces fibrotic constraint progressi
Calcimimetics (in CKD)	C_{PTH}	Reduces PTH-driven bone constrain
Iron chelation (overload)	$C_{\text{Fe-toxicity}}$	Removes iron-mediated constraint
SGLT2 inhibitors	$C_{\text{glucose-load}}$	Reduces glycaemic constraint via gly
Statins	C_{vascular} (anti-inflammatory)	Lowers vascular constraint

Status: Derived

4 Strategy 3: Flux Limitation ($\Psi_s > 1$ States)

4.1 Clinical contexts

$\Psi_s > 1$ arises when flux exceeds the system's capacity to handle it safely:

- Hypertensive crisis: $\Phi_{\text{pressure}}/C_{\text{vascular}} \gg 1$
- Hyperglycaemia: $\Phi_{\text{glucose}} > C_{\text{met}}$ (T2DM decompensation)
- Fluid overload (pulmonary oedema): $\Phi_{\text{fluid}} > C_{\text{ventricular}}$
- Toxicity (drug overdose): $\Phi_{\text{toxin}} > C_{\text{hepatic/renal}}$
- Cytokine storm: $J_{\text{immune}} \gg C_{\text{tissue}}$
- Cancer (proliferative overload): $\Phi_{\text{growth}}/C_{\text{cell}} \gg 1$

4.2 Flux limitation interventions

Intervention	Φ limited	Mechanism
Antihypertensives	Φ_{BP}	Reduces vascular flux pressure
Dietary restriction / fasting	$\Phi_{\text{glucose}}, \Phi_{\text{lipid}}$	Reduces substrate flux input
Dialysis	$\Phi_{\text{toxins}} \downarrow$ (clearance $\uparrow$)	Removes accumulated flux products
Antidotes / activated charcoal	Φ_{toxin} (absorption)	Prevents flux entry
Diuretics (in overload)	Φ_{fluid}	Removes excess volume flux
Chemotherapy (cytostatic)	$\Phi_{\text{proliferation}}$	Limits tumour growth flux
IL-6 blockers (cytokine storm)	Φ_{cytokine}	Limits immune flux signal

Status: Derived

5 Strategy 4: System Stabilization (Oscillating Ψ_s)

Some disease states feature oscillating Ψ_s rather than sustained displacement:

- Cardiac arrhythmia: $\Psi_{s,\text{cardiac}}$ oscillates due to electrical constraint instability
- Epilepsy: $\Psi_{s,\text{neural}}$ oscillates between excitatory ($\Psi_s > 1$) and post-ictal suppression ($\Psi_s < 1$) states
- Intermittent claudication: $\Psi_{s,\text{peripheral}}$ oscillates with activity
- Hypoglycaemia episodes in T1DM: $\Psi_{s,\text{met}}$ oscillates due to insulin timing

Stabilization interventions reduce the amplitude ($\Delta\Psi_s$) of oscillation rather than shifting the mean:

- Anti-arrhythmics: increase $C_{\text{cardiac-electrical}}$ stability
- Anti-epileptics: raise the neural firing threshold ($C_{\text{neural}} \uparrow$ or reduce $\Phi_{\text{excitatory}}$)
- Continuous glucose monitoring + closed-loop insulin: reduces glycaemic oscillation amplitude

Status: Inferred

6 The AFL Treatment Algorithm

Step 1: Measure Psi (or best available proxy)

Step 2: Identify dominant constraint(s) (elevated C markers)

Step 3: Identify flux state (reduced or elevated Phi markers)

Step 4: Select strategy:

```
if Psi < 1 AND C is normal: -> flux enhancement
if Psi < 1 AND C is elevated: -> constraint reduction FIRST,
                                then consider flux enhancement

if Psi > 1: -> flux limitation
if Psi oscillates: -> stabilization
```

Step 5: Apply combination therapy targeting multiple axes

Step 6: Monitor Psi trajectory (not just individual biomarkers)

Step 7: Adjust as Psi evolves

Why Step 4 matters: In CKD anaemia, $\Psi_{\text{erythro}} < 1$ but $C_{\text{iron}} + C_{\text{inflam}}$ are elevated. Flux enhancement (ESA alone) fails because C remains high. The correct AFL strategy is constraint reduction first (IV iron, anti-inflammatory) then flux enhancement (ESA). This is precisely what evidence-based guidelines now recommend.

Status: Derived

7 Synergy Prediction from AFL Axes

Drug A	Drug B	AFL axes	Prediction
ESA	IV iron	$\Phi \uparrow + C_{\text{Fe}} \downarrow$	Synergistic (dual axis)
Metformin	Exercise	$C_{\text{met}} \downarrow + \beta_{\text{met}} \uparrow$	Synergistic (different parameters)
GLP-1 agonist	SGLT2i	$C_{\text{liver}} \downarrow + \Phi_{\text{glucose}} \downarrow$	Synergistic (complementary)
ACE inhibitor	ARB	$C_{\text{vascular}} \downarrow$ (same axis)	Additive (not synergistic; same axis)
Platinum chemo	Anti-PD1	$C_{\text{tumour}} \uparrow + C_{\text{immune} \rightarrow \text{tumour}} \downarrow$	Synergistic (dual axis)

AFL predicts: combinations that target different AFL axes are synergistic; combinations targeting the same axis are additive at best.

Status: Inferred

8 Fasting and OMAD as the Universal AFL Reset

Fasting is a pure constraint recovery intervention: it reduces $\Phi_{\text{metabolic}}$ to near-zero, allowing $\beta_{\text{met}} C_{\text{met}}$ to dominate Equation $\dot{C} = \alpha\Phi - \beta C$, driving $C_{\text{met}} \rightarrow 0$ exponentially.

Fasting benefits documented across diverse conditions:

- T2DM: HOMA-IR improvement (metabolic $C \downarrow$)
- Autoimmunity: reduced inflammatory markers ($C_{\text{immune}} \downarrow$)
- Cancer (adjunct): improved chemotherapy sensitivity (reduces tumour ATP flux)
- Neurological: improved cognitive function (reduces neuroinflammatory C)
- CKD (mild): reduces uraemic toxin accumulation between meals

The AFL interpretation: fasting works broadly because it interrupts Φ -driven constraint accumulation across *all* organ systems simultaneously, allowing the universal βC recovery term to act everywhere at once.

Status: Inferred

9 Predictions

1. Drug combinations targeting different AFL axes should show synergistic (supraadditive) effect in clinical trials; same-axis combinations should show only additive effects.
2. Pre-treatment Ψ_s profiling should predict individual drug response: patients with C -dominant disease state should respond better to constraint-reduction drugs; those with Φ -deficit should respond better to flux-enhancement.
3. Fasting duration should correlate with multi-system C reduction independently of caloric restriction, supporting therapeutic fasting protocols in non-metabolic diseases.
4. The AFL treatment algorithm (Step 1–7) applied to CKD anaemia should outperform empirical ESA dosing titration for time-to-target haemoglobin.

Status: Open

10 Falsifiability

The AFL therapeutic framework would be falsified if: cross-axis drug combinations are not more synergistic than same-axis combinations; pre-treatment Ψ_s profile does not predict drug class response; fasting shows no multi-system benefit in well-controlled clinical trials; or the AFL treatment algorithm underperforms empirical clinical titration.

Status: Open

11 Deep Analysis: Derivation of the Universal Saturation Law and Bridge Function Family

Detailed derivation placeholder retained for future expansion in this preprint release.

11.1 Biological Translation of Mujjabi Hysteresis and Alpha-Jitter Tests

To empirically validate AFL in clinical settings, we map the Mujjabi Hysteresis Test and Alpha-Jitter Test to therapeutic outcomes. Hysteresis appears as disease states requiring significantly lower Φ to maintain the pathology than was required to trigger it. The Alpha-Jitter equates to residual structural shifts in C driven by high-frequency flux intensity spikes, measurable via continuous biomarker monitoring.

12 Biological Mujjabi Laws & Discoveries

12.1 The Mujjabi Cycling-Retention Test [Mujjabi \[2026\]](#) (Biology)

Translating the Part II environmental fluctuation theorems, the **Mujjabi Cycling-Retention Test** [Mujjabi \[2026\]](#) proposes that biological resilience depends not only on peak steady-state throughput, but also on retention or clearance of structural memory (M_s) across repeated stress-recovery cycles.

12.1.1 Runtime Diagnostic: Cycling and Recovery

Release-local runtime diagnostics compare cyclic and chronic toy loads. The useful result is qualitative: recovery windows can change retained memory and the operating ratio. The diagnostic does not prove that any fasting, exercise, or treatment schedule is required for health.

13 Translational Medicine: The "AFL Intervention" Algorithm

To translate these model hypotheses into researchable clinical workflows, we outline a therapeutic decision tree based on the proposed operating ratio Ψ_s :

- **State A (Drive Overload):** If $\Psi_s > 1$ due to excessive input Φ (e.g., hypercaloric feeding), the strict intervention axis is *Deprescribing* or *Fasting* (Reduce Φ). Escalating pharmacological clearance forces the system into higher constraint.
- **State B (Constraint Saturation):** If $\Psi_s > 1$ due to high constraint C with normal Φ (e.g., sarcopenia, heart failure), the intervention axis is *Hormetic Pulsing*. Intermittent, controlled spikes in Φ (Exercise) must be applied to stimulate β relaxation and expand the C capacity boundaries.
- **State C (Pharmacological Masking):** A formal warning: Any intervention that suppresses a symptomatic readout of Φ without addressing the underlying M_s structural drift violates the Mujjabi Capacity Law and will inevitably trigger the Flux-Shunting cascade.

14 Clinical Mapping of Symbols

Table 1: AFL symbol map for therapy and intervention systems.

Symbol	Therapeutic meaning	Measurable proxy
Φ	Physiological drive targeted by intervention	Disease-specific function metric (GFR, VO2)
C	Therapeutic constraint / pathway resistance	Biomarker of disease-specific barrier (e.g., fibrosis)
M_s	Structural memory / chronicity	Duration of disease, epigenetic load, tissue memory
Ψ_s	Therapeutic operating ratio	Clinical composite score, response index

15 Known Medicine Baseline

The following guidelines and frameworks take absolute clinical precedence over any AFL therapy taxonomy:

- **National Institute for Health and Care Excellence (NICE) Clinical Guidelines:** Evidence-based treatment guidance for each disease domain are mandatory; AFL intervention-axis classification is a supplementary conceptual layer only.
- **FDA Drug Approval and Labelling:** All pharmacological interventions must follow approved indications and labelling; AFL taxonomy does not constitute prescribing authority.
- **ADA Standards of Medical Care in Diabetes (2026):** Metabolic intervention sequencing follows ADA guidelines; AFL metabolic-flux framing is a hypothesis for future alignment only.
- **KDIGO 2024 CKD Guidelines:** Renal intervention management follows KDIGO; AFL renal-constraint framing is a modelling hypothesis.
- **Surviving Sepsis Campaign (SSC) 2021:** Immune/inflammatory intervention follows SSC bundles; AFL immune-throughput framing is supplementary.

No AFL therapy output should guide drug selection, combination therapy, dietary intervention, or lifestyle plan without guideline-concordant clinical assessment, specialist oversight, and patient-specific safety evaluation.

16 Intervention-Axis Map

The AFL therapy taxonomy maps interventions to the axis they primarily target, without implying that axis-mapping alone is sufficient for prescribing. The four candidate intervention axes are:

1. **Flux enhancement:** Candidate when $\Psi_s < 1$ due to under-delivery (e.g., EPO supplementation for erythropoietic deficit, insulin in absolute deficiency). Requires confirmation that constraint is not the limiting factor.
2. **Constraint reduction:** Candidate when C is the primary driver (e.g., SGLT2i for renal-tubular constraint, anti-inflammatory biologics for immune constraint). Requires assessment of whether flux can safely increase.
3. **Flux limitation:** Candidate when $\Psi_s > 1$ due to drive overload (e.g., caloric restriction, beta-blockade in tachyarrhythmia). Requires ruling out compensatory over-drive.
4. **System stabilisation:** Candidate when Ψ_s oscillates (e.g., pacemaker therapy, mood-stabiliser, anticonvulsant). Requires biomarker evidence of oscillation rather than static displacement.

All four axes are candidate hypotheses. Synergy is predicted when a combination targets different axes; empirical validation is required before clinical use.

17 Experiments and Future Validation

1. **Step 1 — Mathematical consistency:** Verify intervention-axis toy simulations and cycling-recovery diagnostics (complete — `AFL_Biological_Mujjabi_Tests.py`, outputs in `outputs/`).
2. **Step 2 — Synthetic perturbation:** Reproduce known drug synergy curves (e.g., RAAS + SGLT2i in CKD) as AFL cross-axis combinations; test whether same-axis combinations show predicted redundancy.
3. **Step 3 — Retrospective dataset:** Apply AFL axis classification to published combination therapy RCT data; test whether cross-axis combinations show superior outcomes to same-axis combinations across disease domains.
4. **Step 4 — Prospective observational:** Track Ψ_s proxies before and after intervention in a multimorbidity cohort; test whether baseline AFL state predicts treatment response class.
5. **Step 5 — Pilot intervention study:** AFL-guided combination selection versus empirical clinical selection in a randomised pilot; primary endpoint: time to therapeutic target, adverse events, axis-orthogonality score.

6. **Step 6 — Regulatory alignment:** Engage with FDA Project Optimus and NICE health-technology-assessment frameworks to define the evidence standard required for AFL-informed prescribing tools.

Validation ladder status: Step 1 complete; Steps 2–6 open.

18 Clinical Safety Boundary

This paper provides a mathematical taxonomy of intervention axes. It contains no prescribing instructions, drug recommendations, dietary plans, or dosing guidance. The “AFL Intervention Algorithm” sections in this paper describe candidate research hypotheses, not clinical protocols. Specifically:

- **State A (Drive Overload)** suggestions (fasting, deprescribing) are candidate research hypotheses; actual de-escalation decisions require clinical assessment of competing risks and guideline review.
- **State B (Constraint Saturation)** suggestions (hormetic pulsing, exercise) are candidate research hypotheses; actual exercise plans require cardiac, renal, and musculoskeletal safety assessment.
- **State C (Pharmacological Masking)** is a theoretical warning concept; it does not authorise medication discontinuation without specialist oversight.

19 Runtime Diagnostic

The release-local therapy synthesis script is `supplementary/AFL_Biological_Mujjabi_Tests.py`. It simulates intervention-axis comparisons, cycling-recovery diagnostics, and combination-synergy toy scenarios, producing trajectory plots and a `summary.json` in `outputs/`.

All outputs carry the label “toy diagnostic — not a clinical recommendation.”

20 Conclusion

AFL provides a meta-level intervention taxonomy that classifies treatments by their effect on flux–constraint balance rather than by molecular target alone. The framework predicts synergy from axis-orthogonality, explains monotherapy failure in multi-constraint disease, and positions dietary, lifestyle, and pharmacological interventions within a single mathematical language.

Clinical translation requires prospective validation across multiple disease domains, regulatory engagement, and subspecialist review before any AFL intervention-axis output influences prescribing. This paper establishes the taxonomy and generates testable hypotheses; clinical validity remains fully open.

21 Reproducibility Note

Release-local script: `supplementary/AFL_Biological_Mujjabi_Tests.py`. Dependencies: NumPy, matplotlib. Runtime < 2s. All parameters are set in-file. Outputs written to `outputs/`. No proprietary data required.

References

Steve Bico Mujjabi. Cdfd part ii: Origins of life and tri-regime bioenergetics. 2026. doi: 10.5281/zenodo.20264779. URL <https://doi.org/10.5281/zenodo.20264779>.