

Paper F-04: Nuclear Dynamics: A Constraint-Driven Flux Dynamics (CDFD) Perspective

Steve Bico Mujjabi, MD

Independent Researcher

ORCID: 0009-0001-0556-5516

May 2026

Abstract

This paper asks whether Constraint-Driven Flux Dynamics (CDFD) and Adaptive Flux Limitation (AFL) can give the named system a sharper audit language. The working variables are driving flux Φ , constraint C , surface responsiveness S , and structural memory M_s . The useful question is plain: what can be measured, what would count as overload, and what result would make the mapping fail?

1 Series Context

Part I sets the flow–constraint language used here [1]. Part II develops the Life Number and the origins-of-life use of the same notation [2]. Part III applies AFL to biology and medicine [3]. Part IV is the cross-domain stress test: it asks where the notation clarifies measurement, and where it becomes only a metaphor.

2 Introduction

Notation. In Part IV, C names the active constraint or capacity burden in the system being discussed. Φ is the driving flow, S is the surface response, and M_s is retained structural memory. Λ appears only where the Part II Life Number is being referenced [2]. Other Greek symbols are local coefficients whose meaning is set by the equation in which they appear.

2.1 The Mujjabi Universal Capacity Law

Building directly upon the Fundamental Physics (Part I) [1], the **Mujjabi Universal Capacity Law** proposes that across all scales—from black hole event horizons to global financial markets and power grids—driving high flux (Φ) against a rigid topological constraint (C) can drive system saturation. When Φ/C approaches critical thresholds, the system does not scale linearly; it undergoes a topological phase transition.

2.2 The Mujjabi Universal Memory Law

The **Mujjabi Universal Memory Law** frames persistent systemic collapse. When a complex network fails to clear its structural memory (M_s) and its relaxation parameter decays ($\tau_{relax} \rightarrow \infty$), temporary stressors become permanent rigidities. This is the proposed CDFD mechanism behind ecological death, economic depressions, and infrastructural gridlock.

At the femtometer scale, the atomic nucleus is a highly dense, dynamic droplet of nucleons bound together by the residual strong force. The balance of forces within the nucleus dictates the stability of all matter in the universe. While quantum chromodynamics (QCD) describes the fundamental quark-gluon interactions, describing bulk nuclear phenomena (such as fission and alpha decay) requires a macro-scale thermodynamic approach.

CDFD offers a unified thermodynamic ontology. Whether analyzing a living biological cell or an atomic nucleus, the system must maintain sufficient internal throughput to offset its structural constraints. In the nucleus, this throughput is the continuous exchange of virtual mesons (primarily pions) between nucleons.

3 The Nuclear Adaptive Surface

We model the atomic nucleus as an active surface characterized by the CDFD scaling equation:

$$\Psi_s = \left(\frac{\Phi}{C} \right) \cdot S \cdot M_s \quad (1)$$

For a nucleus composed of Z protons and N neutrons, the parameters map as follows:

- **Flux** (Φ): The strong force mediated nucleon-nucleon exchange. This flux scales with the total number of adjacent nucleon pairs, roughly proportional to the nuclear volume $A = Z + N$.
- **Constraint** (C): The disruptive Coulomb repulsion between protons, which scales as $Z^2/A^{1/3}$, along with the Pauli exclusion principle which forces nucleons into higher energy states.
- **Responsiveness** (S): The binding energy per nucleon, indicating the depth of the local potential well and the surface tension of the nuclear droplet.
- **Memory** (M_s): Nuclear shell structure. Nuclei with "magic numbers" of protons or neutrons exhibit closed shells, locking the nuclear topology into a highly efficient, rigid state (perfect memory, $M_s \rightarrow 1.0$).

3.1 Nuclear Decay as Adaptive Failure

A stable nucleus maintains $\Psi_s \approx 1$. The cohesive flux of the strong force perfectly regulates the outward pressure of the Coulomb constraint.

As we move to heavy elements (increasing Z), the Coulomb constraint $C \propto Z^2$ grows much faster than the strong force flux $\Phi \propto A$. Consequently, the ratio (Φ/C) begins to plummet.

To prevent the system from entering a catastrophic decay regime ($\Psi_s \ll 1$), the nucleus must adapt by increasing its neutron-to-proton ratio, effectively boosting Φ without adding to C .

However, eventually, the surface responsiveness S reaches a saturation limit. The nucleus can no longer bind the excess neutrons effectively due to the Pauli constraint. The system memory M_s locks into a state of chronic overload. The nucleus violently sheds the excess constraint through Alpha decay or spontaneous fission, fracturing the system to restore $\Psi_s \approx 1$ in the resulting daughter nuclei.

4 Measurement Layer

The first job in any domain is to choose proxies before drawing conclusions. Φ may be a rate of material, energy, information, capital, or signal flow. C is the bottleneck that slows or redirects that flow. S measures the system's ability to reconfigure under stress, and M_s records the past load that still changes present behavior. A useful application gives those four terms observable definitions, a time scale, and a failure condition. If a standard domain model explains the same data more cleanly, the CDFD/AFL mapping should give way.

4.1 Current Runtime Diagnostic: Universal Network Cascade

The diagnostic script `Part_E_Synthesis/supplementary/run_partiv_discovery.py` keeps this paper tied to the current CDFD Runtime [4], including RK4 field updates, surface dynamics, structural-memory diffusion, and a finite-value audit. In the May 2026 run, the localized hub-drive stress test ended with hub $\Psi_s = 14.8$, peak network $\Psi_s = 14.8$, 21 nodes above $\Psi_s > 1.5$, and 37 maximum memory-locked nodes. I treat those numbers as a runtime sanity check and as a target for later domain measurement, not as a substitute for field data.

5 Conclusion

Radioactive decay is not a random violation of stasis; it is the necessary, model-predicted response of an active system failing to maintain adequate throughput against rising constraints. By framing nuclear dynamics as an Adaptive Flux Limited system, we bridge the gap between subatomic quantum mechanics and universal macroscopic thermodynamics, revealing that the atomic nucleus breathes and regulates its structural topology exactly like an organic cell.

Conflict of Interest

The author declares no conflict of interest.

Data Availability

Release-local runtime diagnostics are available under each Part's `outputs/` folder, with release-wide synthesis artifacts under `Part_E_Synthesis/supplementary/`, `Part_E_Synthesis/outputs/`, and `Part_E_Synthesis/figures/`.

6 Reference Layer

References

- [1] Steve Bico Mujjabi. Cdfd part i: Fundamental physics - constraint-driven field theory and flux dynamics, 2026.
- [2] Steve Bico Mujjabi. Cdfd part ii: Origins of life and tri-regime bioenergetics - constraint-driven flux dynamics, 2026.
- [3] Steve Bico Mujjabi. Cdfd part iii: Adaptive flux limitation biology and medicine - constraint-driven flux dynamics, 2026.
- [4] Steve Bico Mujjabi. Cdfd runtime: Constraint-driven flux dynamics and cdfl execution engine, 2026.