

Paper D-04: Epigenetics: A Constraint-Driven Flux Dynamics (CDFD) Perspective

Steve Bico Mujjabi, MD

Independent Researcher

Founder, Vura Labs

Kampala, Uganda

ORCID: 0009-0001-0556-5516

August 2026

Shared Part IV notation and evidence boundaries are defined in PART_IV_METHODS_AND_CLAIM_BOUNDARY.md. This paper states only its local observables, comparison, and failure conditions.

Abstract

This paper develops a falsifiable CDFD/AFL model of Epigenetics. It maps developmental signals, environmental exposures, hormones, and metabolism to driving flux Φ , chromatin access, transcriptional capacity, repair, and cellular context to active constraint C , remodeling, feedback, compensation, and lineage-specific regulation to responsiveness S , and epigenetic marks, lineage, exposure history, and developmental timing to structural memory M_s . The target outcome is gene expression, phenotype, persistence, and reversibility, tested against multi-omics, lineage tracing, perturbation, exposure cohorts, and longitudinal phenotypes. The paper treats universality as a shared model architecture rather than an assertion that unlike systems have the same material mechanism. It states the negative result that would make the mapping unnecessary and places the domain claim inside the cumulative CDFD programme and its reproducible runtime boundary.

1 Introduction

The genome is not a static library; it is a bustling factory floor. RNA polymerases continuously scan the DNA, transcribing genetic code into the mRNA that builds the organism. However, not all parts of the factory floor are accessible. The cell physically tightly coils segments of DNA around histone proteins, locking them away from the transcriptional machinery.

In CDFD, this physical coiling is the quintessential mechanism of Adaptive Flux Limitation (AFL) at the molecular level.

2 The Genomic Adaptive Surface

We apply the Adaptive Operating Ratio to the epigenetic landscape:

$$\Psi_s = \left(\frac{\Phi}{C} \right) \cdot S \cdot M_s \quad (1)$$

For a specific genetic locus:

- **Flux** (Φ): The rate of transcription, measured by the volume of mRNA produced and the subsequent protein synthesis cascade.
- **Constraint** (C): Chromatin compaction. Heterochromatin represents a high constraint state where genes are silenced; euchromatin represents a low constraint state where genes are freely accessible.
- **Responsiveness** (S): The availability of transcription factors, RNA polymerase, and cellular ATP required to drive the expression process.
- **Memory** (M_s): Epigenetic marks (e.g., DNA methylation and histone acetylation). These biochemical tags act as chronic memory locks. They ensure that a cell "remembers" its constraint topology even after it divides, maintaining cellular identity (e.g., keeping a skin cell a skin cell).

2.1 Trauma, Adaptation, and Transgenerational Inheritance

During normal cellular differentiation, the adaptive ratio Ψ_s is carefully modulated to create distinct tissue types.

However, if the organism experiences severe environmental trauma (e.g., starvation, intense stress, or chronic inflammation), the organism must adapt to survive. The system aggressively alters its epigenetic memory (M_s). It may heavily methylate (lock down) certain metabolic genes to conserve energy, drastically increasing the local constraint C and dropping the transcription flux Φ .

Because these epigenetic marks constitute structural memory (M_s), they are incredibly persistent. The system locks itself into a defensive, highly constrained topology ($\Psi_s \ll 1$). Remarkably, this memory can be passed down to offspring. The progeny inherit the chronic constraints, leaving them predisposed to metabolic disorders (like obesity or anxiety) even in safe, resource-rich environments, because their genetic surface is pathologically over-constrained by their ancestral memory.

3 Measurement Layer

4 Cross-Domain Model Upgrade: Mechanism, Scale, and Tests

4.1 Observable Translation

The paper-specific translation begins with an observable chain rather than a verbal analogy. For Epigenetics, the proposed drive is developmental signals, environmental exposures, hormones,

and metabolism. The active limitation is chromatin access, transcriptional capacity, repair, and cellular context. Responsiveness is carried by remodeling, feedback, compensation, and lineage-specific regulation, while retained state is represented by epigenetic marks, lineage, exposure history, and developmental timing. The outcome to explain is gene expression, phenotype, persistence, and reversibility.

Table 1: Operational CDFD/AFL map for Paper D-04.

Term	Paper-specific observable meaning	Measurement question
Φ	developmental signals, environmental exposures, hormones, and metabolism	What rate, load, gradient, or demand enters the focal system?
C	chromatin access, transcriptional capacity, repair, and cellular context	Which measured bottleneck limits transfer, function, or recovery?
S	remodeling, feedback, compensation, and lineage-specific regulation	Which process changes effective capacity or reroutes the load?
M_s	epigenetic marks, lineage, exposure history, and developmental timing	Which retained state makes equal present loads produce different futures?
Y	gene expression, phenotype, persistence, and reversibility	Which outcome is predicted out of sample, with uncertainty?




Figure 1: Paper D-04 observable CDFD/AFL architecture for Epigenetics. Solid arrows give the proposed forward chain; dashed arrows make history-dependent feedback explicit.

4.2 Paper-Specific Causal Hypothesis

For this paper, the hypothesized causal sequence is specific. A change in developmental signals, environmental exposures, hormones, and metabolism arrives faster than remodeling, feedback, compensation, and lineage-specific regulation can compensate. This raises or redistributes chromatin access, transcriptional capacity, repair, and cellular context. If the load persists, the retained state encoded by epigenetic marks, lineage, exposure history, and developmental timing changes the next response, so a later event of equal magnitude can produce a different trajectory in gene expression, phenotype, persistence, and reversibility. The scientific task is to estimate that sequence against the best ordinary model of the domain, not merely to redescribe the outcome after it occurs.

4.3 Cross-Scale Translation Without Mechanistic Erasure

For the domain applications family, the cross-domain translation is a disciplined test across systems with very different material mechanisms. A valid mapping preserves the baseline science of the domain, declares the scale of every variable, and earns its place through a discriminating prediction rather than resemblance alone. [2, 4, 3, 1] The required scale declaration for this paper is minutes to generations; loci to organisms. At short scales, the key question is whether the incoming drive is transmitted, buffered, or diverted. At intermediate scales, response changes effective capacity and network exposure. At long scales, retained structure can alter the state space itself. A result is genuinely cross-scale only when the aggregation rule is stated and the same conclusion is not an artifact of averaging.

4.4 Evidence Ladder and Discriminating Tests

Table 2: Evidence ladder for Paper D-04. Each rung can fail independently.

Test	Design	Discriminating result
Proxy audit	Measure multi-omics, lineage tracing, perturbation, exposure cohorts, and longitudinal phenotypes and preregister how each record maps to Φ , C , S , M_s , and Y .	The mapping fails if proxies are non-identifiable, circular, or change meaning across cases.
Load ramp	Apply or observe a controlled exposure followed by removal or reprogramming while estimating the dominant constraint and response time.	A threshold claim requires a reproducible nonlinear change and uncertainty bounds, not a visually chosen breakpoint.
Recovery and memory	Match present load across units with different histories, then compare recovery paths.	The memory claim requires history-dependent divergence after current conditions are controlled.
Model comparison	Compare the baseline domain model with and without the CDFD/AFL state variables using held-out data.	The paper’s stated falsifier is: past exposure does not alter response after current molecular state is controlled.

4.5 Research Programme and Boundary Conditions

The immediate research programme has four steps. First, construct a documented dataset from multi-omics, lineage tracing, perturbation, exposure cohorts, and longitudinal phenotypes and retain raw units before normalization. Second, fit a baseline domain model and report its calibration and out-of-sample error. Third, add the smallest identifiable CDFD/AFL state extension and test whether it improves prediction of gene expression, phenotype, persistence, and reversibility. Fourth, repeat the analysis across the declared scale range and publish negative cases as part of the cross-domain translation audit.

5 Conclusion

Epigenetic state is a concrete molecular form of biological memory whose mechanisms must be studied on their own terms. CDFD supplies a high-level language for testing history-dependent

constraint and response, but it does not erase the mechanistic gap between chromatin regulation, trauma, and economic systems.

Conflict of Interest

The author declares no conflict of interest.

Data Availability

Part-specific runtime diagnostics are under each Part's `outputs/` directory.

Release-wide code: `Part_E_Synthesis/supplementary/`.

Generated summaries: `Part_E_Synthesis/outputs/`.

Figures: `Part_E_Synthesis/figures/`.

6 Reference Layer

References

- [1] Per Bak, Chao Tang, and Kurt Wiesenfeld. Self-organized criticality: An explanation of the $1/f$ noise. *Physical Review Letters*, 59(4):381–384, 1987. DOI: <https://doi.org/10.1103/PhysRevLett.59.381>.
- [2] C. S. Holling. Resilience and stability of ecological systems. *Annual Review of Ecology and Systematics*, 4:1–23, 1973. DOI: <https://doi.org/10.1146/annurev.es.04.110173.000245>.
- [3] Claude E. Shannon. A mathematical theory of communication. *Bell System Technical Journal*, 27(3):379–423, 1948. DOI: <https://doi.org/10.1002/j.1538-7305.1948.tb01338.x>.
- [4] A. M. Turing. The chemical basis of morphogenesis. *Philosophical Transactions of the Royal Society of London. Series B*, 237(641):37–72, 1952. DOI: <https://doi.org/10.1098/rstb.1952.0012>.