

Fractal Dimension: Theory and Computation

Box-Counting Estimation via Hilbert Curve Analysis

ME

May 28, 2026

Contents

1	Introduction	2
2	Measure Theory	2
3	Hausdorff Measure	4
3.1	Definition	4
3.2	Phase Transition	4
4	Hausdorff Dimension	5
4.1	Basic Properties	5
5	Box-counting Dimension	5
5.1	Definition	5
5.2	Relationship to Hausdorff dimension	6

1 Introduction

This entire paper is basically based on [\[Fal03\]](#).

Perhaps not a very serious paper since its just what I found when I wanted to look into this area.

2 Measure Theory

The material in this section is standard and follows the presentation in [\[Men26\]](#).

First to define measure theory (I didn't take Lebesgue so I will be using the above notes...)

A measure is a way to assign a "size" to subsets of a set X . Let $\mathcal{F}$ be a σ -algebra on X .

Definition 2.1. (Measure). Let $(X, \mathcal{F})$ be a measurable space. A measure on $(X, \mathcal{F})$ is a function $\mu : \mathcal{F} \rightarrow [0, \infty]$ satisfying:

1. $\mu(\emptyset) = 0$
2. **Countable additivity:** if $A_1, A_2, \dots \in \mathcal{F}$ are pairwise disjoint, then

$$\mu\left(\bigsqcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mu(A_i)$$

The triple $(X, \mathcal{F}, \mu)$ is called a **measure space**.

Proposition 2.2 (Basic properties). *Let $(X, \mathcal{F}, \mu)$ be a measure space.*

1. **Monotonicity:**

$$A \subseteq B \Rightarrow \mu(A) \leq \mu(B)$$

2. **Countable subadditivity:**

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) \leq \sum_{i=1}^{\infty} \mu(A_i)$$

3. **Continuity from below:** if $A_1 \subseteq A_2 \subseteq \dots$ then

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) = \lim_{i \rightarrow \infty} \mu(A_i)$$

4. **Continuity from above:** if $A_1 \supseteq A_2 \supseteq \dots$ and $\mu(A_1) < \infty$ then

$$\mu\left(\bigcap_{i=1}^{\infty} A_i\right) = \lim_{i \rightarrow \infty} \mu(A_i)$$

The canonical example is the **Lebesgue measure** λ^n on $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$, where $\mathcal{B}(\mathbb{R}^n)$ is the Borel σ -algebra — the smallest σ -algebra containing all open sets. Lebesgue measure assigns to each box $[a_1, b_1] \times \dots \times [a_n, b_n]$ its volume $(b_1 - a_1) \dots (b_n - a_n)$, and extends uniquely to all Borel sets. It formalises the intuitive notions of length, area, and volume.

The inadequacy of Lebesgue measure for fractal geometry is immediate: a curve $\gamma \subset \mathbb{R}^2$ has $\lambda^2(\gamma) = 0$ (zero area) but $\lambda^1(\gamma)$ may be infinite (infinite length). Neither gives useful information about the “size” of γ . What is needed is a family of measures parametrised by a continuous exponent $s \geq 0$.

To construct such a family, we need the notion of an outer measure, which is slightly weaker than a measure — it is defined on *all* subsets of X , not just measurable ones, and only requires subadditivity rather than full additivity.

Definition 2.3 (Outer measure). A function $\mu^* : 2^X \rightarrow [0, \infty]$ is an **outer measure** on X if:

1. $\mu^*(\emptyset) = 0$
2. **Monotonicity:** $A \subseteq B \Rightarrow \mu^*(A) \leq \mu^*(B)$
3. **Countable subadditivity:**

$$\mu^*\left(\bigcup_{i=1}^{\infty} A_i\right) \leq \sum_{i=1}^{\infty} \mu^*(A_i)$$

The relationship between outer measures and measures is given by the Hahn-Carathéodory extension theorem: given an outer measure μ^* on X , the collection

$$\Sigma = \{A \subseteq X : \mu^*(B) = \mu^*(B \cap A) + \mu^*(B \setminus A) \text{ for all } B \subseteq X\}$$

is a σ -algebra, and $\mu^*|_{\Sigma}$ is a measure. Sets in Σ are called **μ^* -measurable**. In particular, all Borel sets are μ^* -measurable whenever μ^* is constructed from a Borel-compatible pre-measure — which will be the case for Hausdorff measure.

The standard way to construct an outer measure is via coverings. If $\mathcal{K} \subseteq 2^X$ covers X (with $\emptyset \in \mathcal{K}$) and $\tilde{\mu} : \mathcal{K} \rightarrow [0, \infty]$ assigns sizes to sets in $\mathcal{K}$ with $\tilde{\mu}(\emptyset) = 0$, then

$$\mu^*(A) = \inf \left\{ \sum_{i=1}^{\infty} \tilde{\mu}(K_i) \mid K_i \in \mathcal{K}, A \subseteq \bigcup_{i=1}^{\infty} K_i \right\}$$

defines an outer measure on X . This is precisely the construction used to define Hausdorff measure — with $\mathcal{K}$ being all subsets of diameter at most δ , and $\tilde{\mu}(U) = |U|^s$.

3 Hausdorff Measure

The material in sections 3 and 4 follows [Fal03].

3.1 Definition

Let (X, d) denote a metric space, $S \subseteq X$ a Borel set.

For a set $U \subseteq X$, write $|U| := \text{diam}(U) = \sup_{x,y \in U} d(x, y)$

Definition 3.1. Let $s \geq 0$, and $\delta > 0$. Define

$$\mathcal{H}_\delta^s(S) = \inf \left\{ \sum_{i=1}^{\infty} |U_i|^s \mid S \subseteq \bigcup_{i=1}^{\infty} U_i, |U_i| \leq \delta \right\}$$

This is the infimum over all countable covers of S by sets of diameter at most δ .

As $\delta \rightarrow 0$, we allow fewer and fewer covers, so $\mathcal{H}_\delta^s(S)$ is non-decreasing in $1/\delta$. The limit

$$\mathcal{H}^s(S) = \lim_{\delta \rightarrow 0} \mathcal{H}_\delta^s(S) \in [0, \infty]$$

is the s -dimensional Hausdorff measure of S . This is possibly $+\infty$.

$\mathcal{H}^s$ is an outer measure on X , and its restriction to the Borel σ -algebra is a measure.

3.2 Phase Transition

Observe that $\mathcal{H}^s(S)$ exhibits a phase transition in s :

Theorem 3.2. *If $\mathcal{H}^s(S) < \infty$ for some s , then $\mathcal{H}^t(S) = 0$ for all $t > s$.*

Proof. Suppose $\mathcal{H}^s(S) < \infty$. For any cover $\{U_i\}$ with $|U_i| \leq \delta$. For $t > s$:

$$\sum |U_i|^t = \sum |U_i|^{t-s} \cdot |U_i|^s \leq \delta^{t-s} \sum |U_i|^s$$

Taking the infimum over all covers gives

$$\mathcal{H}_\delta^t(S) \leq \delta^{t-s} \cdot \mathcal{H}_\delta^s(S) \leq \delta^{t-s} \cdot \mathcal{H}^s(S)$$

where the second inequality holds since $\mathcal{H}_\delta^s(S)$ is non-decreasing in $1/\delta$.

Now let $\delta \rightarrow 0$, $t - s > 0$, and $\mathcal{H}^s(S) < \infty$ is a fixed finite constant. So the right hand side goes to 0, hence $\mathcal{H}^t(S) = 0$. $\square$

Corollary 3.3. *There exists a unique value $d \in [0, \infty]$ such that*

$$\mathcal{H}^s(S) = \begin{cases} \infty & s < d \\ 0 & s > d \end{cases}$$

At $s = d$, the measure $\mathcal{H}^d(S)$ may take any value in $[0, +\infty]$.

4 Hausdorff Dimension

Definition 4.1. (Hausdorff Dimension). The **Hausdorff Dimension** of S is the critical exponent

$$\dim_{\mathcal{H}}(S) = \inf\{s \geq 0 : \mathcal{H}^s(S) = 0\} = \sup\{s \geq 0 : \mathcal{H}^s(S) = +\infty\}$$

This is the “true” dimension of a fractal.

4.1 Basic Properties

Proposition 4.2 (Basic properties). *Hausdorff dimension satisfies the following:*

1. **Monotonicity:** $A \subseteq B \implies \dim_{\mathcal{H}}(A) \leq \dim_{\mathcal{H}}(B)$
2. **Countable stability:** $\dim_{\mathcal{H}}(\cup_{i=1}^{\infty} A_i) = \sup_i \dim_{\mathcal{H}}(A_i)$
3. **Countable sets:** if S is countable, $\dim_{\mathcal{H}}(S) = 0$.
4. **Smooth manifolds**¹: if $M \subset \mathbb{R}^n$ is a smooth k -dimensional submanifold, then $\dim_{\mathcal{H}}(M) = k$.

The following proposition is a useful tool we’ll need later down the line:

Proposition 4.3. *Lipschitz invariance* Let $f : A \rightarrow B$ be Lipschitz with constant C . Then $\dim_{\mathcal{H}}(f(A)) \leq \dim_{\mathcal{H}}(A)$. In particular, bi-Lipschitz maps preserve Hausdorff dimensions.

Proof. If $\{U_i\}$ covers A with $\sum |U_i|^s < \infty$, then $\{f(U_i)\}$ covers $f(A)$ with $|f(U_i)| \leq C|U_i|$ (by Lipschitz), so $\sum |f(U_i)|^s \leq C^s \sum |U_i|^s < \infty$. Hence $\mathcal{H}^s(f(A)) \leq C^s \mathcal{H}^s(A)$, and the dimension inequality follows. $\square$

5 Box-counting Dimension

5.1 Definition

A.K.A Minkowski–Bouligand dimension.

We need a way to be able to effectively compute the dimension of a fractal (i.e. the Hausdorff dimension). A perhaps easy way to do this is to cover said shape with boxes of size ε and count them. We check to see how the number of boxes hitting the shape $N(\varepsilon)$ changes w.r.t the length ε . We use the following formula:

$$\dim_{\text{box}}(S) := \lim_{\varepsilon \rightarrow 0} \frac{\log N(\varepsilon)}{\log(1/\varepsilon)} = - \lim_{\varepsilon \rightarrow 0} \frac{\log N(\varepsilon)}{\log(\varepsilon)}$$

This is roughly the dimension d such that $N(\varepsilon) \approx C\varepsilon^{-d}$.

Intuitively, we choose this to represent the dimension because a d -dimensional box of side

¹A smooth k -dimensional manifold is just a set that locally looks like $\mathbb{R}^k$

ε has a “ d -dimensional volume” ε^d . So we end up with some $N(\varepsilon) \approx \frac{V}{\varepsilon^d} = C\varepsilon^{-d}$. We want to find a d for a fractal i.e. finding the d such that the same scaling property holds.

5.2 Relationship to Hausdorff dimension

Proposition 5.1. *For any bounded $S \subseteq \mathbb{R}^n$, $\dim_{\mathcal{H}}(S) \leq \dim_{\text{box}}(S)$.*

Proof. Let $d > \dim_{\text{box}}(S)$. Then $N(\varepsilon) \leq C\varepsilon^{-\dim_{\text{box}}(S)}$ for some constant C and all small ε . Covering S with $N(\varepsilon)$ hypercubes of diameter $\varepsilon\sqrt{n}$:

$$\mathcal{H}_{\varepsilon\sqrt{n}}^d(S) \leq N(\varepsilon) \cdot (\varepsilon\sqrt{n})^d \leq C(\sqrt{n})^d \cdot \varepsilon^{d-\dim_{\text{box}}(S)} \rightarrow 0$$

as $\varepsilon \rightarrow 0$. Hence $\mathcal{H}^d(S) = 0$, so $\dim_{\mathcal{H}}(S) \leq d$. Since $d > \dim_{\text{box}}(S)$ was arbitrary, $\dim_{\mathcal{H}}(S) \leq \dim_{\text{box}}(S)$. $\square$

This is a very useful property as will be shown in the next few chapters. With this we now have a best upper bound for calculating the Hausdorff dimension. The equality actually holds on “well-behaved” fractals which is exactly what we need.

References

- [Fal03] Kenneth Falconer. *Fractal Geometry: Mathematical Foundations and Applications*. 2nd. Wiley, 2003.
- [Men26] Angeliki Menegaki. *Measure Theory and Integration: Lecture Notes MATH50006*. 2026.