

# Symmetry of Magnetically Induced Currents

MAGIC 2022 | *Peterhouse, Cambridge, UK*

Bang C. Huynh<sup>†</sup>, Andrew M. Teale<sup>†</sup>

<sup>†</sup>School of Chemistry, University of Nottingham, UK

13<sup>th</sup> September 2022



University of  
Nottingham  
UK | CHINA | MALAYSIA



**MAGIC 2022**  
4<sup>th</sup> workshop on MAGnetically  
Induced Currents in molecules

# Introduction

# Overview

- ⚙ Symmetry and pseudo-symmetry *groups* in magnetic fields
- ⚙ *Unitary* representation analysis on *linear spaces*
- ⚙ *Wavefunction* and *current density* symmetries
  - ⚙ *Relationships* between wavefunction and current density symmetries
  - ⚙ Symmetry *descent* and symmetry *breaking* in magnetic fields

# Groups in magnetic fields

# The electronic Hamiltonian

- ⚙ For an  $N_e$ -electron system in a **uniform** magnetic field  $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$ , consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\hat{\mathcal{H}} = \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k$$

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

# The electronic Hamiltonian

- ⚙ For an  $N_e$ -electron system in a **uniform** magnetic field  $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$ , consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\hat{\mathcal{H}} = \boxed{\frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2} + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k$$

↑  
kinetic

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

# The electronic Hamiltonian

- ⚙ For an  $N_e$ -electron system in a **uniform** magnetic field  $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$ , consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\hat{\mathcal{H}} = \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \underbrace{\sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k)}_{\text{external potential}} + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k$$

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

# The electronic Hamiltonian

- ⚙ For an  $N_e$ -electron system in a **uniform** magnetic field  $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$ , consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\hat{\mathcal{H}} = \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \underbrace{\frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|}}_{\text{electron-electron interaction}} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k$$

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).



# The electronic Hamiltonian

- ⚙ For an  $N_e$ -electron system in a **uniform** magnetic field  $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$ , consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\hat{\mathcal{H}} = \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \boxed{\frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k}$$

↑  
spin Zeeman interaction

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

# The electronic Hamiltonian

- ⚙ For an  $N_e$ -electron system in a **uniform** magnetic field  $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$ , consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\begin{aligned}\hat{\mathcal{H}} &= \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k \\ &= \frac{1}{2} \sum_{k=1}^{N_e} \hat{p}_k^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k)\end{aligned}$$

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

# The electronic Hamiltonian

- ⚙ For an  $N_e$ -electron system in a **uniform** magnetic field  $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$ , consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\begin{aligned}\hat{\mathcal{H}} &= \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k \\ &= \underbrace{\frac{1}{2} \sum_{k=1}^{N_e} \hat{\mathbf{p}}_k^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|}}_{\text{Zero-field Hamiltonian, } \hat{\mathcal{H}}_0(v_{\text{ext}})} + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k)\end{aligned}$$

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

# The electronic Hamiltonian

- ⚙ For an  $N_e$ -electron system in a **uniform** magnetic field  $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$ , consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\begin{aligned}\hat{\mathcal{H}} &= \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k \\ &= \frac{1}{2} \sum_{k=1}^{N_e} \hat{p}_k^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \underbrace{\mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k}_{\text{Linear}} + \underbrace{\frac{1}{2} A^2(\mathbf{r}_k)}_{\text{Quadratic}}\end{aligned}$$

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

# The electronic Hamiltonian

- ⚙ For an  $N_e$ -electron system in a **uniform** magnetic field  $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$ , consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\begin{aligned}\hat{\mathcal{H}} &= \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k \\ &= \frac{1}{2} \sum_{k=1}^{N_e} \hat{p}_k^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k) \\ &= \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).\end{aligned}$$

- ⚙ How does each term transform **spatially** and **temporally**?

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. et al. **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. et al. **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

# Tensor transformations

⚙ Let  $\mathbf{v}$  be a rank- $k$  Cartesian tensor in three dimensions.

---

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al. Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

# Tensor transformations

- Let  $\mathbf{v}$  be a rank- $k$  Cartesian tensor in three dimensions.
- Spatially*, let  $u \in O(3)$  be any **proper** or **improper rotation** that acts on an orthogonal basis  $\{\mathbf{e}_i\}$  spanning  $\mathbb{R}^3$  according to

$$\hat{u}\mathbf{e}_i = \mathbf{e}_j U_{ji}.$$

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al.* *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

# Tensor transformations

- Let  $\mathbf{v}$  be a rank- $k$  Cartesian tensor in three dimensions.
- Spatially*, let  $u \in O(3)$  be any **proper** or **improper rotation** that acts on an orthogonal basis  $\{\mathbf{e}_i\}$  spanning  $\mathbb{R}^3$  according to

$$\hat{u}\mathbf{e}_i = \mathbf{e}_j U_{ji}$$

**$U$** : representation matrix for  $u$  in  $\{\mathbf{e}_i\}$

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al. Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).



# Tensor transformations

- Let  $\mathbf{v}$  be a rank- $k$  Cartesian tensor in three dimensions.
- Spatially*, let  $u \in O(3)$  be any **proper** or **improper rotation** that acts on an orthogonal basis  $\{\mathbf{e}_i\}$  spanning  $\mathbb{R}^3$  according to

$$\hat{u}\mathbf{e}_i = \mathbf{e}_j U_{ji}.$$

To consider the **linear** action of  $u$  on  $\mathbf{v}$ , let  $\mathbf{v}' = \hat{u}\mathbf{v}$ .

- $\mathbf{v}$  is a **polar** tensor if

$$v'_{ab\dots k} = U_{ap}U_{bq}\dots U_{kz} v_{pq\dots z}.$$

- $\mathbf{v}$  is an **axial** tensor if

$$v'_{ab\dots k} = |\mathbf{U}| U_{ap}U_{bq}\dots U_{kz} v_{pq\dots z}.$$

Birss, R. R. **Symmetry and Magnetism**. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. **Rev. Mod. Phys.** **40**, 359–379 (April 1968).

Lazzeretti, P. *et al.* **Nucl. Magn. Shield. Mol. Struct.** (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

# Tensor transformations

- Let  $\mathbf{v}$  be a rank- $k$  Cartesian tensor in three dimensions.
- Spatially*, let  $u \in O(3)$  be any **proper** or **improper rotation** that acts on an orthogonal basis  $\{\mathbf{e}_i\}$  spanning  $\mathbb{R}^3$  according to

$$\hat{u}\mathbf{e}_i = \mathbf{e}_j U_{ji}.$$

To consider the *linear* action of  $u$  on  $\mathbf{v}$ , let  $\mathbf{v}' = \hat{u}\mathbf{v}$ .

- $\mathbf{v}$  is a **polar** tensor if

$$v'_{ab\dots k} = U_{ap} U_{bq} \dots U_{kz} v_{pq\dots z}.$$

- $\mathbf{v}$  is an **axial** tensor if

$$v'_{ab\dots k} = |\mathbf{u}| U_{ap} U_{bq} \dots U_{kz} v_{pq\dots z}.$$

+1 for proper rotations, -1 for improper rotations

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al.* *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

# Tensor transformations

- ⚙ Let  $\mathbf{v}$  be a rank- $k$  Cartesian tensor in three dimensions.
- ⚙ *Temporally*, let  $\theta$  be the **time reversal** that acts on an orthogonal basis  $\{\zeta_1, \zeta_2\}$  of a two-component spinor according to

$$\hat{\theta} \begin{bmatrix} \zeta_1 & \zeta_2 \end{bmatrix} = \begin{bmatrix} \zeta_2^* & -\zeta_1^* \end{bmatrix}.$$

---

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al. Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

# Tensor transformations

- Let  $\mathbf{v}$  be a rank- $k$  Cartesian tensor in three dimensions.
- Temporally*, let  $\theta$  be the **time reversal** that acts on an orthogonal basis  $\{\zeta_1, \zeta_2\}$  of a two-component spinor according to

$$\hat{\theta}[\zeta_1 \quad \zeta_2] = [\zeta_2^* \quad -\zeta_1^*].$$

To consider the *antilinear* action of  $\theta$  on  $\mathbf{v}$ , let  $\mathbf{v}'' = \hat{\theta}\mathbf{v}$ .

- $\mathbf{v}$  is a **time-even** tensor (*i*-tensor) if

$$\mathbf{v}'' = \mathbf{v}.$$

- $\mathbf{v}$  is a **time-odd** tensor (*c*-tensor) if

$$\mathbf{v}'' = -\mathbf{v}.$$

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

# Tensor classifications

⚙ There are *four* types of tensors under spatial-temporal transformations.

⚙ **polar time-even** tensors

⚙ position vectors  $\mathbf{r}$

⚙ **axial time-even** tensors

⚙ **polar time-odd** tensors

⚙ linear momentum vectors  $\mathbf{p}$

⚙ magnetic vector potentials  $\mathbf{A}$

⚙ current densities  $\mathbf{j}$

⚙ **axial time-odd** tensors

⚙ angular momentum vectors  $\mathbf{l}$

⚙ magnetic field vectors  $\mathbf{B}$

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al.* *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

# Symmetry and pseudo-symmetry groups

## Definition (symmetry group)

All transformations  $\hat{T}$  that leave the electronic Hamiltonian  $\hat{\mathcal{H}}$  invariant, i.e.  $\hat{T}\hat{\mathcal{H}}\hat{T}^{-1} = \hat{\mathcal{H}}$ , form the **symmetry group** of  $\hat{\mathcal{H}}$ . Such transformations are called **symmetry transformations**.

- ⚙ Symmetry transformations impose **constraints** on the eigenfunctions of  $\hat{\mathcal{H}}$  and properties calculated from them.

# Symmetry and pseudo-symmetry groups

## Definition (symmetry group)

All transformations  $\hat{T}$  that leave the electronic Hamiltonian  $\hat{\mathcal{H}}$  invariant, i.e.  $\hat{T}\hat{\mathcal{H}}\hat{T}^{-1} = \hat{\mathcal{H}}$ , form the **symmetry group** of  $\hat{\mathcal{H}}$ . Such transformations are called **symmetry transformations**.

- ⚙ Symmetry transformations impose **constraints** on the eigenfunctions of  $\hat{\mathcal{H}}$  and properties calculated from them.

## Definition (pseudo-symmetry group)

Consider a term  $\hat{\mathcal{H}}' \subset \hat{\mathcal{H}}$ . All transformations  $\hat{T}$  that leave  $\hat{\mathcal{H}}'$  invariant but not the full  $\hat{\mathcal{H}}$  form a **pseudo-symmetry group** of  $\hat{\mathcal{H}}$ . Such transformations are called **pseudo-symmetry transformations**.

- ⚙ Pseudo-symmetry transformations provide ways to understand eigenfunctions of  $\hat{\mathcal{H}}$  (**complicated**) from the perspective of  $\hat{\mathcal{H}}'$  (**simpler**).

# Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$

Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).



# Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \underbrace{\hat{\mathcal{H}}_0(v_{\text{ext}})}_{\text{polar time-even}} + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$

---

Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).  
 Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).  
 Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).  
 Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

# Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \underbrace{\hat{\mathcal{H}}_0(v_{\text{ext}})}_{\mathcal{G} + \theta\mathcal{G}} + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$



Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

# Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \underbrace{\mathbf{A}(\mathbf{r}_k)}_{\substack{\text{polar} \\ \text{time-odd}}} \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$



Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

# Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$

↑  
polar time-odd



Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

# Groups in magnetic fields

⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \underbrace{\mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k}_{\text{O}(3) + \theta\text{O}(3)} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \underbrace{\frac{1}{2}A^2(\mathbf{r}_k)}_{\text{O}(3) + \theta\text{O}(3)}.$$



Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

# Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \underbrace{\frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k}_{\text{axial time-odd}} + \frac{1}{2} A^2(\mathbf{r}_k).$$



Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).


Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

# Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \underbrace{\frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k}_{\substack{\mathcal{H} + \theta u \mathcal{H} \\ (u \in \mathcal{G}, \hat{u} \mathbf{B} = -\mathbf{B})}} + \frac{1}{2} A^2(\mathbf{r}_k).$$



Wigner, E. **Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra**. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. **Rev. Mod. Phys.** **40**, 359–379 (April 1968).


Lazzeretti, P. et al. **Nucl. Magn. Shield. Mol. Struct.** (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. **J. Chem. Phys.** **99**, 3669–3682 (1993).

# Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$



Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).




# Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$

- ⚙ If  $\mathcal{M} = \mathcal{H} + \theta u\mathcal{H}$  is isomorphic to a **unitary group**  $\mathcal{M}'$ , we write  $\mathcal{M} = \mathcal{M}'(\mathcal{H})$ .



Wigner, E. **Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra**. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. **Nucl. Magn. Shield. Mol. Struct.** (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

# Unitary representation analysis

# Group unitary representations on linear spaces

- ⚙ Consider a **group**  $\mathcal{G}$  that acts **unitarily** on a **linear space**  $V$ .
- ⚙ Let  $\mathbf{v}$  be an element in  $V$ . The unitary action of  $\mathcal{G}$  on  $\mathbf{v}$  generates an **orbit**

$$\mathcal{G} \cdot \mathbf{v} = \{\hat{u}_i \mathbf{v} \mid u_i \in \mathcal{G}\}$$

which spans a **representation subspace**  $\Gamma \subseteq V$ .

For simplicity, we will assume that  $\mathcal{G} \cdot \mathbf{v}$  is a linearly independent basis.

# Group unitary representations on linear spaces

- ⚙ Consider a **group**  $\mathcal{G}$  that acts **unitarily** on a **linear space**  $V$ .
- ⚙ Let  $\mathbf{v}$  be an element in  $V$ . The unitary action of  $\mathcal{G}$  on  $\mathbf{v}$  generates an **orbit**

$$\mathcal{G} \cdot \mathbf{v} = \{\hat{u}_i \mathbf{v} \mid u_i \in \mathcal{G}\}$$

which spans a **representation subspace**  $\Gamma \subseteq V$ .

For simplicity, we will assume that  $\mathcal{G} \cdot \mathbf{v}$  is a linearly independent basis.

- ⚙ We seek to decompose  $\Gamma$  into known **irreducible** representations of  $\mathcal{G}$  on  $V$ .  
 $\hookrightarrow$  This **quantifies** the **symmetry** of  $\mathbf{v}$  under the action of  $\mathcal{G}$ .
- ⚙ To this end, we pick a reference element  $\mathbf{v}_i = \hat{u}_i \mathbf{v}$  in  $\mathcal{G} \cdot \mathbf{v}$  and define the **representation matrices**  $D^\Gamma(u_k)$  for all  $u_k \in \mathcal{G}$ :

$$\hat{u}_k \mathbf{v}_i = \sum_j \mathbf{v}_j D_{ji}^\Gamma(u_k).$$

Their traces,  $\chi^\Gamma(u_k)$ , give the **characters** required for the decomposition of  $\Gamma$ .

# Representation matrix determination

- ⚙ We opt for a projection method.
  - ↔ This requires  $V$  to be *endowed* with an *inner product*  $\langle \cdot | \cdot \rangle$ .

# Representation matrix determination

- ⚙ We opt for a projection method.  
 ↪ This requires  $V$  to be *endowed* with an **inner product**  $\langle \cdot | \cdot \rangle$ .
- ⚙ Define a non-orthogonal projection operator

$$\hat{\mathcal{P}}_I = \sum_j |\mathbf{v}_i\rangle (\mathbf{S}^{-1})_{ij} \langle \mathbf{v}_j|$$

where  $S_{ij} = \langle \mathbf{v}_i | \mathbf{v}_j \rangle$  such that  $\hat{\mathcal{P}}_I |\mathbf{v}_j\rangle = \delta_{ij} |\mathbf{v}_i\rangle$ .

# Representation matrix determination

- ⚙ We opt for a projection method.  
 ↪ This requires  $V$  to be **endowed** with an **inner product**  $\langle \cdot | \cdot \rangle$ .
- ⚙ Define a non-orthogonal projection operator

$$\hat{\mathcal{P}}_l = \sum_j |\mathbf{v}_i\rangle (\mathbf{S}^{-1})_{ij} \langle \mathbf{v}_j|$$

where  $S_{ij} = \langle \mathbf{v}_i | \mathbf{v}_j \rangle$  such that  $\hat{\mathcal{P}}_l |\mathbf{v}_j\rangle = \delta_{ij} |\mathbf{v}_i\rangle$ .

- ⚙ Application of  $\hat{\mathcal{P}}$  on the defining equation for  $\mathbf{D}^\Gamma(u_k)$  yields

$$\mathbf{D}^\Gamma(u_k) = \mathbf{S}^{-1} \mathbf{T}(u_k),$$

where

$$T_{ij}(u_k) = \langle \mathbf{v}_i | \hat{u}_k \mathbf{v}_j \rangle.$$

# Representation matrix determination

- ⚙ We opt for a projection method.  
 ↪ This requires  $V$  to be **endowed** with an **inner product**  $\langle \cdot | \cdot \rangle$ .
- ⚙ Define a non-orthogonal projection operator

$$\hat{\mathcal{P}}_i = \sum_j |\mathbf{v}_i\rangle (\mathbf{S}^{-1})_{ij} \langle \mathbf{v}_j|$$

where  $S_{ij} = \langle \mathbf{v}_i | \mathbf{v}_j \rangle$  such that  $\hat{\mathcal{P}}_i |\mathbf{v}_j\rangle = \delta_{ij} |\mathbf{v}_i\rangle$ .

- ⚙ Application of  $\hat{\mathcal{P}}$  on the defining equation for  $\mathbf{D}^\Gamma(u_k)$  yields

$$\mathbf{D}^\Gamma(u_k) = \mathbf{S}^{-1} \mathbf{T}(u_k),$$

where

$$|\mathcal{G}|^3 \text{ elements} \rightarrow T_{ij}(u_k) = \langle \mathbf{v}_i | \hat{u}_k \mathbf{v}_j \rangle.$$

- ⚙ **Closure** of  $\mathcal{G} \implies T_{ij}(u_k)$  can be **mapped to**  $T_{m1}(\mathbf{e})$  for some  $m = 1, \dots, |\mathcal{G}|$ .  
 ↪  $\mathbf{T}(u_k)$  can be computed with  $\mathcal{O}(|\mathcal{G}|)$  time complexity.



# Current density linear space

- ⚙ The current densities  $\mathbf{j}(\mathbf{r})$  with  $\mathbf{r} \in \mathbb{R}^3$  form a linear space  $V_j$ .
- ⚙ Define an inner product  $\langle \cdot | \cdot \rangle$  on  $V_j$  as

$$\langle \mathbf{j}_m | \mathbf{j}_n \rangle = \int \mathbf{j}_m(\mathbf{r})^\dagger \mathbf{j}_n(\mathbf{r}) d\mathbf{r}.$$

↔ Enables projection-based unitary representation analysis on  $V_j$

# Current density linear space

⚙ The current densities  $\mathbf{j}(\mathbf{r})$  with  $\mathbf{r} \in \mathbb{R}^3$  form a linear space  $V_j$ .

⚙ Define an inner product  $\langle \cdot | \cdot \rangle$  on  $V_j$  as

$$\langle \mathbf{j}_m | \mathbf{j}_n \rangle = \int \mathbf{j}_m(\mathbf{r})^\dagger \mathbf{j}_n(\mathbf{r}) d\mathbf{r}.$$

↔ Enables projection-based unitary representation analysis on  $V_j$

⚙ Given a current density  $\mathbf{j}(\mathbf{r})$  and a symmetry or pseudo-symmetry **unitary** group  $\mathcal{G}$ , the required overlap matrix elements for the symmetry analysis of  $\mathbf{j}(\mathbf{r})$  are of the form

$$T_{m1}(e) = \langle \hat{u}_m \mathbf{j} | \mathbf{j} \rangle = \int (\hat{u}_m \mathbf{j})(\mathbf{r})^\dagger \mathbf{j}(\mathbf{r}) d\mathbf{r}, \quad u_m \in \mathcal{G}.$$

# Current density linear space

⚙ The current densities  $\mathbf{j}(\mathbf{r})$  with  $\mathbf{r} \in \mathbb{R}^3$  form a linear space  $V_j$ .

⚙ Define an inner product  $\langle \cdot | \cdot \rangle$  on  $V_j$  as

$$\langle \mathbf{j}_m | \mathbf{j}_n \rangle = \int \mathbf{j}_m(\mathbf{r})^\dagger \mathbf{j}_n(\mathbf{r}) d\mathbf{r}.$$

↔ Enables projection-based unitary representation analysis on  $V_j$

⚙ Given a current density  $\mathbf{j}(\mathbf{r})$  and a symmetry or pseudo-symmetry **unitary** group  $\mathcal{G}$ , the required overlap matrix elements for the symmetry analysis of  $\mathbf{j}(\mathbf{r})$  are of the form

(pseudo-)symmetry-transformed current density

$$T_{m1}(e) = \langle \mathbf{u}_m \mathbf{j} | \mathbf{j} \rangle = \int (\mathbf{u}_m \mathbf{j})(\mathbf{r})^\dagger \mathbf{j}(\mathbf{r}) d\mathbf{r}, \quad \mathbf{u}_m \in \mathcal{G}.$$

# Non-perturbative current densities

- ⚙ Non-perturbative calculations in arbitrarily strong magnetic fields are performed in a basis of **London atomic orbitals**:

$$\omega_{\mu}(\mathbf{r}; \mathbf{R}_{\mu}) = \varphi_{\mu}(\mathbf{r}; \mathbf{R}_{\mu}) \exp[-i\mathbf{A}(\mathbf{R}_{\mu}) \cdot \mathbf{r}].$$

- ⚙ In this basis, the current density can be partitioned into the **diamagnetic** and **paramagnetic** contributions with the *non-perturbative* forms:

$$\mathbf{j}(\mathbf{r}) = \mathbf{j}_d(\mathbf{r}) + \mathbf{j}_p(\mathbf{r})$$

$$\mathbf{j}_d(\mathbf{r}) = -\mathbf{A}(\mathbf{r}) \sum_{\sigma} \omega_{\mu}^{*}(\mathbf{r}) \omega_{\nu}(\mathbf{r}) P_{\sigma}^{\nu\mu}, \quad \mathbf{j}_p(\mathbf{r}) = \frac{i}{2} \sum_{\sigma} \nabla \omega_{\mu}^{*}(\mathbf{r}) \omega_{\nu}(\mathbf{r}) P_{\sigma}^{\nu\mu} + \text{c.c.}$$

Soncini, A. & Fowler, P. W. *Chem. Phys. Lett.* **396**, 174–181 (2004).

Tellgren, E. I. *et al. J. Chem. Phys.* **129**, 154114 (October 2008), Tellgren, E. I. *et al. J. Chem. Phys.* **140**, 034101 (January 2014).

Irons, T. J. P. *et al. Chemistry (MDPI)*. **3**, 916–934 (August 2021).

# Non-perturbative current densities

- Non-perturbative calculations in arbitrarily strong magnetic fields are performed in a basis of **London atomic orbitals**:

$$\omega_{\mu}(\mathbf{r}; \mathbf{R}_{\mu}) = \varphi_{\mu}(\mathbf{r}; \mathbf{R}_{\mu}) \exp[-i\mathbf{A}(\mathbf{R}_{\mu}) \cdot \mathbf{r}].$$

- In this basis, the current density can be partitioned into the **diamagnetic** and **paramagnetic** contributions with the *non-perturbative* forms:

$$\mathbf{j}(\mathbf{r}) = \mathbf{j}_d(\mathbf{r}) + \mathbf{j}_p(\mathbf{r})$$

$$\mathbf{j}_d(\mathbf{r}) = -\mathbf{A}(\mathbf{r}) \sum_{\sigma} \omega_{\mu}^{*}(\mathbf{r}) \omega_{\nu}(\mathbf{r}) P_{\sigma}^{\nu\mu}, \quad \mathbf{j}_p(\mathbf{r}) = \frac{i}{2} \sum_{\sigma} \nabla \omega_{\mu}^{*}(\mathbf{r}) \omega_{\nu}(\mathbf{r}) P_{\sigma}^{\nu\mu} + \text{c.c.}$$

- This partition depends on the gauge origin **G** which manifests itself in

$$\mathbf{A}(\mathbf{r}) = \frac{1}{2} \mathbf{B} \times (\mathbf{r} - \mathbf{G}).$$

Soncini, A. & Fowler, P. W. *Chem. Phys. Lett.* **396**, 174–181 (2004).

Tellgren, E. I. *et al. J. Chem. Phys.* **129**, 154114 (October 2008), Tellgren, E. I. *et al. J. Chem. Phys.* **140**, 034101 (January 2014).

Irons, T. J. P. *et al. Chemistry (MDPI)*. **3**, 916–934 (August 2021).

# Ipsocentric *DZ*

- ⚙ We employ the **ipsocentric *DZ*** method which makes use of a continuous set of gauge transformations:

$$\mathbf{G} \equiv \mathbf{G}(\mathbf{r}) = \mathbf{r},$$

so that  $\mathbf{j}_d(\mathbf{r})$  vanishes and  $\mathbf{j}(\mathbf{r}) = \mathbf{j}_p(\mathbf{r})$  formally.

---

Keith, T. A. & Bader, R. F. *Chem. Phys. Lett.* **210**, 223–231 (1993).

Soncini, A. & Fowler, P. W. *Chem. Phys. Lett.* **396**, 174–181 (2004).

# Ipsocentric $DZ$

- ⚙ We employ the **ipsocentric  $DZ$**  method which makes use of a continuous set of gauge transformations:

$$\mathbf{G} \equiv \mathbf{G}(\mathbf{r}) = \mathbf{r},$$

so that  $\mathbf{j}_d(\mathbf{r})$  vanishes and  $\mathbf{j}(\mathbf{r}) = \mathbf{j}_p(\mathbf{r})$  formally.

- ⚙ In addition,

$$\begin{aligned}\omega_\mu(\mathbf{r}; \mathbf{R}_\mu) &= \varphi_\mu(\mathbf{r}; \mathbf{R}_\mu) \exp\left[-\frac{i}{2}(\mathbf{B} \times (\mathbf{R}_\mu - \mathbf{r})) \cdot \mathbf{r}\right] \\ &= \varphi_\mu(\mathbf{r}; \mathbf{R}_\mu) \exp\left[-\frac{i}{2}(\mathbf{B} \times \mathbf{R}_\mu) \cdot \mathbf{r}\right],\end{aligned}$$

which is the same as keeping  $\mathbf{G}$  at the space-fixed origin.

Keith, T. A. & Bader, R. F. *Chem. Phys. Lett.* **210**, 223–231 (1993).

Soncini, A. & Fowler, P. W. *Chem. Phys. Lett.* **396**, 174–181 (2004).

# Integrals

- ⚙ The required overlap matrix elements for the symmetry analysis of  $\mathbf{j}(\mathbf{r})$  may now be cast in a computable form:

$$\begin{aligned}
 T_{m1}(e) &= \int (\hat{u}_m \mathbf{j}_p)(\mathbf{r})^\dagger \mathbf{j}_p(\mathbf{r}) d\mathbf{r} \\
 &= \frac{1}{4} \sum_{\sigma\sigma'} (P_\sigma^{v\mu})^* P_{\sigma'}^{v'\mu'} \\
 &\quad \int (\hat{u}_m \omega_v^*)(\mathbf{r}) \nabla (\hat{u}_m \omega_\mu)(\mathbf{r})^\dagger [\nabla \omega_{\mu'}^*(\mathbf{r}) \omega_{v'}(\mathbf{r}) - \nabla \omega_{\mu'}(\mathbf{r}) \omega_{v'}^*(\mathbf{r})] d\mathbf{r} + \text{c.c.}
 \end{aligned}$$



# Integrals

- ⚙ The required overlap matrix elements for the symmetry analysis of  $\mathbf{j}(\mathbf{r})$  may now be cast in a computable form:

$$\begin{aligned}
 T_{m1}(e) &= \int (\hat{u}_m \mathbf{j}_p)(\mathbf{r})^\dagger \mathbf{j}_p(\mathbf{r}) d\mathbf{r} \\
 &= \frac{1}{4} \sum_{\sigma\sigma'} (P_\sigma^{v\mu})^* P_{\sigma'}^{v'\mu'} \\
 &\quad \int (\hat{u}_m \omega_v^*)(\mathbf{r}) \nabla(\hat{u}_m \omega_\mu)(\mathbf{r})^\dagger [\nabla \omega_{\mu'}^*(\mathbf{r}) \omega_{v'}(\mathbf{r}) - \nabla \omega_{\mu'}(\mathbf{r}) \omega_{v'}^*(\mathbf{r})] d\mathbf{r} + \text{c.c.}
 \end{aligned}$$

second derivatives of four-centre overlap integrals

# Integrals

- ⚙ The required overlap matrix elements for the symmetry analysis of  $\mathbf{j}(\mathbf{r})$  may now be cast in a computable form:

$$\begin{aligned}
 T_{m1}(e) &= \int (\hat{u}_m \mathbf{j}_p)(\mathbf{r})^\dagger \mathbf{j}_p(\mathbf{r}) d\mathbf{r} \\
 &= \frac{1}{4} \sum_{\sigma\sigma'} (P_\sigma^{v\mu})^* P_{\sigma'}^{v'\mu'} \\
 &\quad \int (\hat{u}_m \omega_v^*)(\mathbf{r}) \nabla (\hat{u}_m \omega_\mu)(\mathbf{r})^\dagger [\nabla \omega_{\mu'}^*(\mathbf{r}) \omega_{v'}(\mathbf{r}) - \nabla \omega_{\mu'}(\mathbf{r}) \omega_{v'}^*(\mathbf{r})] d\mathbf{r} + \text{c.c.}
 \end{aligned}$$

(pseudo-)symmetry-transformed London orbitals