

Symmetry of Magnetically Induced Currents

MAGIC 2022 | *Peterhouse, Cambridge, UK*

Bang C. Huynh[†], Andrew M. Teale[†]

[†]School of Chemistry, University of Nottingham, UK

13th September 2022



MAGIC 2022
4th workshop on MAGnetically
Induced Currents in molecules

Introduction

Overview

- ⚙ Symmetry and pseudo-symmetry *groups* in magnetic fields
- ⚙ *Unitary* representation analysis on *linear spaces*
- ⚙ *Wavefunction* and *current density* symmetries
 - ⚙ *Relationships* between wavefunction and current density symmetries
 - ⚙ Symmetry *descent* and symmetry *breaking* in magnetic fields

Groups in magnetic fields

The electronic Hamiltonian

- ⚙ For an N_e -electron system in a **uniform** magnetic field $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$, consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\hat{\mathcal{H}} = \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k$$

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

The electronic Hamiltonian

- ⚙ For an N_e -electron system in a **uniform** magnetic field $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$, consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\hat{\mathcal{H}} = \boxed{\frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2} + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k$$

↑
kinetic

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

The electronic Hamiltonian

- ⚙ For an N_e -electron system in a **uniform** magnetic field $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$, consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\hat{\mathcal{H}} = \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \underbrace{\sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k)}_{\text{external potential}} + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k$$

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

The electronic Hamiltonian

- ⚙ For an N_e -electron system in a **uniform** magnetic field $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$, consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\hat{\mathcal{H}} = \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \underbrace{\frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|}}_{\text{electron-electron interaction}} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k$$

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

The electronic Hamiltonian

- ⚙ For an N_e -electron system in a **uniform** magnetic field $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$, consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\hat{\mathcal{H}} = \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \boxed{\frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k}$$

↑
spin Zeeman interaction

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

The electronic Hamiltonian

- ⚙ For an N_e -electron system in a **uniform** magnetic field $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$, consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\begin{aligned}\hat{\mathcal{H}} &= \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k \\ &= \frac{1}{2} \sum_{k=1}^{N_e} \hat{p}_k^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k)\end{aligned}$$

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

The electronic Hamiltonian

- ⚙ For an N_e -electron system in a **uniform** magnetic field $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$, consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\hat{\mathcal{H}} = \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k$$

$$= \underbrace{\frac{1}{2} \sum_{k=1}^{N_e} \hat{\mathbf{p}}_k^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|}}_{\substack{\text{Zero-field Hamiltonian, } \hat{\mathcal{H}}_0(v_{\text{ext}})}} + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k)$$

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

The electronic Hamiltonian

- ⚙ For an N_e -electron system in a **uniform** magnetic field $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$, consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\begin{aligned}\hat{\mathcal{H}} &= \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k \\ &= \frac{1}{2} \sum_{k=1}^{N_e} \hat{p}_k^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \underbrace{\mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k}_{\text{Linear}} + \underbrace{\frac{1}{2} A^2(\mathbf{r}_k)}_{\text{Quadratic}}\end{aligned}$$

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. *et al.* **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. *et al.* **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

The electronic Hamiltonian

- ⚙ For an N_e -electron system in a **uniform** magnetic field $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r})$, consider the **Schrödinger–Pauli Hamiltonian** in atomic units:

$$\begin{aligned}\hat{\mathcal{H}} &= \frac{1}{2} \sum_{k=1}^{N_e} |-\hat{\mathbf{p}}_k + \mathbf{A}(\mathbf{r}_k)|^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k \\ &= \frac{1}{2} \sum_{k=1}^{N_e} \hat{p}_k^2 + \sum_{k=1}^{N_e} v_{\text{ext}}(\mathbf{r}_k) + \frac{1}{2} \sum_{k \neq l}^{N_e} \frac{1}{|\mathbf{r}_k - \mathbf{r}_l|} + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k) \\ &= \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).\end{aligned}$$

- ⚙ How does each term transform **spatially** and **temporally**?

Weil, J. A. & Bolton, J. R. **Electron Paramagnetic Resonance**. 2nd (John Wiley & Sons, Inc., Hoboken, New Jersey, 2007).

Tellgren, E. I. et al. **J. Chem. Phys.** **148**, 024101 (January 2018).

Irons, T. J. P. et al. **Chemistry (MDPI)**. **3**, 916–934 (August 2021).

Tensor transformations

⚙ Let \mathbf{v} be a rank- k Cartesian tensor in three dimensions.

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al.* *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Tensor transformations

- Let \mathbf{v} be a rank- k Cartesian tensor in three dimensions.
- Spatially*, let $u \in O(3)$ be any **proper** or **improper rotation** that acts on an orthogonal basis $\{\mathbf{e}_i\}$ spanning \mathbb{R}^3 according to

$$\hat{u}\mathbf{e}_i = \mathbf{e}_j U_{ji}.$$

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al.* *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Tensor transformations

- Let \mathbf{v} be a rank- k Cartesian tensor in three dimensions.
- Spatially*, let $u \in O(3)$ be any **proper** or **improper rotation** that acts on an orthogonal basis $\{\mathbf{e}_i\}$ spanning \mathbb{R}^3 according to

$$\hat{u}\mathbf{e}_i = \mathbf{e}_j U_{ji}$$

U : representation matrix for u in $\{\mathbf{e}_i\}$

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al.* *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Tensor transformations

- Let \mathbf{v} be a rank- k Cartesian tensor in three dimensions.
- Spatially*, let $u \in O(3)$ be any **proper** or **improper rotation** that acts on an orthogonal basis $\{\mathbf{e}_i\}$ spanning \mathbb{R}^3 according to

$$\hat{u}\mathbf{e}_i = \mathbf{e}_j U_{ji}.$$

To consider the *linear* action of u on \mathbf{v} , let $\mathbf{v}' = \hat{u}\mathbf{v}$.

- \mathbf{v} is a **polar** tensor if

$$v'_{ab\dots k} = U_{ap}U_{bq}\dots U_{kz} v_{pq\dots z}.$$

- \mathbf{v} is an **axial** tensor if

$$v'_{ab\dots k} = |\mathbf{U}| U_{ap}U_{bq}\dots U_{kz} v_{pq\dots z}.$$

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al.* *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Tensor transformations

- Let \mathbf{v} be a rank- k Cartesian tensor in three dimensions.
- Spatially*, let $u \in O(3)$ be any **proper** or **improper rotation** that acts on an orthogonal basis $\{\mathbf{e}_i\}$ spanning \mathbb{R}^3 according to

$$\hat{u}\mathbf{e}_i = \mathbf{e}_j U_{ji}.$$

To consider the *linear* action of u on \mathbf{v} , let $\mathbf{v}' = \hat{u}\mathbf{v}$.

- \mathbf{v} is a **polar** tensor if

$$v'_{ab\dots k} = U_{ap} U_{bq} \dots U_{kz} v_{pq\dots z}.$$

- \mathbf{v} is an **axial** tensor if

$$v'_{ab\dots k} = |\mathbf{u}| U_{ap} U_{bq} \dots U_{kz} v_{pq\dots z}.$$

+1 for proper rotations, -1 for improper rotations

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al.* *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Tensor transformations

- ⚙ Let \mathbf{v} be a rank- k Cartesian tensor in three dimensions.
- ⚙ *Temporally*, let θ be the **time reversal** that acts on an orthogonal basis $\{\zeta_1, \zeta_2\}$ of a two-component spinor according to

$$\hat{\theta} \begin{bmatrix} \zeta_1 & \zeta_2 \end{bmatrix} = \begin{bmatrix} \zeta_2^* & -\zeta_1^* \end{bmatrix}.$$

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al. Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Tensor transformations

- Let \mathbf{v} be a rank- k Cartesian tensor in three dimensions.
- Temporally*, let θ be the **time reversal** that acts on an orthogonal basis $\{\zeta_1, \zeta_2\}$ of a two-component spinor according to

$$\hat{\theta}[\zeta_1 \quad \zeta_2] = [\zeta_2^* \quad -\zeta_1^*].$$

To consider the *antilinear* action of θ on \mathbf{v} , let $\mathbf{v}'' = \hat{\theta}\mathbf{v}$.

- \mathbf{v} is a **time-even** tensor (*i*-tensor) if

$$\mathbf{v}'' = \mathbf{v}.$$

- \mathbf{v} is a **time-odd** tensor (*c*-tensor) if

$$\mathbf{v}'' = -\mathbf{v}.$$

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Tensor classifications

⚙ There are *four* types of tensors under spatial-temporal transformations.

⚙ **polar time-even** tensors

⚙ position vectors \mathbf{r}

⚙ **axial time-even** tensors

⚙ **polar time-odd** tensors

⚙ linear momentum vectors \mathbf{p}

⚙ magnetic vector potentials \mathbf{A}

⚙ current densities \mathbf{j}

⚙ **axial time-odd** tensors

⚙ angular momentum vectors \mathbf{l}

⚙ magnetic field vectors \mathbf{B}

Birss, R. R. *Symmetry and Magnetism*. (North-Holland Pub. Co., Amsterdam, 1966).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. *et al.* *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Symmetry and pseudo-symmetry groups

Definition (symmetry group)

All transformations \hat{T} that leave the electronic Hamiltonian $\hat{\mathcal{H}}$ invariant, i.e. $\hat{T}\hat{\mathcal{H}}\hat{T}^{-1} = \hat{\mathcal{H}}$, form the **symmetry group** of $\hat{\mathcal{H}}$. Such transformations are called **symmetry transformations**.

- ⚙ Symmetry transformations impose **constraints** on the eigenfunctions of $\hat{\mathcal{H}}$ and properties calculated from them.

Symmetry and pseudo-symmetry groups

Definition (symmetry group)

All transformations \hat{T} that leave the electronic Hamiltonian $\hat{\mathcal{H}}$ invariant, i.e. $\hat{T}\hat{\mathcal{H}}\hat{T}^{-1} = \hat{\mathcal{H}}$, form the **symmetry group** of $\hat{\mathcal{H}}$. Such transformations are called **symmetry transformations**.

- ⚙ Symmetry transformations impose **constraints** on the eigenfunctions of $\hat{\mathcal{H}}$ and properties calculated from them.

Definition (pseudo-symmetry group)

Consider a term $\hat{\mathcal{H}}' \subset \hat{\mathcal{H}}$. All transformations \hat{T} that leave $\hat{\mathcal{H}}'$ invariant but not the full $\hat{\mathcal{H}}$ form a **pseudo-symmetry group** of $\hat{\mathcal{H}}$. Such transformations are called **pseudo-symmetry transformations**.

- ⚙ Pseudo-symmetry transformations provide ways to understand eigenfunctions of $\hat{\mathcal{H}}$ (**complicated**) from the perspective of $\hat{\mathcal{H}}'$ (**simpler**).

Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$

Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \underbrace{\hat{\mathcal{H}}_0(v_{\text{ext}})}_{\substack{\text{polar} \\ \text{time-even}}} + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$

Wigner, E. **Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra**. 386 (Academic Press, London, 1959).
 Bradley, C. J. & Davies, B. L. **Rev. Mod. Phys.** **40**, 359–379 (April 1968).
 Lazzeretti, P. et al. **Nucl. Magn. Shield. Mol. Struct.** (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).
 Keith, T. A. & Bader, R. F. J. **Chem. Phys.** **99**, 3669–3682 (1993).

Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \underbrace{\hat{\mathcal{H}}_0(v_{\text{ext}})}_{\mathcal{G} + \theta\mathcal{G}} + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$



Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \underbrace{\mathbf{A}(\mathbf{r}_k)}_{\substack{\text{polar} \\ \text{time-odd}}} \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$



Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$

↑
polar time-odd



Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \underbrace{\mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k}_{\mathcal{O}(3) + \theta\mathcal{O}(3)} + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \underbrace{\frac{1}{2}A^2(\mathbf{r}_k)}_{\mathcal{O}(3) + \theta\mathcal{O}(3)}.$$



Wigner, E. **Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra**. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. **Rev. Mod. Phys.** **40**, 359–379 (April 1968).

Lazzeretti, P. et al. **Nucl. Magn. Shield. Mol. Struct.** (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. **Chem. Phys.** **99**, 3669–3682 (1993).

Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \boxed{\frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k} + \frac{1}{2} A^2(\mathbf{r}_k).$$

↑
axial time-odd



Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).


Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \underbrace{\frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k}_{\substack{\mathcal{H} + \theta u \mathcal{H} \\ (u \in \mathcal{G}, \hat{u} \mathbf{B} = -\mathbf{B})}} + \frac{1}{2} A^2(\mathbf{r}_k).$$



Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).


Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. *J. Chem. Phys.* **99**, 3669–3682 (1993).

Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$



Wigner, E. *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra*. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. *Nucl. Magn. Shield. Mol. Struct.* (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).


Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

Groups in magnetic fields

- ⚙ Let us revisit the electronic Hamiltonian in a uniform magnetic field:

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0(v_{\text{ext}}) + \mathbf{A}(\mathbf{r}_k) \cdot \hat{\mathbf{p}}_k + \frac{g_s}{2} \sum_{k=1}^{N_e} \mathbf{B} \cdot \hat{\mathbf{s}}_k + \frac{1}{2} A^2(\mathbf{r}_k).$$

- ⚙ If $\mathcal{M} = \mathcal{H} + \theta u\mathcal{H}$ is isomorphic to a **unitary group** \mathcal{M}' , we write $\mathcal{M} = \mathcal{M}'(\mathcal{H})$.



Wigner, E. **Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra**. 386 (Academic Press, London, 1959).

Bradley, C. J. & Davies, B. L. *Rev. Mod. Phys.* **40**, 359–379 (April 1968).

Lazzeretti, P. et al. **Nucl. Magn. Shield. Mol. Struct.** (ed Tossell, J. A.) 163 (Springer Science+Business Media, B.V., Maryland, 1993).

Keith, T. A. & Bader, R. F. J. *Chem. Phys.* **99**, 3669–3682 (1993).

Unitary representation analysis

Group unitary representations on linear spaces

- ⚙ Consider a **group** \mathcal{G} that acts **unitarily** on a **linear space** V .
- ⚙ Let \mathbf{v} be an element in V . The unitary action of \mathcal{G} on \mathbf{v} generates an **orbit**

$$\mathcal{G} \cdot \mathbf{v} = \{\hat{u}_i \mathbf{v} \mid u_i \in \mathcal{G}\}$$

which spans a **representation subspace** $\Gamma \subseteq V$.

For simplicity, we will assume that $\mathcal{G} \cdot \mathbf{v}$ is a linearly independent basis.

Group unitary representations on linear spaces

- ⚙ Consider a **group** \mathcal{G} that acts **unitarily** on a **linear space** V .
- ⚙ Let \mathbf{v} be an element in V . The unitary action of \mathcal{G} on \mathbf{v} generates an **orbit**

$$\mathcal{G} \cdot \mathbf{v} = \{\hat{u}_i \mathbf{v} \mid u_i \in \mathcal{G}\}$$

which spans a **representation subspace** $\Gamma \subseteq V$.

For simplicity, we will assume that $\mathcal{G} \cdot \mathbf{v}$ is a linearly independent basis.

- ⚙ We seek to decompose Γ into known **irreducible** representations of \mathcal{G} on V .
 \hookrightarrow This **quantifies** the **symmetry** of \mathbf{v} under the action of \mathcal{G} .
- ⚙ To this end, we pick a reference element $\mathbf{v}_i = \hat{u}_i \mathbf{v}$ in $\mathcal{G} \cdot \mathbf{v}$ and define the **representation matrices** $D^\Gamma(u_k)$ for all $u_k \in \mathcal{G}$:

$$\hat{u}_k \mathbf{v}_i = \sum_j \mathbf{v}_j D_{ji}^\Gamma(u_k).$$

Their traces, $\chi^\Gamma(u_k)$, give the **characters** required for the decomposition of Γ .

Representation matrix determination

- ⚙ We opt for a projection method.
 - ↔ This requires V to be *endowed* with an *inner product* $\langle \cdot | \cdot \rangle$.

Representation matrix determination

- ⚙ We opt for a projection method.
 ↪ This requires V to be *endowed* with an **inner product** $\langle \cdot | \cdot \rangle$.
- ⚙ Define a non-orthogonal projection operator

$$\hat{\mathcal{P}}_I = \sum_j |\mathbf{v}_i\rangle (\mathbf{S}^{-1})_{ij} \langle \mathbf{v}_j|$$

where $S_{ij} = \langle \mathbf{v}_i | \mathbf{v}_j \rangle$ such that $\hat{\mathcal{P}}_I |\mathbf{v}_j\rangle = \delta_{ij} |\mathbf{v}_i\rangle$.

Representation matrix determination

- ⚙ We opt for a projection method.
 ↪ This requires V to be *endowed* with an *inner product* $\langle \cdot | \cdot \rangle$.
- ⚙ Define a non-orthogonal projection operator

$$\hat{\mathcal{P}}_i = \sum_j |\mathbf{v}_i\rangle (\mathbf{S}^{-1})_{ij} \langle \mathbf{v}_j|$$

where $S_{ij} = \langle \mathbf{v}_i | \mathbf{v}_j \rangle$ such that $\hat{\mathcal{P}}_i |\mathbf{v}_j\rangle = \delta_{ij} |\mathbf{v}_i\rangle$.

- ⚙ Application of $\hat{\mathcal{P}}$ on the defining equation for $\mathbf{D}^\Gamma(u_k)$ yields

$$\mathbf{D}^\Gamma(u_k) = \mathbf{S}^{-1} \mathbf{T}(u_k),$$

where

$$T_{ij}(u_k) = \langle \mathbf{v}_i | \hat{u}_k \mathbf{v}_j \rangle.$$

Representation matrix determination

- ⚙ We opt for a projection method.
 \hookrightarrow This requires V to be **endowed** with an **inner product** $\langle \cdot | \cdot \rangle$.
- ⚙ Define a non-orthogonal projection operator

$$\hat{\mathcal{P}}_i = \sum_j |\mathbf{v}_i\rangle (\mathbf{S}^{-1})_{ij} \langle \mathbf{v}_j|$$

where $S_{ij} = \langle \mathbf{v}_i | \mathbf{v}_j \rangle$ such that $\hat{\mathcal{P}}_i |\mathbf{v}_j\rangle = \delta_{ij} |\mathbf{v}_i\rangle$.

- ⚙ Application of $\hat{\mathcal{P}}$ on the defining equation for $\mathbf{D}^\Gamma(u_k)$ yields

$$\mathbf{D}^\Gamma(u_k) = \mathbf{S}^{-1} \mathbf{T}(u_k),$$

where

$$|\mathcal{G}|^3 \text{ elements} \rightarrow T_{ij}(u_k) = \langle \mathbf{v}_i | \hat{u}_k \mathbf{v}_j \rangle.$$

- ⚙ **Closure** of $\mathcal{G} \implies T_{ij}(u_k)$ can be **mapped to** $T_{m1}(\mathbf{e})$ for some $m = 1, \dots, |\mathcal{G}|$.
 $\hookrightarrow \mathbf{T}(u_k)$ can be computed with $\mathcal{O}(|\mathcal{G}|)$ time complexity.

Current density linear space

- ⚙ The current densities $\mathbf{j}(\mathbf{r})$ with $\mathbf{r} \in \mathbb{R}^3$ form a linear space V_j .
- ⚙ Define an inner product $\langle \cdot | \cdot \rangle$ on V_j as

$$\langle \mathbf{j}_m | \mathbf{j}_n \rangle = \int \mathbf{j}_m(\mathbf{r})^\dagger \mathbf{j}_n(\mathbf{r}) d\mathbf{r}.$$

↔ Enables projection-based unitary representation analysis on V_j

Current density linear space

- ⚙ The current densities $\mathbf{j}(\mathbf{r})$ with $\mathbf{r} \in \mathbb{R}^3$ form a linear space V_j .
- ⚙ Define an inner product $\langle \cdot | \cdot \rangle$ on V_j as

$$\langle \mathbf{j}_m | \mathbf{j}_n \rangle = \int \mathbf{j}_m(\mathbf{r})^\dagger \mathbf{j}_n(\mathbf{r}) d\mathbf{r}.$$

↔ Enables projection-based unitary representation analysis on V_j

- ⚙ Given a current density $\mathbf{j}(\mathbf{r})$ and a symmetry or pseudo-symmetry **unitary** group \mathcal{G} , the required overlap matrix elements for the symmetry analysis of $\mathbf{j}(\mathbf{r})$ are of the form

$$T_{m1}(e) = \langle \hat{u}_m \mathbf{j} | \mathbf{j} \rangle = \int (\hat{u}_m \mathbf{j})(\mathbf{r})^\dagger \mathbf{j}(\mathbf{r}) d\mathbf{r}, \quad u_m \in \mathcal{G}.$$

Current density linear space

- ⚙ The current densities $\mathbf{j}(\mathbf{r})$ with $\mathbf{r} \in \mathbb{R}^3$ form a linear space V_j .
- ⚙ Define an inner product $\langle \cdot | \cdot \rangle$ on V_j as

$$\langle \mathbf{j}_m | \mathbf{j}_n \rangle = \int \mathbf{j}_m(\mathbf{r})^\dagger \mathbf{j}_n(\mathbf{r}) d\mathbf{r}.$$

↔ Enables projection-based unitary representation analysis on V_j

- ⚙ Given a current density $\mathbf{j}(\mathbf{r})$ and a symmetry or pseudo-symmetry **unitary** group \mathcal{G} , the required overlap matrix elements for the symmetry analysis of $\mathbf{j}(\mathbf{r})$ are of the form

$$T_{m1}(e) = \langle \hat{u}_m \mathbf{j} | \mathbf{j} \rangle = \int (\hat{u}_m \mathbf{j})(\mathbf{r})^\dagger \mathbf{j}(\mathbf{r}) d\mathbf{r}, \quad u_m \in \mathcal{G}.$$

(pseudo-)symmetry-transformed current density

Non-perturbative current densities

- Non-perturbative calculations in arbitrarily strong magnetic fields are performed in a basis of **London atomic orbitals**:

$$\omega_{\mu}(\mathbf{r}; \mathbf{R}_{\mu}) = \varphi_{\mu}(\mathbf{r}; \mathbf{R}_{\mu}) \exp[-i\mathbf{A}(\mathbf{R}_{\mu}) \cdot \mathbf{r}].$$

- In this basis, the current density can be partitioned into the **diamagnetic** and **paramagnetic** contributions with the *non-perturbative* forms:

$$\mathbf{j}(\mathbf{r}) = \mathbf{j}_d(\mathbf{r}) + \mathbf{j}_p(\mathbf{r})$$

$$\mathbf{j}_d(\mathbf{r}) = -\mathbf{A}(\mathbf{r}) \sum_{\sigma} \omega_{\mu}^{*}(\mathbf{r}) \omega_{\nu}(\mathbf{r}) P_{\sigma}^{V\mu}, \quad \mathbf{j}_p(\mathbf{r}) = \frac{i}{2} \sum_{\sigma} \nabla \omega_{\mu}^{*}(\mathbf{r}) \omega_{\nu}(\mathbf{r}) P_{\sigma}^{V\mu} + \text{c.c.}$$

Soncini, A. & Fowler, P. W. *Chem. Phys. Lett.* **396**, 174–181 (2004).

Tellgren, E. I. *et al. J. Chem. Phys.* **129**, 154114 (October 2008), Tellgren, E. I. *et al. J. Chem. Phys.* **140**, 034101 (January 2014).

Irons, T. J. P. *et al. Chemistry (MDPI)*. **3**, 916–934 (August 2021).

Non-perturbative current densities

- Non-perturbative calculations in arbitrarily strong magnetic fields are performed in a basis of **London atomic orbitals**:

$$\omega_{\mu}(\mathbf{r}; \mathbf{R}_{\mu}) = \varphi_{\mu}(\mathbf{r}; \mathbf{R}_{\mu}) \exp[-i\mathbf{A}(\mathbf{R}_{\mu}) \cdot \mathbf{r}].$$

- In this basis, the current density can be partitioned into the **diamagnetic** and **paramagnetic** contributions with the *non-perturbative* forms:

$$\mathbf{j}(\mathbf{r}) = \mathbf{j}_d(\mathbf{r}) + \mathbf{j}_p(\mathbf{r})$$

$$\mathbf{j}_d(\mathbf{r}) = -\mathbf{A}(\mathbf{r}) \sum_{\sigma} \omega_{\mu}^{*}(\mathbf{r}) \omega_{\nu}(\mathbf{r}) P_{\sigma}^{\nu\mu}, \quad \mathbf{j}_p(\mathbf{r}) = \frac{i}{2} \sum_{\sigma} \nabla \omega_{\mu}^{*}(\mathbf{r}) \omega_{\nu}(\mathbf{r}) P_{\sigma}^{\nu\mu} + \text{c.c.}$$

- This partition depends on the gauge origin **G** which manifests itself in

$$\mathbf{A}(\mathbf{r}) = \frac{1}{2} \mathbf{B} \times (\mathbf{r} - \mathbf{G}).$$

Soncini, A. & Fowler, P. W. *Chem. Phys. Lett.* **396**, 174–181 (2004).

Tellgren, E. I. *et al. J. Chem. Phys.* **129**, 154114 (October 2008), Tellgren, E. I. *et al. J. Chem. Phys.* **140**, 034101 (January 2014).

Irons, T. J. P. *et al. Chemistry (MDPI)*. **3**, 916–934 (August 2021).

Ipsocentric *DZ*

- ⚙ We employ the **ipsocentric *DZ*** method which makes use of a continuous set of gauge transformations:

$$\mathbf{G} \equiv \mathbf{G}(\mathbf{r}) = \mathbf{r},$$

so that $\mathbf{j}_d(\mathbf{r})$ vanishes and $\mathbf{j}(\mathbf{r}) = \mathbf{j}_p(\mathbf{r})$ formally.

Keith, T. A. & Bader, R. F. *Chem. Phys. Lett.* **210**, 223–231 (1993).

Soncini, A. & Fowler, P. W. *Chem. Phys. Lett.* **396**, 174–181 (2004).

Ipsocentric DZ

- ⚙ We employ the **ipsocentric DZ** method which makes use of a continuous set of gauge transformations:

$$\mathbf{G} \equiv \mathbf{G}(\mathbf{r}) = \mathbf{r},$$

so that $\mathbf{j}_d(\mathbf{r})$ vanishes and $\mathbf{j}(\mathbf{r}) = \mathbf{j}_p(\mathbf{r})$ formally.

- ⚙ In addition,

$$\begin{aligned}\omega_\mu(\mathbf{r}; \mathbf{R}_\mu) &= \varphi_\mu(\mathbf{r}; \mathbf{R}_\mu) \exp\left[-\frac{i}{2}(\mathbf{B} \times (\mathbf{R}_\mu - \mathbf{r})) \cdot \mathbf{r}\right] \\ &= \varphi_\mu(\mathbf{r}; \mathbf{R}_\mu) \exp\left[-\frac{i}{2}(\mathbf{B} \times \mathbf{R}_\mu) \cdot \mathbf{r}\right],\end{aligned}$$

which is the same as keeping \mathbf{G} at the space-fixed origin.

Keith, T. A. & Bader, R. F. *Chem. Phys. Lett.* **210**, 223–231 (1993).

Soncini, A. & Fowler, P. W. *Chem. Phys. Lett.* **396**, 174–181 (2004).

Integrals

- ⚙ The required overlap matrix elements for the symmetry analysis of $\mathbf{j}(\mathbf{r})$ may now be cast in a computable form:

$$\begin{aligned}
 T_{m1}(e) &= \int (\hat{u}_m \mathbf{j}_p)(\mathbf{r})^\dagger \mathbf{j}_p(\mathbf{r}) d\mathbf{r} \\
 &= \frac{1}{4} \sum_{\sigma\sigma'} (P_{\sigma}^{v\mu})^* P_{\sigma'}^{v'\mu'} \\
 &\quad \int (\hat{u}_m \omega_v^*)(\mathbf{r}) \nabla(\hat{u}_m \omega_\mu)(\mathbf{r})^\dagger [\nabla \omega_{\mu'}^*(\mathbf{r}) \omega_{v'}(\mathbf{r}) - \nabla \omega_{\mu'}(\mathbf{r}) \omega_{v'}^*(\mathbf{r})] d\mathbf{r} + \text{c.c.}
 \end{aligned}$$

Integrals

- ⚙ The required overlap matrix elements for the symmetry analysis of $\mathbf{j}(\mathbf{r})$ may now be cast in a computable form:

$$\begin{aligned}
 T_{m1}(e) &= \int (\hat{u}_m \mathbf{j}_p)(\mathbf{r})^\dagger \mathbf{j}_p(\mathbf{r}) d\mathbf{r} \\
 &= \frac{1}{4} \sum_{\sigma\sigma'} (P_{\sigma}^{\nu\mu})^* P_{\sigma'}^{\nu'\mu'} \\
 &\quad \int (\hat{u}_m \omega_{\nu}^*)(\mathbf{r}) \nabla (\hat{u}_m \omega_{\mu})(\mathbf{r})^T [\nabla \omega_{\mu'}^*(\mathbf{r}) \omega_{\nu'}(\mathbf{r}) - \nabla \omega_{\mu'}(\mathbf{r}) \omega_{\nu'}^*(\mathbf{r})] d\mathbf{r} + \text{c.c.}
 \end{aligned}$$

second derivatives of four-centre overlap integrals

Integrals

- ⚙ The required overlap matrix elements for the symmetry analysis of $\mathbf{j}(\mathbf{r})$ may now be cast in a computable form:

$$\begin{aligned}
 T_{m1}(e) &= \int (\hat{u}_m \mathbf{j}_p)(\mathbf{r})^\dagger \mathbf{j}_p(\mathbf{r}) d\mathbf{r} \\
 &= \frac{1}{4} \sum_{\sigma\sigma'} (P_\sigma^{v\mu})^* P_{\sigma'}^{v'\mu'} \\
 &\quad \int (\hat{u}_m \omega_v^*)(\mathbf{r}) \nabla (\hat{u}_m \omega_\mu)(\mathbf{r})^\dagger [\nabla \omega_{\mu'}^*(\mathbf{r}) \omega_{v'}(\mathbf{r}) - \nabla \omega_{\mu'}(\mathbf{r}) \omega_{v'}^*(\mathbf{r})] d\mathbf{r} + \text{c.c.}
 \end{aligned}$$

(pseudo-)symmetry-transformed London orbitals

Current density symmetry

UHF electronic structure