

1 Intro

1.1 Template

2 Data Structures

2.1 Template

3 Dynamic Programming

3.1 Template

4 Graphs

4.1 Template

5 Strings

5.1 Template

6 Mathematics

6.1 Template

7 Geometry

7.1 Template

8 Miscellaneous

8.1 Template

9 Math Extra

9.1 Seldom used combinatorics

- Fibonacci in $O(\log(N))$ with memoization is:

$$f(0) = f(1) = 1$$

$$f(2k) = f(k)^2 + f(k-1)^2$$

$$f(2k+1) = f(k) \times (f(k) + 2 * f(k-1))$$

- Wilson's Theorem Extension:

$$B = b_1 b_2 \dots b_m \pmod n = \pm 1, \text{ all } b_i \leq n \text{ such that } \gcd(b_i, n) = 1.$$

If $n \leq 4$ or $n = (\text{odd prime})^k$ or $n = 2(\text{odd prime})^k$ then $B = -1$ for any k .

Else $B = 1$.

- Stirling numbers of the second kind, denoted by $S(n, k)$ = number of ways to split n numbers into k non-empty sets.

$$S(n, 1) = S(n, n) = 1$$

$$S(n, k) = kS(n-1, k) + S(n-1, k-1)$$

$S(n-d+1, k-d+1) = S(n, k)$ where if indexes i, j belong to the same set, then $|i-j| \geq d$.

- Burnside's Lemma: $|\text{classes}| = \frac{1}{|G|} \sum_{\forall g \in G} (K^{C(g)})$, where:
 G = different permutations possible,

$C(g)$ = number of cycles on the permutation g ,

and K = Number of states for each element

- Different ways to paint a necklace with N beads and K colors:

$$G = (1, 2, \dots, N), (2, 3, \dots, N, 1), \dots, (N, 1, \dots, N-1)$$

$g_i = (i, i+1, \dots, i+N)$, (taking mod N to get it right) $i = 1 \dots N$
 with i per step, that is, $i \rightarrow 2i \rightarrow 3i \dots$

Cycles in g_i all have size $n/\text{gcd}(i, n)$, so

$$C(g_i) = \text{gcd}(i, n)$$

$$\text{ans} = \frac{1}{N} \sum_{i=1 \dots N} (K^{\text{gcd}(i, n)})$$

$$\text{ans} = \frac{1}{N} \sum_{\forall d|N} (\phi(\frac{N}{d}) K^d)$$

9.2 Combinatorial formulas

$$\sum_{k=0}^n k^2 = n(n+1)(2n+1)/6$$

$$\sum_{k=0}^n k^3 = n^2(n+1)^2/4$$

$$\sum_{k=0}^n k^4 = (6n^5 + 15n^4 + 10n^3 - n)/30$$

$$\sum_{k=0}^n k^5 = (2n^6 + 6n^5 + 5n^4 - n^2)/12$$

$$\sum_{k=0}^n x^k = (x^{n+1} - 1)/(x - 1)$$

$$\sum_{k=0}^n kx^k = (x - (n+1)x^{n+1} + nx^{n+2})/(x-1)^2$$

$$\binom{n}{k} = \frac{n!}{(n-k)!k!}$$

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}$$

$$\binom{n}{k} = \frac{n}{n-k} \binom{n-1}{k}$$

$$\binom{n}{k} = \frac{n-k+1}{k} \binom{n}{k-1}$$

$$\binom{n+1}{k} = \frac{n+1}{n-k+1} \binom{n}{k}$$

$$\binom{n}{k+1} = \frac{n-k}{k+1} \binom{n}{k}$$

$$\sum_{k=1}^n k \binom{n}{k} = n2^{n-1}$$

$$\sum_{k=1}^n k^2 \binom{n}{k} = (n+n^2)2^{n-2}$$

$$\binom{m+n}{r} = \sum_{k=0}^r \binom{m}{k} \binom{n}{r-k}$$

$$\binom{n}{k} = \prod_{i=1}^k \frac{n-k+i}{i}$$

9.3 Number theory identities

Lucas' Theorem: For non-negative integers m and n and a prime p ,

$$\binom{m}{n} \equiv \prod_{i=0}^k \binom{m_i}{n_i} \pmod{p},$$

where

$$m = m_k p^k + m_{k-1} p^{k-1} + \cdots + m_1 p + m_0$$

is the base p representation of m , and similarly for n .

9.4 Stirling Numbers of the second kind

Number of ways to partition a set of n numbers into k non-empty subsets.

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\} = \frac{1}{k!} \sum_{j=0}^k (-1)^{(k-j)} \binom{k}{j} j^n$$

Recurrence relation:

$$\left\{ \begin{matrix} 0 \\ 0 \end{matrix} \right\} = 1$$

$$\left\{ \begin{matrix} n \\ 0 \end{matrix} \right\} = \left\{ \begin{matrix} 0 \\ n \end{matrix} \right\} = 1$$

$$\left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\} = k \left\{ \begin{matrix} n \\ k \end{matrix} \right\} + \left\{ \begin{matrix} n \\ k-1 \end{matrix} \right\}$$

9.5 Burnside's Lemma

Let G be a finite group that acts on a set X . For each g in G let X^g denote the set of elements in X that are fixed by g , which means $X^g = \{x \in X | g(x) = x\}$. Burnside's lemma asserts the following formula for the number of orbits, denoted $|X/G|$:

$$|X/G| = \frac{1}{|G|} \sum_{g \in G} |X^g|$$

9.6 Numerical integration

RK4: to integrate $\dot{y} = f(t, y)$ with $y_0 = y(t_0)$, compute

$$k_1 = f(t_n, y_n)$$

$$k_2 = f(t_n + \frac{h}{2}, y_n + \frac{h}{2} k_1)$$

$$k_3 = f(t_n + \frac{h}{2}, y_n + \frac{h}{2} k_2)$$

$$k_4 = f(t_n + h, y_n + h k_3)$$

$$y_{n+1} = y_n + \frac{h}{6} (k_1 + 2k_2 + 2k_3 + k_4)$$