



McCOMBS SCHOOL OF BUSINESS

Salem Center for Policy

Probability

David Puelz

September 13, 2021



The basics and conditional probability

Independence

Paradoxes, mixtures, and the rule of total probability

Random variables, distributions, and simulation

What is probability?



- A measure of **uncertainty**
- Answering the question: “How likely is a given event?”
- As with any mathematical concept, there are a set of **axioms** setting the “ground rules”
- Separately, there are different ways to interpret probability ...
 - (i) **frequentist**: limit of relative frequency after repeating an experiment an infinite number of times (coin flip!)
 - (ii) **Bayesian**: subjective belief about the likelihood of an event occurrence



If A denotes some event, then $P(A)$ is the probability that this event occurs:

- $P(\text{coin lands heads}) = 0.5$
- $P(\text{rainy day in Ireland}) = 0.85$
- $P(\text{cold day in Hell}) = 0.0000001$

And so on...



If A denotes some event, then $P(A)$ is the probability that this event occurs:

- $P(\text{coin lands heads}) = 0.5$
- $P(\text{rainy day in Ireland}) = 0.85$
- $P(\text{cold day in Hell}) = 0.0000001$

And so on...



Some probabilities are estimated from direct experience over the long run:

- $P(\text{newborn baby is a boy}) = \frac{106}{206}$
- $P(\text{death due to car accident}) = \frac{11}{100,000}$
- $P(\text{death due to any cause}) = 1$

Some probabilities are estimated from direct experience over the long run:

- $P(\text{newborn baby is a boy}) = \frac{106}{206}$
- $P(\text{death due to car accident}) = \frac{11}{100,000}$
- $P(\text{death due to any cause}) = 1$

Others are synthesized from our best judgments about unique events:

- $P(\text{Apple stock goes up after next earnings call}) = 0.54$
- $P(\text{Djokovic wins next US Open}) = 0.4$ (6 to 4 odds)
- etc.



A conditional probability is the chance that one thing happens, given that some other thing has already happened.

A great example is a weather forecast: if you look outside this morning and see gathering clouds, you might assume that rain is likely and carry an umbrella.

We express this judgment as a conditional probability: e.g. “the conditional probability of rain this afternoon, given clouds this morning, is 60%.”



In statistics, we write this a bit more compactly:

- $P(\text{rain this afternoon} \mid \text{clouds this morning}) = 0.6$
- That vertical bar means “given” or “conditional upon.”
- The thing on the left of the bar is the event we’re interested in.
- The thing on the right of the bar is our knowledge, also called the “conditioning event” or “conditioning variable”: what we believe or assume to be true.

$P(A \mid B)$: “the probability of A, given that B occurs.”



Conditional probabilities are how we express judgments in a way that reflects our partial knowledge.

- You just gave *Manifest* a high rating. What's the conditional probability that you will like *The Bourne Identity* or *Saving Private Ryan*?



Conditional probabilities are how we express judgments in a way that reflects our partial knowledge.

- You just gave *Manifest* a high rating. What's the conditional probability that you will like *The Bourne Identity* or *Saving Private Ryan*?
- You just bought organic dog food on Amazon. What's the conditional probability that you will also buy a GPS-enabled dog collar?



Conditional probabilities are how we express judgments in a way that reflects our partial knowledge.

- You just gave *Manifest* a high rating. What's the conditional probability that you will like *The Bourne Identity* or *Saving Private Ryan*?
- You just bought organic dog food on Amazon. What's the conditional probability that you will also buy a GPS-enabled dog collar?
- You follow Lionel Messi (@leomessi) on Instagram. What's the conditional probability that you will respond to a suggestion to follow Cristiano Ronaldo (@cristiano)?



Conditional probabilities are how we express judgments in a way that reflects our partial knowledge.

- You just gave *Manifest* a high rating. What's the conditional probability that you will like *The Bourne Identity* or *Saving Private Ryan*?
- You just bought organic dog food on Amazon. What's the conditional probability that you will also buy a GPS-enabled dog collar?
- You follow Lionel Messi (@leomessi) on Instagram. What's the conditional probability that you will respond to a suggestion to follow Cristiano Ronaldo (@cristiano)?



A really important fact is that conditional probabilities are **not symmetric**:

$$P(A \mid B) \neq P(B \mid A)$$

As a quick counter-example, let the events A and B be as follows:

- A: “you can dribble a basketball”
- B: “you play in the NBA”

Probability basics: conditioning



- A: “you can dribble a basketball”
- B: “you play in the NBA”



Clearly $P(A \mid B) = 1$: every NBA player can dribble a basketball!

Probability basics: conditioning



- A : “you can dribble a basketball”
- B : “you play in the NBA”



But $P(B \mid A)$ is nearly zero!



An **uncertain outcome** (more formally called a “random process”) has two key properties:

1. The set of possible outcomes, called the sample space, *is known* beforehand.
 2. The particular outcome that occurs is *not known* beforehand.
- We denote the **sample space** as Ω , and some particular element of the sample space as $\omega \in \Omega$

Uncertain outcomes and probability models



Examples:

1. NBA finals, Golden State vs. Toronto:

$$\Omega = \{4-0, 4-1, 4-2, 4-3, 3-4, 2-4, 1-4, 0-4\}$$

Uncertain outcomes and probability models



Examples:

1. NBA finals, Golden State vs. Toronto:

$$\Omega = \{4-0, 4-1, 4-2, 4-3, 3-4, 2-4, 1-4, 0-4\}$$

2. Temperature in degrees F in Austin on a random day:

$$\Omega = [10, 115]$$

Uncertain outcomes and probability models



Examples:

1. NBA finals, Golden State vs. Toronto:

$$\Omega = \{4-0, 4-1, 4-2, 4-3, 3-4, 2-4, 1-4, 0-4\}$$

2. Temperature in degrees F in Austin on a random day:

$$\Omega = [10, 115]$$

3. Number of no-shows on an AA flight from Austin to DFW:

$$\Omega = \{0, 1, 2, \dots, N_{\text{seats}}\}$$



Examples:

1. NBA finals, Golden State vs. Toronto:

$$\Omega = \{4-0, 4-1, 4-2, 4-3, 3-4, 2-4, 1-4, 0-4\}$$

2. Temperature in degrees F in Austin on a random day:

$$\Omega = [10, 115]$$

3. Number of no-shows on an AA flight from Austin to DFW:

$$\Omega = \{0, 1, 2, \dots, N_{\text{seats}}\}$$

4. Poker hand

$$\Omega = \text{all possible five-card deals from a 52-card deck}$$



An **event** is a *subset of the sample space*, i.e. $A \subset \Omega$. For example:

1. **NBA finals, Golden State vs. Toronto.** Let A be the event "Toronto wins". Then

$$A = \{3-4, 2-4, 1-4, 0-4\} \subset \Omega$$

2. **Austin weather.** Let A be the event "cooler than 90 degrees". Then

$$A = [10, 90) \subset [10, 115]$$

3. **Flight no-shows.** Let A be "more than 5 no shows":

$$A = \{6, 7, 8, \dots, N_{\text{seats}}\}$$



We need some basic set-theory concepts to make sense of probability, since the sample space Ω is a set, and since “events” are subsets of Ω .

Union: $A \cup B = \{\omega : \omega \in A \text{ or } \omega \in B\}$

Intersection: $A \cap B = \{\omega : \omega \in A \text{ and } \omega \in B\}$

Complement: $A^C = \tilde{A} = \{\omega : \omega \notin A\}$

Difference: $A \setminus B = \{\omega : \omega \in A, \omega \notin B\}$

Disjointness: A and B are disjoint if $A \cap B = \emptyset$ (the empty set).

Axioms of probability (Kolmogorov)



These are the **ground rules**!

Consider an uncertain outcome with sample space Ω . “Probability” $P(\cdot)$ is a set function that maps Ω to the real numbers, such that:

1. **Non-negativity**: For any event $A \subset \Omega$, $P(A) \geq 0$.
2. **Normalization**: $P(\Omega) = 1$ and $P(\emptyset) = 0$.
3. **Finite additivity**: If A and B are disjoint, then $P(A \cup B) = P(A) + P(B)$.
- 3a. **Finite additivity (general)**: For any sets A and B ,
$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

(bonus: prove this with set theory!)

Not that intuitive! Notice no mention of frequencies...



- **Uncertain outcome/“random process”**: we know the possibilities ahead of time, just not the specific one that occurs
- **Sample space**: the set of possible outcomes
- **Event**: a subset of the sample space
- **Probability**: a function that maps events to real numbers and that obeys Kolmogorov's axioms

OK, so how do we actually *calculate* probabilities?



Now that we have an understanding of the axioms, notation, and interpretation, how do we **calculate** probabilities?

Counting!



(review 6.1.3-6.1.4 in the QSS book ... ways to count objects in structured sets are discussed)

Suppose our sample space Ω is a finite set consisting of N elements $\omega_1, \dots, \omega_N$.

Suppose further that $P(\omega_i) = 1/N$: each outcome is equally likely, i.e. we have a discrete uniform distribution over possible outcomes.

Then for each set $A \subset \Omega$,


$$P(A) = \frac{|A|}{N} = \frac{\text{Number of elements in } A}{\text{Number of elements in } \Omega}$$

That is, to compute $P(A)$, we just need to count how many elements are in A .

Counting example



Someone deals you a five-card poker hand from a 52-card deck.
What is the probability of a flush (all five cards the same suit)?



Note: this is a very historically accurate illustration of probability, given its origins among bored French aristocrats!

Counting example



- Our sample space has $N = \binom{52}{5} = 2,598,960$ possible poker hands, each one equally likely.
- How many possible flushes are there? Let's start with hearts:
→ There are 13 hearts



- Our sample space has $N = \binom{52}{5} = 2,598,960$ possible poker hands, each one equally likely.
- How many possible flushes are there? Let's start with hearts:
 - There are 13 hearts
 - To make a flush with hearts, you need any 5 of these 13 cards.



- Our sample space has $N = \binom{52}{5} = 2,598,960$ possible poker hands, each one equally likely.
- How many possible flushes are there? Let's start with hearts:
 - There are 13 hearts
 - To make a flush with hearts, you need any 5 of these 13 cards.
 - Thus there are $\binom{13}{5} = 1287$ possible flushes with hearts.



- Our sample space has $N = \binom{52}{5} = 2,598,960$ possible poker hands, each one equally likely.
- How many possible flushes are there? Let's start with hearts:
 - There are 13 hearts
 - To make a flush with hearts, you need any 5 of these 13 cards.
 - Thus there are $\binom{13}{5} = 1287$ possible flushes with hearts.
 - The same argument works for all four suits, so there are $4 \times 1287 = 5,148$ flushes. Thus:



- Our sample space has $N = \binom{52}{5} = 2,598,960$ possible poker hands, each one equally likely.
- How many possible flushes are there? Let's start with hearts:
 - There are 13 hearts
 - To make a flush with hearts, you need any 5 of these 13 cards.
 - Thus there are $\binom{13}{5} = 1287$ possible flushes with hearts.
 - The same argument works for all four suits, so there are $4 \times 1287 = 5,148$ flushes. Thus:

$$P(\text{flush}) = \frac{|A|}{|\Omega|} = \frac{5148}{2598960} = 0.00198079$$